

Oenothera humifusa

seabeach evening primrose

Onagraceae



Credit: Josiah Raymer, UF/IFAS

Seabeach evening primrose is found in beach dunes throughout coastal counties in Florida, west to Louisiana, and as far north as New Jersey.

General Description

Seabeach evening primrose is an herbaceous perennial with horizontal stems forming a rounded, mounding canopy. *Leaves* are simple and alternate and wider than 1 mm with short, silky pubescence. *Stems* are prostrate, grow up to 3 ft long, and are covered with silky appressed pubescence. *Flowers* are axillary with 4 yellow petals that are 0.2 to 0.6 in long, 4 sepals that are 0.1 to 0.4 in long, and flattened bracts. They have a hypanthium that is longer than the ovary. *Fruits* are 10- to 35-mm-long dehiscent capsules with appressed hairs that give rise to numerous rounded seeds.

Propagation

No published seed propagation information is presently available for *O. humifusa*; however, other *Oenothera* species

are well studied and have been shown to require light to germinate (Greiner and Köhl 2014). The authors collected stem cuttings in the fall and found the seeds present within the capsules already on the cuttings readily germinated in propagation flats under intermittent mist and natural photoperiod post-dehiscence. Seedlings initially form a low, broad rosette and are easily transplanted as small plugs after two weeks' growth in the propagation flat. Seedling growth is eventually lateral becoming multibranched. Seedlings will fill 4-in pots in 4 weeks when grown outdoors with overhead irrigation and fertilized with slow-release fertilizer.



Credit: Mack Thetford, UF/IFAS

are well studied and have been shown to require light to germinate (Greiner and Köhl 2014). The authors collected stem cuttings in the fall and found the seeds present within the capsules already on the cuttings readily germinated in propagation flats under intermittent mist and natural photoperiod post-dehiscence. Seedlings initially form a low, broad rosette and are easily transplanted as small plugs after two weeks' growth in the propagation flat. Seedling growth is eventually lateral becoming multibranched. Seedlings will fill 4-in pots in 4 weeks when grown outdoors with overhead irrigation and fertilized with slow-release fertilizer.

The authors found stem cuttings of *Oenothera* rapidly and readily produce roots without auxin application. Apical stem cuttings taken in late April 2017 from Santa Rosa Island, Florida, rooted and filled 162-cell flats in less than 4 weeks while under intermittent mist and natural photoperiod in a controlled-environment greenhouse. Cuttings grown in 72-cell flats

filled 4-in pots in 4 weeks when grown outdoors under overhead irrigation and fertilized with slow-release fertilizer. In nursery culture, this plant may become infested with flea beetles (*Altica* sp). The preferred plant hosts of this insect are likely limited to the members of the evening primrose family, Onagraceae (Clark et al. 2004). The beetle is a problem on *Oenothera* in the spring (May–June in north Florida) when larvae and adults appear in large numbers and can feed on the host plant until it dies. The authors have not noted this insect to be a serious pest of *Oenothera* growing within coastal plant communities, likely related to

1. This document is SGEB-75-16, one of a series of the Florida Sea Grant College Program, UF/IFAS Extension. Original publication date June 2018. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Debbie Miller, professor, Wildlife Ecology and Conservation Department, UF/IFAS West Florida Research and Education Center; Mack Thetford, associate professor, Environmental Horticulture Department, UF/IFAS West Florida Research and Education Center; Christina Verlinde, UF/IFAS Extension Florida Sea Grant Agent, Santa Rosa County; Gabriel Campbell, graduate research assistant, UF/IFAS West Florida Research and Education Center; and Ashlynn Smith, graduate research assistant, UF/IFAS West Florida Research and Education Center, Milton, FL 32583.



the harsh environment and presence of natural enemies. Commercial pest management recommendations are available to manage the pest in nursery culture.

Outplanting



Credit: Steven Wagner, UF/IFAS

Plants in 4-in pots with fully developed rootballs and canopies have successfully been outplanted by the authors. In a preliminary trial, when planted in July on open beach areas behind low frontal dunes, transplant success was 43%. However, in subsequent plantings in early March survival was 100% after 2 months. Plants were outplanted on backdunes midslope at least 12 in from each other and existing perennial grasses. Fertilizer application ($\frac{1}{2}$ tsp Osmocote 18-6-12) improved the aesthetic appearance of plants and increased biomass for this plant but was not necessary for survival.

Literature Cited

- Clark, S.M., D.G. LeDoux, T.N. Seeno, E.G. Riley, A.J. Gilbert, and J.M. Sullivan. 2004. "Host plants of leaf beetle species occurring in the United States and Canada (*Coleoptera*: *Megalopodidae*, *Orsodacnidae*, *Chrysomelidae* exclusive of *Bruchinae*)."
Coleopterists Society, Special Publication no. 2. 476 pp.
- Greiner, S., and K. Köhl. 2014. "Growing evening primrose (*Oenothera*)."
Frontiers in Plant Science. 5(38):1–12.

