

Chapter 12. Onion, Leek, and Chive Production in Florida

Peter J. Dittmar, Eugene J. McAvoy, Monica Ozores-Hampton, Richard N. Raid, Pamela Roberts, Hugh A. Smith, Xavier Martini, Johan Desaegeer, Joseph W. Noling, Shouan Zhang, and Lincoln Zotarelli

Botany and Planting

Onion - *Allium cepa*, Alliaceae (Amaryllidaceae)

Leek - *Allium ampeloprasum*

Chive - *Allium schoenoprasum*

Bunching onion - *Allium fistulosum*

Table 12.1. Planting information for onion and allies.

Planting dates	Seeded	Transplanted
North Florida	Mid Sept - mid Nov	Nov - Jan
Central Florida	Oct	Dec - Jan
South Florida	Oct	Dec - Jan
Planting information		
Distance between rows (in)	14 - 18	14 - 18
Distance between plants (in)	3 - 4	4 - 6
Seeding depth (in)	0.25 - 0.5	-
Seed per acre (lb)	3 - 4	1
Days to maturity	100 - 130	100 - 1302
Plant populations (acre)	149,343	112,123

Cultivars

Table 12.2. Onion and leek cultivars.

Onion		
Dawn (H)	Prowler (H)	Sweet Harvest (H)
Granex 33 (H)	Red Hunter	Sweet Success (H)
Plethora (H)	Savannah Sweet (H)	Vidurra (H)
Pirate (H)	Sweet Caroline (H)	
Bunching Onion		
Perfecto Blanco	Tokyo Long White	White Portugal
Leeks		
Jolant	Chives	Tivi
King Richard	Staro	Tornado
Lancelot	Megathon	Verina
Chives		
Staro		
¹ H=hybrid		

Table 12.3. Herbicides approved for managing weeds in onion.

Labels change frequently. Be sure to read a current product label before applying any chemical.

Active ingredient lb. a.i./A	Trade name product/A	MOA code	Crops	Weeds controlled / remarks
*** PREEMERGENCE ***				
Bensulide 5.0 - 6.0	(Prefar) 4 E 5 - 6 qt.	8	Dry bulb vegetables: onion, garlic, shallot	Annual broadleaf and grass control. Incorporate or irrigate 1 to 2 in. deep within 36 hr. of application. Consult label for rotational restrictions.
Bromoxynil 0.25 - 0.38	(Buctril) 2 EC 1.0 - 1.5 pt. (Buctril) 4 EC 0.5 - 0.75 pt.	6	Onion (dry bulb)	Broadleaf weeds. Preemergence is restricted to muck soils containing greater than 10% organic matter. Apply at least 3 to 4 days prior to emergence. Weeds should not exceed the 4-leaf stage, 2 in. in height or 1 in. in diameter.
Carfentrazone up to 0.031	(Aim) 1.9 EW up to 2 fl. oz. (Aim) 2.0 EC up to 2 fl. oz.	14	Onion, garlic, leek, chive, shallot	Apply as a preplant burn down for emerged broadleaf weeds. Use crop oil concentrate, methylated seed oil, or nonionic surfactant at recommended rates. Maximum rate of 0.096 lb. a.i./A per season. No pretransplant interval.
DCPA 4.5 - 6.0	(Dacthal) W-75 6 - 8 lb. (Dacthal) 6 F 6 - 8 lb.	3	Onion (dry bulb, green), leek, shallot	Annual grasses and certain broadleaf control. Apply at seeding or planting. Maximum rate of 10.5 lb. a.i./A per season.
Flumioxazin up to 0.06	(Chateau) 51 WDG up to 2 fl. oz.	14	Garlic, Onion (dry bulb)	Broadleaf control. Apply to transplanted onions between the 2- and 6- leaf stage. Apply to direct seeded onions between the 3- and 6- leaf stage.

Table 12.3. Herbicides approved for managing weeds in onion. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.				
Active ingredient lb. a.i./A	Trade name product/A	MOA code	Crops	Weeds controlled / remarks
Glyphosate	(various formulations) consult label	9	Onion, garlic, leek, chive, shallot	Control of emerged broadleaf and grass weeds. Consult individual labels for restrictions.
Oxyfluorfen 2	(Goal 2XL) 2 EC 2.0 pt. (GoalTender) 4 EC 1.0 pt.	14	Onion (dry bulb), garlic (dry bulb)	Certain broadleaf weeds. Transplanted only. Apply within 2 days of transplanting. Necrotic lesions, twisting or stunting of plants can occur if applications are made during cool, wet weather or prior to the full development of the true leaves.
Paraquat 0.5 - 1.0	(Gramoxone) 2 SL 2 - 4 pt.	22	Onion (dry bulb)	Controls emerged weeds. Apply prior, during, or after planting, but before crop emergence. Only 2 applications a season. Use a non-ionic surfactant.
Paraquat 0.63 - 1.0	(Gramoxone) 2 SL 2.5 - 4.0 pt. (Firestorm) 3 SL 1.7 - 2.7 pt.	22	Onion (seeded), garlic	Emerged broadleaf and grass weeds. Apply as a preplant burndown. A maximum of 1 lb. a.i./A per season.
Pelargonic acid	(Scythe) 4.2 EC 3 - 10% v/v	27	Onion, garlic, leek, chive, shallot	Emerged broadleaf and grass weeds. Apply before emergence of crop. Product is a contact, nonselective, foliar applied herbicide. There is no residual activity.
Pendimethalin mineral 0.5 - 0.75 muck 1.0 - 2.0	(Prowl) 3.3 EC mineral 1.2 - 1.8 pt. muck 2.4 - 4.8 pt. (Prowl H20) 3.8 mineral 1.0 - 1.5 pt. muck 2.0 - 4.0 pt.	3	Onion (dry bulb), shallot (dry bulb), garlic	Mineral soils: Apply when crop has 2 to 9 true leaves. Muck soils: May be applied as sequential application: preemergence through loop stage, early postemergence (2 to 6 true leaf stage), late postemergence (6 to 9 true leaf stage). Do not exceed 5.9 lb. a.i./A in a growing season. PHI 45 days.
Pyraflufen 0.001 - 0.003	(ET Herbicide/Defoliant) 0.5 - 2.0 fl. oz.	14	Bulb vegetables	Emerged broadleaf and grass weeds. Apply as a preplant burndown treatment.
*** POSTEMERGENCE ***				
Bromoxynil 0.25 - 0.38	(Buctril) 2 EC 1.0 - 1.5 pt. (Buctril) 4 EC 0.5 - 1.5 pt.	6	Onion (Dry bulb)	Broadleaf weeds. Soil and onion leaves should be dry before application. Waxy coating on leaves reduces chances for injury. Varieties vary in sensitivity and use on a trial basis.
Bromoxynil 0.38 - 0.5	(Buctril) 2 EC 1.5 - 2.0 pt.	6	Garlic	Broadleaf weeds. Apply after emergence but before 12 in. tall. Weeds are most susceptible up to the 4-leaf stage or 2 in. height or 1 in. wide. PHI 112 days.
Carfentrazone up to 0.031	(Aim) 1.9 EW up to 2 fl. oz. (Aim) 2.0 EC up to 2 fl. oz.	14	Onion (dry bulb), garlic, leek, chive, shallot	Emerged broadleaf control. Post-direct hooded application to row middles for burndown of emerged weeds. Use crop oil concentrate or nonionic surfactant at recommended rates. PHI 0 days.
Clethodim 0.09 - 0.25	(Arrow, Select) 2 EC 6 - 16 fl. oz. (Select Max) 1 EC 9 - 32 fl. oz.	1	Onion (dry bulb), garlic, shallot (dry bulb)	Annual and perennial grass control. Some labels include green onion. Consult label for use of a crop oil concentrate or nonionic surfactant. PHI 45 days.
Clethodim 0.09 - 0.13	(Arrow, Select) 2 EC 6 - 8 fl. oz. (Select Max) 1 EC 12 - 16 fl. oz.	1	Chive, leek	Emerged annual and perennial grasses. Consult labels rate of COC or NIS. Do not apply more than 0.5 lb. a.i./A per season. Select Max is not registered in leek. PHI 14 days.
DCPA 4.5 - 6.0	(Dacthal) W-75 6 - 8 lb. (Dacthal) 6 F 6 - 8 pt.	3	Onion (dry bulb, green), leek, shallot	Annual grasses and certain broadleaf control. Apply at layby up to 14 weeks after planting. If weeds have emerged then cultivate or hand weed. Maximum rate of 10.5 lb. a.i./A per season.
Dimethanamid-p up to 1.0	(Outlook) 6 EC up to 21 fl. oz.	15	Onion (dry bulb), garlic, leek, shallot (dry bulb, green)	Broadleaf and grass weeds. Apply after the 2 true-leaf stage. Apply as a single application or split application. Split application of 10 - 14 fl. oz. followed by 7 to 10 fl. oz./A with 14 days between applications. Do not exceed 21 fl. oz./A per growing season. PHI 30 days.
Fluazifop 0.188	(Fusilade DX) 12 fl. oz.	1	Onion (dry bulb), garlic	Annual and perennial grass weeds. Do not apply more than 48 fl. oz./A per season. Include a COC or NIS. PHI 45 days.

Table 12.3. Herbicides approved for managing weeds in onion. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.					
Active ingredient lb. a.i./A	Trade name product/A	MOA code	Crops	Weeds controlled / remarks	
Oxyfluorfen 0.5	(Goal 2 XL) 2 EC 0.5 pt. direct seeded (GoalTender) 4 EC 0.25 pt. direct seeded	14	Onion (dry bulb), garlic (dry bulb)	Certain broadleaf weeds. Direct seeded only. Apply after the crop has 2 true leaves. Multiple treatments can be applied and do not exceed 0.5 lb. a.i./A. Necrotic lesions, twisting or stunting of plants can occur if applications are made during cool, wet weather or prior to development of true leaves. PHI 60 days.	
Sethoxydim 0.19 - 0.28	(Poast) 1.5 EC 1.0 - 1.5 pt.	1	Onion (dry bulb, green), garlic, leek, shallot	Grass weeds. Include a surfactant. Do not apply more than 4.5 pt./A per season. PHI 30 days.	
Treflan 0.35 - 0.5	(Treflan) 4 EC 0.75 - 1.0 pt. (Treflan TR-10) 3.75 - 5.0 lb.	3	Onion (dry bulb)	Broadleaf and grass weeds. Apply at layby to the soil surface between onion rows. Remove weeds that are preent. Do not apply in muck soils. PHI 60 days.	

Table 12.4. Insecticides labeled for management of arthropod pests of garlic, leeks, and shallots.

Labels change frequently. Be sure to read a current product label before applying any chemical.
Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Insect or mite pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate Product/acre	REI (hours)	Days to Harvest	Remarks ²
Aphids	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5-2.5 pt; 8F: 1-1.56 pt	24	3	Leeks, shallots, garlic. Three applications per year for garlic, two for all others.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Leeks, shallots, and garlic.
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (garlic). Do not apply more than 15.36 fl oz per acre per season.
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	BotaniGard ES (<i>Beauveria bassiana</i> strain GHA)	0.25 - 1 quart per acre. Apply in sufficient water to cover foliage, typically 5 - 100 gallons per acre.	4	0	
	-	Grandevo (<i>Chromobacterium</i> <i>subtsugae</i> strain PRAA4-1)	1-3 lb	4	0	Leek, garlic, shallot
	--	M-Pede 49% EC Soap, insecticidal	1-2% V/V	12	0	OMRI-listed.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
Beet Armyworm	--	Trilogy (extract of neem oil)	0.5-2.0% V/V	4	0	Apply morning or evening to reduce potential for leaf burn. Toxic to bees exposed to direct treatment. OMRI-listed.
	1A	*Lannate LV; *SP (methomyl)	LV: 1.5 pt; SP: 0.5 lb	48	7	Garlic only.
	28	Coragen (chlorantraniliprole)	3.5-5.0 fl oz	4	1	No more than 4 foliar applications or 15.4 fl oz of product per acre per crop. For entire Bulb Vegetable Group.
Caterpillars (includes cutworms and armyworms)	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Dry bulb and garlic only. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Garlic. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Leeks, shallots, and garlic.
	3A	*Pounce 25 WP (permethrin)	6.4-12.8 oz—garlic	12	1	Garlic

Table 12.4. Insecticides labeled for management of arthropod pests of garlic, leeks, and shallots. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.

Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Insect or mite pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate Product/acre	REI (hours)	Days to Harvest	Remarks ²
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (garlic). Do not apply more than 15.36 fl oz per acre per season.
	5	Entrust SC (spinosad)	3-8 fl oz	4	1	No more than 5 applications per year (29 fl oz product). For leeks, garlic, and shallots. OMRI-listed.
	5	Radiant SC (spinetoram)	5-10 fl oz	4	1	Use with an adjuvant.
	11A	Agree WG (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	1.0-2.0 lb	4	0	Apply when larvae are small for best control.
	11A	Biobit HP (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5-2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. Can be used in the greenhouse. For organic production.
	11A	Crymax WDG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5-2.0 lb	4	0	Use high rate for armyworms. Treat when larvae are young.
	11A	Deliver (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25-1.5 lb	4	0	Use higher rates for armyworms. OMRI-listed.
	11A	DiPel DF (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25-2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. OMRI-listed.
	11A	Javelin WG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.12-1.5 lb	4	0	Treat when larvae are young. Thorough coverage is essential. OMRI-listed.
	11A	Xentari DF (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	0.5-2.0 lb	4	0	Treat when larvae are young. Thorough coverage is essential. May be used in the greenhouse. Can be used in organic production.
	18	Intrepid 2F (methoxyfenozide)	4-12 fl oz	4	1	
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	Grandevo (<i>Chromobacterium subtsugae</i> strain PRAA4-1)	1-3 lb	4	0	Leek, garlic, and shallots
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
Fire ants	7A	Extinguish (S)-methoprene)	1-1.5 lb	4	0	Slow-acting IGR (insect growth regulator). Best applied early spring and fall where crop will be grown. Colonies will be reduced after three weeks and eliminated after 8 to 10 weeks. May be applied by ground equipment or aerially.
	7C	Esteem Ant Bait (pyriproxyfen)	1.5-2.0 lb	12	1	Dry bulb only.
Flea beetles	4A	Scorpion 35SL (dinotefuran)	foliar: 3.5-7.0 fl oz, soil: 8.75-10.5 fl oz	12	foliar: 1, soil: N/A	Garlic (Crop Groups 3-07A and 3-07B)
	5	Entrust SC (spinosad)	3-8 fl oz	4	1	No more than 5 applications per year (29 fl oz product). For leeks, garlic, and shallots. OMRI-listed.
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
Liriomyza Leafminers	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Garlic only. Label is registered only for garlic. Maximum of 2 lb ai/acre per season.

Table 12.4. Insecticides labeled for management of arthropod pests of garlic, leeks, and shallots. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.
Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Insect or mite pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate Product/acre	REI (hours)	Days to Harvest	Remarks ²
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Garlic. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Leeks, shallots, garlic.
	4A	Scorpion 35SL (dinotefuran)	foliar: 3.5-7.0 fl oz, soil: 8.75-10.5 fl oz	12	foliar: 1, soil: N/A	Garlic (Crop Groups 3-07A and 3-07B)
	5	Entrust SC (spinosad)	3-8 fl oz	4	1	No more than 5 applications per year (29 fl oz product). For leeks, garlic, and shallots. OMRI-listed.
	5	Radiant SC (spinetoram)	5-10 fl oz	4	1	Use with an adjuvant.
	6	*Agri-Mek SC (abamectin)	1.75-3.5 fl oz	12	30	Must be used with a non-ionic activator type wetting, spreading and/or penetrating adjuvant., not a binder sticker type adjuvant. Maximum of 10.25 fl oz per acre per season.
	17	Trigard (cyromazine)	2.66 oz	12	7	Maximum of 6 applications per crop.
	28	Exirel (cyazypyr)	13.5-20.5 fl oz	12	1	Do not apply more than 0.4 lb ai per acre of cyazypyr or cyantraniliprole-containing products per season. Toxic to bees. Do not allow drift to blooming crops or weeds.
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
Onion maggot	1B	*Diazinon AG500, *Diazinon 50W (diazinon)	AG500: 2-4 qt; 50W: 4-8 lb	72	preplant	Do not make more than one soil application per year. For leeks, garlic, and shallots.
	1B	Lorsban 75WG; *Advanced (chlorpyrifos)	75WG: 1.33 lb; Advanced: depends on row spacing	24	60	Dry bulb only. One application per year.
	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5-2.5 pt; 8F: 1-1.56 pt	leeks, shallots, garlic: 24	3	Leeks, shallots, and garlic. Three applications per year for garlic, two for all others.
	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Garlic only. Label is registered only for garlic. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Garlic. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Leeks, shallots, and garlic.
	3A	*Pounce 25 WP (permethrin)	6.4-19.2 oz—onions 6.4-12.8 oz—garlic	12	1	Garlic
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (garlic). Do not apply more than 15.36 fl oz per acre per season.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
Plant bugs	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Garlic. Maximum of 0.77 pt per acre per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (garlic). Do not apply more than 15.36 fl oz per acre per season.
	--	M-Pede 49% EC Soap, insecticidal	1-2% V/V	12	0	OMRI-listed.

Table 12.4. Insecticides labeled for management of arthropod pests of garlic, leeks, and shallots. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.

Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

See label to make sure for: Preproposed and other alternative products labeled for disease management.						
Insect or mite pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate Product/acre	REI (hours)	Days to Harvest	Remarks ²
Stink bugs	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Garlic only. Label is registered only for garlic. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Garlic. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Leeks, shallots, and garlic.
	3A	*Pounce 25 WP (permethrin)	6.4-19.2 oz—onions 6.4-12.8 oz—garlic	12	1	Garlic
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (garlic). Do not apply more than 15.36 fl oz per acre per season.
	4A	Scorpion 35SL (dinotefuran)	foliar: 3.5-7.0 fl oz, soil: 8.75-10.5 fl oz	12	foliar: 1, soil: N/A	Garlic (Crop Groups 3-07A and 3-07B)
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
Thrips (check label for species controlled)	1A	*Lannate LV; *SP (methomyl)	LV: 1.5-3.0 pt; SP: 0.5-1.0 lb	48	7	
	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5-2.5 pt; 8F: 1-1.56 pt	leeks, shallots, garlic: 24	3	Leeks, shallots, and garlic. Three applications per year for garlic, two for all others.
	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Garlic only. Label is registered only for garlic. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Garlic. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Leeks, shallots, and garlic.
	3A	*Pounce 25 WP (permethrin)	6.4-19.2 oz—onions 6.4-12.8 oz—garlic	12	1	Bulb onions and garlic
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (garlic). Do not apply more than 15.36 fl oz per acre per season.
	4A	Admire Pro (imidacloprid)	14.0 fl oz	12	21	Apply no more than 14 fl oz per acre per season.
	4A	Assail 30SG (acetamiprid)	5.0-8.0 oz	12	7	No more than 4 applications or 32 oz of product per acre per season.
	4A	Scorpion 35SL (dinotefuran)	foliar: 3.5-7.0 fl oz, soil: 8.75-10.5 fl oz	12	foliar: 1, soil: N/A	Garlic (Crop Groups 3-07A and 3-07B)
	5	Entrust SC (spinosad)	3-8 fl oz	4	1	No more than 5 applications per year (29 fl oz product). For leeks, garlic, and shallots. OMRI-listed.
	5	Radiant SC (spinetoram)	5-10 fl oz	4	1	Use with an adjuvant.
	6	*Agri-Mek SC (abamectin)	1.75-3.5 fl oz	12	30	Must be used with a non-ionic activator type wetting, spreading and/or penetrating adjuvant., not a binder sticker type adjuvant. Maximum of 10.25 fl oz per acre per season.
	7C	Knack IGR (pyriproxyfen)	8 fl oz	12	3	Maximum of 2 applications, at least 14 days apart. No activity against adult insects.
	28	Exirel (cyantraniliprole)	13.5-20.5 fl oz	12	1	Label contains pollinator protection warnings.
	28	Verimark (cyantraniliprole)	6.75-10 fl oz drip; 10-13.5 fl oz in-furrow at planting.	4	1	
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	BotaniGard ES (<i>Beauveria bassiana</i> strain GHA)	0.25 - 1 quart per acre. Apply in sufficient water to cover foliage, typically 5 - 100 gallons per acre.		0	4
	--	Grandevo (<i>Chromobacterium subtsugae</i> strain PRAA4-1)	1-3 lb	4	0	Leek, garlic, and shallot

Table 12.4. Insecticides labeled for management of arthropod pests of garlic, leeks, and shallots. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.
Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Insect or mite pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate Product/acre	REI (hours)	Days to Harvest	Remarks ²
	--	M-Pede 49% EC Soap, insecticidal	1-2% V/V	12	0	OMRI-listed.
	--	Requiem EC (extract of <i>Chenopodium ambrosioides</i>)	1.5-3.0 qt	4	0	Begin as soon as thrips are seen.
	--	Trilogy (extract of neem oil)	0.5-2.0% V/V	4	0	Apply morning or evening to reduce potential for leaf burn. Toxic to bees exposed to direct treatment. OMRI-listed.
Wireworms	1B	*Diazinon AG500, *Diazinon 50W (diazinon)	AG500: 2-4 qt; 50W: 4-8 lb	72	preplant	Do not make more than one soil application per year. For leeks, garlic, and shallots.

¹ Mode of Action (MOA) codes for plant pest insecticides from the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification v. 8.2, March 2017. Number codes (1 through 29) are used to distinguish the main insecticide mode of action groups, with additional letters for certain sub-groups within each main group. All insecticides within the same group (with same number) indicate same active ingredient or similar mode of action. This information must be considered for the insecticide resistance management decisions. - = unknown, or a mode of action that has not been classified yet.

² Information provided in this table applies only to Florida. Be sure to read a current product label before applying any product. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by the University of Florida Cooperative Extension Service nor discrimination against similar products or services not mentioned. OMRI listed: Listed by the Organic Materials Review Institute for use in organic production.

* Restricted use insecticide.

Table 12.5. Insecticides registered for managing insect pests of onions.

Labels change frequently. Be sure to read a current product label before applying any chemical.
Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate (Product/acre)	REI (hours)	Days to Harvest	Remarks ²
Aphids	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5-2.5 pt; 8F: 1.5 pt	12	3	Onions, bulb and green. Two applications per year. Suppression only.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Onions, bulb and green.
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (onions and garlic), not green onions. Do not apply more than 15.36 fl oz per acre per season. Suppression only.
	--	Azaguard (azadirachtin)	10-16 fl oz	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	BotaniGard ES (<i>Beauveria bassiana</i>)	0.5-2 qt/100 gal	4	0	May be used in greenhouses. Contact dealer for recommendations if an adjuvant must be used. Not compatible in tank mix with fungicides.
	--	Grandevo (<i>Chromobacterium subtsugae</i> strain PRAA4-1)	1-3 lb	4	0	Onion (bulb and green), and shallot
	--	M-Pede (potassium salts of fatty acids)	1-2% V/V	12	0	OMRI-listed.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	4	0	OMRI-listed.
	--	Trilogy (extract of neem oil)	0.5-2.0% V/V	4	0	Apply morning or evening to reduce potential for leaf burn. Toxic to bees exposed to direct treatment. OMRI-listed.
Caterpillars (includes cutworms and armyworms)	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Dry bulb and garlic only. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Bulb onions. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	6.4-19.2 oz	12	1	Bulb onions. Maximum 1 lb ai/acre per season.

Table 12.5. Insecticides registered for managing insect pests of onions. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.

Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate (Product/acre)	REI (hours)	Days to Harvest	Remarks ²
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (onions), not green onions. Do not apply more than 15.36 fl oz per acre per season.
	5	Entrust SC (spinosad)	3-6 fl oz	4	1	No more than 5 applications per year and no more than 29 fl oz product. For onions, bulb and green. OMRI-listed.
	5	Radiant SC (spinetoram)	5-10 fl oz	4	1	Use with an adjuvant. No more than 30 fl oz per acre per year.
	11A	Agree WG (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	1.0-2.0 lb	4	0	Apply when larvae are small for best control.
	11A	Biobit HP (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5-2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. Can be used in the greenhouse. For organic production.
	11A	Crymax WDG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.5-2.0 lb	4	0	Use high rate for armyworms. Treat when larvae are young.
	11A	Deliver (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25-1.5 lb	4	0	Use higher rates for armyworms. OMRI-listed.
	11A	DiPel DF (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.25-2.0 lb	4	0	Treat when larvae are young. Good coverage is essential. OMRI-listed.
	11A	Javelin WG (<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i>)	0.12-1.5 lb	4	0	Treat when larvae are young. Thorough coverage is essential. OMRI-listed.
	11A	Xentari DF (<i>Bacillus thuringiensis</i> subspecies <i>aizawai</i>)	0.5-2.0 lb	4	0	Treat when larvae are young. Thorough coverage is essential. May be used in the greenhouse. Can be used in organic production.
	18	Intrepid 2F (methoxyfenozide)	4-8 fl oz	4	1	Green onion subgroup only. Do not make more than 6 applications per year and do not apply more than 64 fl oz per acre per year.
	28	Coragen (chlorantraniliprole)	3.5-7.5 fl oz	4	1	Beet armyworm only. No more than 4 foliar applications or 15.4 fl oz of product per acre per crop.
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	4	0	OMRI-listed.
Fire ants	7A	Extinguish ((S)-methoprene)	1-1.5 lb	4	0	Slow-acting IGR (insect growth regulator). Best applied early spring and fall where crop will be grown. Colonies will be reduced after three weeks and eliminated after 8 to 10 weeks. May be applied by ground equipment or aerially.
	7C	Esteem Ant Bait (pyriproxyfen)	1.5-2.0 lb	12	1	Dry bulb only.
Flea beetles	4A	Scorpion 35SL (dinotefuran)	foliar: 3.5-7.0 fl oz, .	12	foliar: 1 soil: N/A	Bulb and green onions (Crop Groups 3-07A and 3-07B). Do not apply more than 10.5 fl oz per acre per season.
	5	Entrust SC (spinosad)	3-6 fl oz	4	1	No more than 5 applications per year and no more than 29 fl oz product. For onions, bulb and green. OMRI-listed.
Leafminers	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Dry bulb only. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Bulb onions. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	6.4-19.2 oz—onions	12	1	Bulb onions. Maximum 1 lb ai/acre per season.
	4A	Scorpion 35SL (dinotefuran)	foliar: 5.25-7.0 fl oz, soil: 8.75-10.5 fl oz	12	foliar: 1 soil: N/A	Bulb and green onions (Crop Groups 3-07A and 3-07B). Do not apply more than 10.5 fl oz per acre per season.
	5	Entrust SC (spinosad)	3-6 fl oz	4	1	No more than 5 applications per year and no more than 29 fl oz product. For onions, bulb and green. OMRI-listed.
	5	Radiant SC (spinetoram)	6-10 fl oz	4	1	Use with an adjuvant. No more than 30 fl oz per acre per year.

Table 12.5. Insecticides registered for managing insect pests of onions. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.
Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate (Product/acre)	REI (hours)	Days to Harvest	Remarks ²
	6	*Agri-Mek SC (abamectin)	1.75-3.5 fl oz	12	30	Must be used with a non-ionic activator type wetting, spreading and/or penetrating adjuvant., not a binder sticker type adjuvant. Maximum of 10.25 fl oz per acre per season.
	17	Trigard (cyromazine)	2.66 oz	12	7	Maximum of 6 applications per crop.
	28	Exirel (cyazypyr)	13.5-20.5 fl oz	12	1	Do not apply more than 0.4 lb ai per acre of cyazypyr or cyantraniliprole-containing products per season. Toxic to bees. Do not allow drift to blooming crops or weeds.
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
Onion maggot	1B	*Diazinon AG500, *Diazinon 50W (diazinon)	AG500: 2-4 qt, 50W: 4-8 lb	72	preplant	Do not make more than one soil application per year. For onions, bulb and green.
	1B	Lorsban 75WG; *Advanced (chlorpyrifos)	75WG: 1.33 lb; Advanced: 32 fl oz	24	60	Dry bulb only. One application per year.
	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5-2.5 pt; 8F: 1.5 pt	12	3	Onions, bulb and green. No more than two applications per year.
	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Dry bulb only. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.28 fl oz	24	14	Bulb onions. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	6.4-19.2 oz	12	1	Bulb onions. Maximum 1 lb ai/acre per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (onions and garlic), not green onions. Do not apply more than 15.36 fl oz per acre per season.
	--	Neemix 4.5 EC (azadirachtin)	4-16 fl oz	12	0	OMRI-listed.
Plant bugs	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Bulb onions. Maximum of 0.77 pt per acre per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (onions and garlic), not green onions. Do not apply more than 15.36 fl oz per acre per season.
	--	M-Pede (potassium salts of fatty acids)	1-2% V/V	12	0	OMRI-listed.
Stink bugs	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Dry bulb only. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Bulb onions. Maximum of 0.77 pt per acre per season.
	3A	*Mustang (zeta-cypermethrin)	2.4-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	6.4-19.2 oz	12	1	Bulb onions. Maximum 1 lb ai/acre per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (onions), not green onions. Do not apply more than 15.36 fl oz per acre per season.
	4A	Scorpion 35SL (dinotefuran)	foliar: 3.5-7.0 fl oz	12	1	Bulb and green onions (Crop Groups 3-07A and 3-07B). Do not apply more than 10.5 fl oz per acre per season.
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
Thrips (check label for species controlled)	1B	Malathion 5EC, 8F (malathion)	5EC: 1.5-2 pt; 8F: 1.56 pt	12	3	Onions, bulb and green. No more than two applications per year.
	3A	*Ambush 25W (permethrin)	6.4-19.2 oz	12	1	Dry bulb. Maximum of 2 lb ai/acre per season.
	3A	*Declare Insecticide (gamma-cyhalothrin)	0.77-1.54 fl oz	24	14	Bulb onions. Maximum of 0.77 pt per acre per season.

Table 12.5. Insecticides registered for managing insect pests of onions. (continued)

Labels change frequently. Be sure to read a current product label before applying any chemical.

Also refer to Table 19.1 for biopesticide and other alternative products labeled for disease management.

Pest	MOA Code ¹	Trade Name (Active Ingredient) *Restricted	Rate (Product/acre)	REI (hours)	Days to Harvest	Remarks ²
	3A	*Mustang (zeta-cypermethrin)	3.2-4.3 oz	12	7	Do not apply more than 21.5 oz per acre per season. Onions, bulb and green.
	3A	*Pounce 25 WP (permethrin)	9.6-19.2 oz	12	1	Bulb onions. Maximum 1 lb ai/acre per season.
	3A	*Warrior II (lambda-cyhalothrin)	0.96-1.92 fl oz	24	14	For bulb crops only (onions), not green onions. Do not apply more than 15.36 fl oz per acre per season.
	4A	Admire Pro (imidacloprid)	14.0 fl oz	12	21	Apply no more than 14 fl oz per acre per season.
	4A	Assail 30SG (acetamiprid)	5.0-8.0 oz	12	7	No more than 4 applications or 32 oz of product per acre per season.
	4A	Scorpion 35SL (dinotefuran)	foliar: 5.25-7.0 fl oz, soil: 8.75-10.5 fl oz	12	foliar: 1 soil: N/A	Bulb and green onions (Crop Groups 3-07A and 3-07B). Do not apply more than 10.5 fl oz per acre per season.
	5	Entrust SC (spinosad)	3-8 fl oz	4	1	No more than 5 applications per year (29 fl oz product). For onions, bulb and green. OMRI-listed.
	5	Radiant SC (spinetoram)	6-10 fl oz	4	1	Use with an adjuvant. No more than 30 fl oz per acre per year.
	6	*Agri-Mek SC (abamectin)	1.75-3.5 fl oz	12	30	Must be used with a non-ionic activator type wetting, spreading and/or penetrating adjuvant., not a binder sticker type adjuvant. Maximum of 10.25 fl oz per acre per season.
	7C	Knack IGR (pyriproxyfen)	8 fl oz	12	3	Maximum of 2 applications, at least 14 days apart. No activity against adult insects. Onion (dry bulb only).
	28	Exirel (cyazypyr)	13.5-20.5 fl oz	12	1	Do not apply more than 0.4 lb ai per acre of cyazypyr or cyantraniliprole-containing products per season. Toxic to bees. Rotate with other effective modes of action. Begin making applications when populations are low. If populations are above threshold, use an effective knockdown product prior to applying Exirel
	28	Verimark (cyantraniliprole)	at planting: 10-13.5 fl oz drip chemigation: 6.75-10 fl oz	1	4	Both at plant and drip chemigation. Do not apply more than 2 drip chemigation applications. Begin drip applications when populations first appear. Rotate with other effective modes of action.
	--	Aza-Direct (azadirachtin)	1-2 pts, up to 3.5, if needed	4	0	Antifeedant, repellent, insect growth regulator. OMRI-listed.
	--	Azatin XL (azadirachtin)	5-21 fl oz	4	0	Antifeedant, repellent, insect growth regulator.
	--	BotaniGard ES (<i>Beauveria bassiana</i>)	0.5-2 qt/100 gal	4	0	May be used in greenhouses. Contact dealer for recommendations if an adjuvant must be used. Not compatible in tank mix with fungicides.
	--	M-Pede (potassium salts of fatty acids)	1-2% V/V	12	0	OMRI-listed.
	--	Requiem EC (extract of <i>Chenopodium ambrosioides</i>)	1.5-3.0 qt	4	0	Begin as soon as thrips are seen.
	--	Trilogy (extract of neem oil)	0.5-2.0% V/V	4	0	Apply morning or evening to reduce potential for leaf burn. Toxic to bees exposed to direct treatment. OMRI-listed.
Wireworms	1B	*Diazinon AG500, *Diazinon 50W (diazinon)	AG500: 3-4 qt, 50W: 4-8 lb	72	preplant	Do not make more than one soil application per year. For onions, bulb and green.

¹ Mode of Action (MOA) codes for plant pest insecticides from the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification v. 8.2, March 2017. Number codes (1 through 29) are used to distinguish the main insecticide mode of action groups, with additional letters for certain sub-groups within each main group. All insecticides within the same group (with same number) indicate same active ingredient or similar mode of action. This information must be considered for the insecticide resistance management decisions. - = unknown, or a mode of action that has not been classified yet.

² Information provided in this table applies only to Florida. Be sure to read a current product label before applying any product. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by the University of Florida Cooperative Extension Service nor discrimination against similar products or services not mentioned. OMRI listed: Listed by the Organic Materials Review Institute for use in organic production.

* **Restricted use insecticide.**

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action.

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical. Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
Bacterial blight <i>(Xanthomonas axonopodis</i> pv. <i>allii</i>)	M1	(copper compounds) Many brands available: Badge SC, Badge X2, Champ DP, Champ F2 FL, Champ WG, Cuprofix Ultra 40D, Cueva,Cuproxat, Kentan DF, Kocide 3000, Kocide 2000, Kocide DF, Kop-hydroxide, Master Cop, Nordox, Nordox 75WG, Nu Cop 3L, Nu Cop 50DF, Nu Cop HB	SEE INDIVIDUAL LABEL		0	Varies by product from 4 hr to 2 days.	Frequent use of copper fungicides may cause foliar burn
	M1 & M3	Mankocide (copper hydroxide + mancozeb)	2.25 lbs	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
(Suppression only)	21	Actigard 50WG (acibenzolar-S-methyl)	1 oz.	4 oz	7	0.5	Do not apply to plants that are stressed or prior or within 5 days transplanting. Supression only
(Suppression only)	27 & 11	Tanos (cymoxanil + famoxadone)	8 oz	84 oz	3	0.5	See label
Black mold <i>(Aspergillus niger)</i>	9 & 12	Switch 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	Do not make more than 2 consecutive appl. Do not plant crops not on the label for 30 days after last application
Botrytis leaf blight <i>(Botrytis squamosa)</i>	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL		7	1	Do not apply to exposed bulbs
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF, Initiate 720, Initiate Zn	SEE INDIVIDUAL LABEL		7 (dry bulb); 14 (green)	1	
	M1 & M3	Mankocide (copper hydroxide + mancozeb)	2.25 lbs	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevada 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL		7	1	
(Suppression only)	3	(propiconazole) Many brands available: Amtide proiconazole 41.8% EC, Bumper 41.8 EC, Bumper ES, Fitness, Propicure, Propimax EC, Propistar EC, Shar-Shield PPZ, Tide Propiconazole EC, Tilt	SEE INDIVIDUAL LABEL		See label	0.5	
	3 & 7	Luna Experience (fluopyram + tebuconazole)	12.8 fl oz	25.6 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	14	Botran 5F (DCNA dicloran)	2 to 3 1/5 lbs	See label	14	See label	See label
	4 & M5	Ridomil Gold Bravo SC, (mefenoxam + chlorothalonil)	See label		7 (dry bulb); 14/21 (green)	2	
	7	Endura (boscalid)	6.8 oz	41 oz	7	0.5	Alternate applications with a different fungicide group.
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	See label
	7 & 9	Luna Tranquility (fluopyram + pyrimethanil)	27 fl oz	54.7 fl oz	7	0.5	Do not make more than 2 applications per year.
	7 & 11	Merivon (fluxapyroxad + pyraclostrobin)	11 fl oz	33 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	9	Vanguard WG (cyprodinil)	10 oz	28 oz	7	0.5	Do not make more than 2 consecutive applications.

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action. (continued)

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical.							
Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
(Suppression only)	9	Scala SC (pyrimethanil)	18 fl oz	54 fl oz	7	0.5	Do not make more than 2 consecutive applications.
	11	Quadris, (azoxystrobin)	See label	92.3 fl oz	See label	4 hr	Alternate applications with a different fungicide group.
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. Make no more than 5 applications/crop.
	9 & 3	Inspire Super (cyprodinil + difenoconazole)	20 fl.oz	See label	14 (green); 7 (dry bulb)	0.5	Do not make more than 2 applications before alternate with a different fungicide group
	11 & M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	See label	14	0.5	Alternate applications with different fungicide groups. Make no more than 3 applications/ crop. Do not tank mix with COC, MSO or silicon adjuvant.
	11 & 3	Quadris Top (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	Make no more than 1 application before alternating to another fungicide with a non- QoI (Group 11) mode of action
	11 & 3	Quilt Xcel (azoxystrobin + propiconazole)	21 fl oz (dry); 26 fl oz (green)	56 fl oz	14 (dry bulb); 0 (green)	0.5	Make only 1 appl. before alternating to a non-group 11 fungicide
	11 & 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	Do not make more than 2 consecutive applications. No more than 6 appl/ crop.
	9 & 12	Switch 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	Do not make more than 2 consecutive appl. Do not plant crops not on the label for 30 days after last appl.
22 & M5	Zing (zoxamide + chlorothanil)	30 fl oz	7.5 qt	7	0.5	Do not make more than 8 applications per year.	
29	Omega 500F (fluazinam)	1 pt	6 pt	7	2		
Botrytis neck rot (Botrytis allii)	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL		7	1	Do not apply to exposed bulbs
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF, Initiate 720, Initiate Zn	SEE INDIVIDUAL LABEL		7 (dry bulb); 14 (green)	1	
(Suppression only)	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL		7	1	
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	
	14	Botran 5F (DCNA dicloran)	3.5 lbs	See label	14	See label	
	9	Vanguard WG (cyprodinil)	10 oz	28 oz	7	0.5	Do not make more than 2 consecutive applications.
(Suppression only)	9	Scala SC (pyrimethanil)	18 fl oz	54 fl oz	7	0.5	Do not make more than 2 consecutive applications.
	11 & M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	See label	14	0.5	Alternate applications with different fungicide groups. Make no more than 3 applications/ crop. Do not tank mix with COC, MSO or silicon adjuvant.

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action. (continued)

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical. Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
(Suppression only)	11 & 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	Do not make more than 2 consecutive applications. No more than 6 applications/ crop
	9 & 12	Switch 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	Do not make more than 2 consecutive appl. Do not plant crops not on the label for 30 days after last appl.
	29	Omega 500F (fluazinam)	1 pt	6 pt	7	2	
	U15 & M5	Orondis Opti (oxathiapiprolin + chlorothalonil)	2.5 pt	10 pt	7	0.5	Do not make more than 2 sequential applications and do not exceed more than 33% of all fungicide applications with Orondis.
Cercospora leaf spot (Cercospora duggiae)	11 & 3	Quadris Top (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	Make no more than 1 application before alternating to another fungicide with a non- QoI (Group 11) mode of action
Cladosporium leaf blotch (Cladosporium allii)	11	Quadris, (azoxystrobin)	See label	92.3 fl oz	See label	4 hr	Alternate applications with a different fungicide group.
	11 & M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	See label	14	0.5	Alternate applications with different fungicide groups. Make no more than 3 applications/ crop. Do not tank mix with COC, MSO or silicon adjuvant.
	11 & 3	Quilt Xcel (azoxystrobin + propiconazole)	21 fl oz (dry); 26 fl oz (green)	56 fl oz	14 (dry bulb); 0 (green)	0.5	Make only 1 appl. before alternating to a non-group 11 fungicide
	11 & 3	Quadris Top (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	Make no more than 1 application before alternating to another fungicide with a non- QoI (Group 11) mode of action
	9 & 3	Inspire Super (cyprodinil + difenoconazole)	20 fl.oz	See label	14 (green); 7 (dry bulb)	0.5	Do not make more than 2 applications before alternate with a different fungicide group
Damping-off	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP	SEE INDIVIDUAL LABEL		7	1	Do not apply to exposed bulbs
	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevada 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL		7	1	
	12	Cannonball WP (fludioxonil)	7 oz	32 oz	7	0.5	
	12	Maxim 4FS (fludioxonil)	0.16 fl oz/ 100 lbs seed				Seed treatment only.
Downy mildew (Peronospora destructor)	M1	(copper compounds) Many brands available: Badge SC, Badge X2, Basic Copper 53, C-O-C-S WDG, Champ DP, Champ F2 FL, Champ WG, C-O-C DF, C-O-C WP, Copper Count N, Cuprofix Ultra 40D, Cueva, Kentan DF, Kocide 3000, Kocide 2000, Kocide DF, Nordox, Nordox 75WP, Nu Cop 3L, Nu Cop 50DF, Nu Cop HB	SEE INDIVIDUAL LABEL		0	Varies from 4 hr to 2 days.	Frequent use of copper fungicides may cause foliar burn
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF. Initiate 720. Initiate Zn	SEE INDIVIDUAL LABEL		7 (dry bulb); 14 (green)	1	

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action. (continued)

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical. Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
(Suppression only)	M1 & M2	Top cop with sulfur (basic copper sulfate + sulfur)	2-3 qts	See label	0	1	Do not use during hot weather or within 4 weeks of an oil application.
	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP	SEE INDIVIDUAL LABEL		7	1	Do not apply to exposed bulbs
	M1 & M3	Mankocide (copper hydroxide + mancozeb)	2.25 lbs	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
	U15 & M5	Orondis Opti (oxathiapiprolin + chlorothalonil)	2.5 pt	10 pt	7	0.5	Do not make more than 2 sequential applications and do not exceed more than 33% of all fungicide applications with Orondis.
	U15 & 40	Orondis Ultra (oxathiapiprolin + mandipropamid)	8 fl oz	32 fl oz (dry); 24 fl oz (green)	7	4 hr	Do not make more than 2 sequential applications and do not exceed more than 33% of all fungicide applications with Orondis.
	4 & M1	Ridomil Gold/Copper (mefenoxam + copper hydroxide)	2 lbs		7 (green); 10 (dry bulb)	2	Maximum 0.4 lbs/A a.i. of metalaxyl or mefenoxam
	4 & M3	Ridomil Gold MZ WG (mefenoxam + mancozeb)	2.5 lbs		7 (dry bulb)	2	
	4 & M5	Ridomil Gold Bravo SC (mefenoxam + chlorothalonil)	See label		7 (dry bulb); 14/21 (green)	2	
	11	Quadris, (azoxystrobin)	See label	92.3 fl oz	See label	4 hr	Alternate applications with a different fungicide group.
	11	Reason 500SC (fenamidone)	5.5 fl oz	22 fl oz	7	0.5	Alternate applications with a different fungicide group.
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. Make no more than 5 appl/crop.
	11 & M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	See label	14	0.5	Alternate applications with different fungicide groups. Make no more than 3 applications/ crop. Do not tank mix with COC, MSO or silicon adjuvant.
	11 & 3	Quilt Xcel (azoxystrobin + propiconazole)	21 fl oz (dry); 26 fl oz (green)	56 fl oz	14 (dry bulb); 0 (green)	0.5	Make only 1 appl. before alternating to a non-group 11 fungicide
	11 & 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	Do not make more than 2 consecutive applications. No more than 6 appl/ crop
	21	Actigard 50WG (acibenzolar-S-methyl)	1 oz.	4 oz	7	0.5	Do not apply to plants that are stressed or prior or within 5 days transplanting.
	21	Ranman (cyazofamid)	3 fl oz	16.5 fl oz	0	0.5	Alternate with fungicides of dissimilar modes of action. Do not make more than 3 sequential applications.
	22 & M5	Zing (zoxamide + chlorothanil)	30 fl oz	7.5 qt	7	0.5	Do not make more than 8 applications per year.
	27 & 11	Tanos (cymoxanil + famoxadone)	8 oz	84 oz	3	0.5	
	29	Omega 500F (fluazinam)	1 pt	6 pt	7	2	
	33	Aliette WDG, Linebacker WDG (fosetyl-Al)	3 lb	7 appl.	See label	See label	Do not tank mix with copper fungicides, adjuvants or foliar fertilizers
33	(mono- and di-potassium salts of phosphorous acid)	SEE INDIVIDUAL LABEL		0	4 hr	See label for details.	

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action. (continued)

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical. Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
		Many brands available: Alude, Confine Extra, Fosiphite, Fungi-phite, K-Phite, Oxiphos, Phiticide, Phostrol, Prophyt, Rampart, Reveille					
	40	Forum (dimethomorph)	6 oz	30 oz	0	0.5	Must be applied in a tank mix with another fungicide active against downy mildew. Do not make more than 2 sequential applications.
	40	Revus (mandipropamid)	8 fl oz	32 fl oz (dry); 24 fl oz (green)	7	4 hr	Make no more than 2 consecutive appl. before switching to a non-group 40 fungicide. A silicon-based adjuvant must be added at recommended rates
	43	Presidio (fluopicolide)	4 fl oz.		2	0.5	Do not make more than 4 applications of Presidio per season.
	21	Actigard 50WG (Acibenzolar-s-methyl)	1 oz.	4 oz	7	0.5	Do not apply to plants that are stressed or prior or within 5 days transplanting. Supression only
	40 & 45	Zampro (dimethomorph + ametoctradin)	14 fl oz	42 fl oz	0	0.5	See label
Iris Yellow Spot (Iris yellow spot virus) (Suppression only)	21	Actigard 50WG (acibenzolar-S-methyl)	1 oz.	4 oz	7	0.5	Do not apply to plants that are stressed or prior or within 5 days transplanting. Supression only.
Powdery mildew	M2	(sulfur) Many brands available: Kumulus DF, Micro Sulf, Microfine Sulfur, Microthiol Disperss, Sulfur 90W, Yellow Jacket Wettable Sulfur	SEE INDIVIDUAL LABEL		0	1	Do not use during hot weather or within 4 weeks of an oil application.
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	
	9 & 3	Inspire Super (cyprodinil + difenoconazole)	20 fl.oz	See label	14 (green); 7 (dry bulb)	0.5	Do not make more than 2 applications before alternate with a different fungicide group
	11 & 3	Quadris Top (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	Make no more than 1 application before alternating to another fungicide with a non- Qol (Group 11) mode for action
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. Make no more than 5 appl/crop.
Purple blotch (Alternaria porri)	M1	(copper compounds) Many brands available: Badge SC, Badge X2, Basic Copper 53, C-O-C-S WDG, Champ DP, Champ F2 FL, Champ WG, C-O-C DF, C-O-C WP, Copper Count N, Cuprofix Ultra 40D, Cueva, Cuproxat, Kentan DF, Kocide 3000, Kocide 2000, Kocide DF, Kop-hydroxide, Nordox, Nordox 75WP, Nu Cop 3L, Nu Cop 50DF, Nu Cop HB	SEE INDIVIDUAL LABEL		0	Varies from 4 hr to 2 days.	Frequent use of copper fungicides may cause foliar burn
	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL		7	1	Do not apply to exposed bulbs
	M5	(chlorothalonil) Many brands available: Bravo Ultrex, Bravo Weather Stik, Bravo Zn, Chloronil 720, Chlorothalonil 720 EC, Echo 720, Echo 90 DF, Echo Zn, Equus 500 Zn, Equus 720 SST, Equus DF, Initiate 720, Initiate Zn	SEE INDIVIDUAL LABEL		7 (dry bulb); 14 (green)	1	
	M1 & M2	Top cop with sulfur (basic copper sulfate + sulfur)	3 qts	See label	0	1	Do not use during hot weather or within 4 weeks of an oil application.

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action. (continued)

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical. Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
	M1 & M3	Mankocide (copper hydroxide + mancozeb)	2.25 lbs	160 lb (per crop)	7	2	Do not apply to exposed bulbs.
	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL		7	1	
	3	(propiconazole) Many brands available: Amtide proiconazole 41.8% EC, Bumper 41.8 EC, Bumper ES, Fitness, Propimax EC, Propistar EC, Tide Propionazole EC, Tilt, Shar-Shield PPZ	SEE INDIVIDUAL LABEL		See label	0.5	
	3	(tebuconazole) Many brands available: Monsoon, Onset 3.6L, Orius 3.6 F, Tebu-Crop 3.6F, Tebustar 3.6 L, Tebuzol 3.6F, Toledo	SEE INDIVIDUAL LABEL		7	0.5	
	3 & 7	Luna Experience (fluopyram + tebuconazole)	12.8 fl oz	25.6 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	3 & M5	Muscle ADV (tebuconazole + chlorothalonil)	1.6 pt	3.2 pt	7	0.5	Apply in a protective fungicide schedule.
	4 & M5	Ridomil Gold Bravo SC, (mefenoxam + chlorothalonil)	See label		7 (dry bulb); 14/21 (green)	2	
	7	Endura (boscalid)	6.8 oz	41 oz	7	0.5	Alternate applications with a different fungicide group.
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	3	0.5	See label
	7 & 9	Luna Tranquility (fluopyram + pyrimethanil)	27 fl oz	54.7 fl oz	7	0.5	Do not make more than 2 applications per year.
	7 & 11	Merivon (fluxapyroxad + pyraclostrobin)	11 fl oz	33 fl oz	7	0.5	Alternate with fungicides of dissimilar modes of action.
	9	Vangard WG (cyprodinil)	10 oz	28 oz	7	0.5	Do not make more than 2 consecutive applications.
	9	Scala SC (pyrimethanil)	18 fl oz	54 fl oz	7	0.5	Do not make more than 2 consecutive applications.
	11	Quadris, (azoxystrobin)	See label	92.3 fl oz	See label	4 hr	Alternate applications with a different fungicide group.
	11	Reason 500SC (fenamidone)	5.5 fl oz	22 fl oz	7	0.5	Alternate applications with a different fungicide group.
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. Make no more than 5 applications/crop.
	9 & 3	Inspire Super (cyprodinil + difenoconazole)	20 fl.oz	See label	14 (green); 7 (dry bulb)	0.5	Do not make more than 2 applications before alternate with a different fungicide group
	11 & M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	See label	14	0.5	Alternate applications with different fungicide groups. Make no more than 3 applications/ crop. Do not tank mix with COC, MSO or silicon adjuvant.
	11 & 3	Quadris Top (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	Make no more than 1 application before alternating to another fungicide with a non- QoI (Group 11) mode for action
	11 & 3	Quilt Xcel (azoxystrobin + propiconazole)	21 fl oz (dry); 26 fl oz (green)	56 fl oz	14 (dry bulb); 0 (green)	0.5	Make only 1 appl. before alternating to a non-group 11 fungicide
	11 & 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	Do not make more than 2 consecutive applications. No more than 6 appl/ crop.

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action. (continued)

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical. Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
(Suppression only)	9 & 12	Switch 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	Do not make more than 2 consecutive appl. Do not plant crops not on the label for 30 days after last appl.
	22 & M5	Zing (zoxamide + chlorothanil)	30 fl oz	7.5 qt	7	0.5	Do not make more than 8 applications per year.
	27 & 11	Tanos (cymoxanil + famoxadone)	8 oz	84 oz	3	0.5	
	29	Omega 500 F (fluazinam)	1 pt	6 pt	7	2	
	33	Aliette WDG, Linebacker WDG (fosetyl-Al)	3 lb	7 appl.	See label	See label	Do not tank mix with copper fungicides, adjuvants or foliar fertilizers
	U15 & M5	Orondis Opti (oxathiapiprolin + chlorothalonil)	2.5 pt	10 pt	7	0.5	Do not make more than 2 sequential applications and do not exceed more than 33% of all fungicide applications with Orondis.
Pythium Damping-off	4	(metalaxyl) Many brands available: Allegiance, Aquire, MetaStar 2E AG, Sebring 2.65 ST, Sebring 318 FS, Sebring 480 FS	SEE INDIVIDUAL LABEL				
	4	(mefenoxam) Many brands available: Apron XL, Ridomil Gold EC, Ridomil Gold SL, Ultra Flourish	SEE INDIVIDUAL LABEL		N/A	2	
	11 & 4	Uniform (azoxystrobin + mefenoxam)	See label				
Rhizoctonia Damping off	11	Dynasty (azoxystrobin)	0.38 fl oz/ 100 lb seed	See label	See label	4 hr	Seed treatment only.
	11	Quadris (azoxystrobin)					
	11 & 4	Uniform (azoxystrobin + mefenoxam)	See label				
Rust (<i>Puccinia allii</i>)	3	(tebuconazole) Many brands available: Monsoon, Onset 3.6L, Orius 3.6 F, Tebu-Crop 3.6F, Tebustar 3.6 L, Tebuzol 3.6F, Toledo	SEE INDIVIDUAL LABEL		7	0.5	
	3 & M5	Muscle ADV (tebuconazole + chlorothalonil)	1.6 pt	3.2 pt	7	0.5	Apply in a protective fungicide schedule.
	11	Quadris, (azoxystrobin)	See label	92.3 fl oz	See label	4 hr	Alternate applications with a different fungicide group.
	11	Cabrio EG (pyraclostrobin)	12 oz	72 oz	7	0.5	Alternate applications with a different fungicide group. Make no more than 5 appl/crop.
	11 & 3	Quilt Xcel (azoxystrobin + propiconazole)	21 fl oz (dry); 26 fl oz (green)	56 fl oz	14 (dry bulb); 0 (green)	0.5	Make only 1 appl. before alternating to a non-group 11 fungicide.
	9 & 3	Inspire Super (cyprodinil + difenoconazole)	20 fl.oz	See label	14 (green); 7 (dry bulb)	0.5	Do not make more than 2 applications before alternate with a different fungicide group.
Seedling blight	M3	(mancozeb) Many brands available: Dithane DF, Dithane F45, Dithane M45, Koverall, Manzate, Manzate Pro-Stik, Penncozeb 4FL, Penncozeb 75DF, Penncozeb 80WP, Roper DF	SEE INDIVIDUAL LABEL		7	1	Do not apply to exposed bulbs
	12	Maxim 4FS Spirato 480 FS (fludioxonil)	0.16 fl oz/ 100 lbs seed	-	-	0.5	

Table 12.6. Onion fungicides ordered by disease and then FRAC group according to their mode of action. (continued)

Pertinent Diseases or Pathogens	Fungicide Group ¹	Chemical (active ingredients)	Max. Rate/Acre		Min. Days to		Remarks ²
			Applic.	Season	Harvest	Reentry	
Labels change frequently. Be sure to read a current product label before applying any chemical. Refer to Table 19.2 for biopesticide and other alternative products labeled for disease management.							
	11	Dynasty (azoxystrobin)	0.38 fl oz/ 100 lb seed	See label	See label	4 hr	
	14	Botran 5F (DCNA dicloran)	3.5 lbs	See label	14	See label	
Stemphylium Blight (<i>Stemphylium vesicarium</i>)	2	(iprodione) Many brands available: Enclosure 4, Iprodione 4L AG, Meteor, Nevado 4F, Rovral 4 Flowable	SEE INDIVIDUAL LABEL		7	1	
	7	Fontelis (Penthiopyrad)	24 fl oz	72 fl oz	3	0.5	
	9 & 3	Inspire Super (cyprodinil + difenoconazole)	20 fl.oz	See label	14 (green); 7 (dry bulb)	0.5	Do not make more than 2 applications before alternate with a different fungicide group
	11 & 3	Quadris Top (azoxystrobin + difenoconazole)	14 fl oz	42 fl oz	7	0.5	Make no more than 1 application before alternating to another fungicide with a non- Qol (Group 11) mode for action
	11 & 7	Pristine (pyraclostrobin + boscalid)	18.5 oz	111 oz	7	0.5	Do not make more than 2 consecutive applications. No more than 6 appl/ crop.
	9 & 12	Switch 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	Do not make more than 2 consecutive appl. Do not plant crops not on the label for 30 days after last appl.
	29	Omega (fluazinam)	1 pt	6 pt	7	1	Begin sprays at first development and continue on 7-14 day schedule.
White rot (<i>Sclerotium cepivorum</i>)	1	(thiophanate-methyl) Many brands available: Incognito 4.5 F, NuFarm T-Methyl 4.5 T, NuFarm T-Methyl 70 WSB, Thiophanate-methyl 85 WDG, Topsin 4.5 FL, Topsin M 70 WP, Topsin M 70 WDG, Topsin M WSB	SEE INDIVIDUAL LABEL			3	
	3	(tebuconazole) Many brands available: Onset 3.6L, Orius 3.6 F, Tebu-Crop 3.6F, Tebustar 3.6 L, Tebuzol 3.6F, Toledo	SEE INDIVIDUAL LABEL		7	0.5	
	7	Fontelis (penthiopyrad)	24 fl oz	72 fl oz	7	0.5	Spray into empty furrow at time of planting.
	11 & M5	Quadris Opti (azoxystrobin + chlorothalonil)	3.7 pt	see label	14	0.5	Alternate applications with different fungicide groups. Make no more than 3 applications/ crop. Do not tank mix with COC, MSO or silicon adjuvant.
	9 & 12	Switch 62.5WG (cyprodinil + fludioxonil)	14 oz	56 oz	7	0.5	Do not make more than 2 consecutive appl. Do not plant crops not on the label for 30 days after last appl.
	12	Cannonball WP (fludioxonil)	7 oz	32 oz	7	0.5	
	12	Maxim 4FS (fludioxonil)	0.16 fl oz/ 100 lbs seed	-	-	0.5	
	14	Botran 5F (DCNA dicloran)	3.5 lbs	See label	14	0.5	

¹ FRAC code (fungicide group): Number (1 through 45) and letters (U and P) are used to distinguish the fungicide mode of action groups. All fungicides within the same group (with same number or letter) indicate same active ingredient or similar mode of action. This information must be considered for the fungicide resistance management decisions. U = unknown, or a mode of action that has not been classified yet and is typically associated with another number; P = host plant defense inducers. Source: FRAC Code List 2018; <http://www.frac.info/> (FRAC = Fungicide Resistance Action Committee).

² Information provided in this table applies only to Florida. Be sure to read a current product label before applying any chemical. The use of brand names and any mention or listing of commercial products or services in the publication does not imply endorsement by the University of Florida Cooperative Extension Service nor discrimination against similar products or services not mentioned.

Table 12.7. Fumigant nematicides for onions in Florida.

Nematicide	Broadcast Application ¹		In the Row Applications
	Gallons or Lbs Per acre	Fl oz /1000 ft / chisel spaced 12" apart	
Telone II ^{2,3}	9 to 12 gal	26 to 35	For any row spacing, application rates given may be concentrated in the row, but shall never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Telone C-17 ^{2,3}	10.8 to 17.1 gal	31.8 to 50.2	For any row spacing, application rates given may be concentrated in the row, but shall never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Telone C-35 ^{2,3}	13 to 20.5 gal	38 to 60	For any row spacing, application rates given may be concentrated in the row, but shall never exceed labeled maximum for broadcast applications. Consult the product label for additional detail.
Pic-Clor 60	19 to 31.5 gal	57 to 90	For any row spacing, application rates should never exceed the labeled maximum for broadcast applications. Consult the product label for additional detail.
Vapam HL	75 gal	-	For drip or in-row chisel fumigation, consult product label for proportionately reduced overall rates, drip concentration and flow modifying directions and procedures.
KPam HL	60 gal	-	For drip or in-row chisel fumigation, consult product label for proportionately reduced overall rates, drip concentration and flow modifying directions.
Allyl Isothiocyanate (AITC) Dominus	40 gal	-	For drip or in-row fumigation and crop termination, consult product label for overall rates, drip concentration and flow modifying directions

¹ Gallons /acre and Fl oz / 1000 feet provided only for mineral soils. Higher rates may be possible for heavier textured (loam, silt, clay) or highly organic soils.

² All of the fumigants mentioned are for retail sale and use only by state certified applicators or persons under their direct supervision. New supplemental labeling for the Telone products must be in the hands of the user at the time of application. See new label details for additional use restrictions based on soil characteristics, buffer zones, requirements for Fumigant Management Plans (FMP) and Personal Protective Equipment (PPE), mandatory good agricultural practices (GAP's), product and applicator training certification, and other use and rate modifying recommendations and restrictions.

³ Higher application rates are possible in the presence of cyst-forming nematodes.

Rates are believed to be correct for products named, and similar products of other brand names, when applied to mineral soils. Higher rates are required for muck (organic) soils. However, the **grower** has the final responsibility to see that each product is used legally; **read the label** of the product to be sure that you are using it properly.