

2018–2019 Florida Citrus Production Guide: Soft-Bodied Insects Attacking Foliage and Fruit¹

P. A. Stansly, J. A. Qureshi, L. L. Stelinski, L. M. Diepenbrock, and M. E. Rogers²

This section is focused on sucking insects which affect foliage, twigs, and fruit of citrus in Florida. Insects covered here include scales, mealybugs, whiteflies, and aphids that can impact the health of young and mature trees and fruit quality. They differ from each other in their biology, generation times, and injury to plant, but approaches to monitoring and management are similar. Individual discussions of some groups are provided, and the tables of management options are organized by active ingredient, with the target pests from this chapter in bold text. Information on the Asian citrus psyllid and Citrus Leafminer can be found in a separate chapter of the Pest Management Guide ([ENY-734](#)).

Scale Insects

There are two major groups of scale insects, soft scales and armored scales. Soft scales generally attain a larger size than armored scales and are somewhat mobile as nymphs. Nymphs and adult female armored scales are completely sessile and adult males of both are tiny gnat-like insects with a single pair of wings. The cover and body of soft scales is attached whereas the cover can be removed from the armored scale, revealing the round body underneath.

Another important distinction is that armored scales produce no honeydew whereas soft scales produce copious amounts of honeydew that attracts ants and serves as substrate for sooty mold which often accumulates on foliage below the infestation.

The most important soft scale species in Florida citrus is probably the Caribbean black scale (*Saisseta neglecta*), followed by green and brown scales (*Coccus viridis* and *C. hesperidium*, respectively), and Florida wax scale (*Ceroplastes floridensis*). Mature females are usually found on scaffold limbs, especially of young trees. Mobile first instars, “crawlers”, move out toward the outer canopy and successive nymphal stages gradually migrate inward.

The most important *armored scale* pests in Florida are snow scale (*Unaspis citri*), Florida red scale (*Chrysomphalus aonidium*), purple scale (*Lepidosaphes beckii*), Glover’s scale (*Lepidosaphes gloveri*), and chaff scale (*Parlatoria pergandii*). Snow scale tends to infest the trunk and scaffold limbs, especially of grapefruit. The name refers to the male nymphs which are white, numerous, and indistinguishable

1. This document is ENY-604, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date December 01, 1995. Revised September 2013, April 2016, and May 2018. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This publication is included in SP-43, 2018–2019 *Florida Citrus Production Guide*. For a copy of this handbook, request information on its purchase at your local UF/IFAS Extension office.
2. P. A. Stansly, professor, Department of Entomology and Nematology, Southwest Florida Research and Education Center; J. A. Qureshi, assistant professor, Department of Entomology and Nematology, Indian River REC; L. L. Stelinski, associate professor, Department of Entomology and Nematology, Citrus REC; L. M. Diepenbrock, assistant professor, Department of Entomology and Nematology, Citrus REC; and M. E. Rogers, professor, Department of Entomology and Nematology, CREC; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition. Use pesticides safely. Read and follow directions on the manufacturer’s label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county’s UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

from males of the related species, lesser snow scale, *Pinaspis strachani*, and fern scale, *P. aspidistrae*. Female snow scales are relatively large, oyster shape, purple in color with a median ridge. Lesser snow scale generally inhabits smaller limbs and fern scale make small round colonies of males with a single female off to the side on leaves and fruit. Florida red scale typically inhabits fruit and leaves while purple, Glover's, and chaff scale may be found in any part of the canopy.

Pest management of both armored and soft scale insects in Florida citrus is based on highly successful action of native and introduced natural enemies, including predators, especially ladybeetles, parasitic wasps, and fungal pathogens. These relatively specific natural enemies co-exist with their hosts in the citrus grove under most conditions and can respond to suppress pest numbers when they periodically increase in individual groves. Thus, scale insects should not be considered key pests in development of seasonal pesticidal programs. However, there are conditions under which natural enemies may not function well. It is in these cases that scale insects achieve importance in an overall Integrated Pest Management program. Factors that are most often responsible for increases in scale populations are: a) weather conditions that disrupt biological control; b) movement of the pest to groves where natural enemies do not occur; and c) disruption of natural enemies by other practices, particularly repeated use of non-selective insecticides during a period when natural enemies are active and exposed. When these disruptions occur, scale populations can increase sufficiently to damage leaves, fruit, twigs, branches, and/or trunks. The sessile nature of scale insects promotes high concentrations of scales in limited areas within the grove, thus building populations can go unnoticed for several generations. Generation times for most scale species require more than one month to progress from egg to adult. Thus, populations do not build quickly like some other pest groups such as mites or aphids.

The first consideration for management should be to determine if the problem is induced by management practices and can be solved by changing those practices. For example, if repeated applications of non-selective pesticides are responsible for scale population increase, then the solution is to desist and allow natural enemies to recover. If, on the other hand, seasonal fluctuations have brought about population levels of concern, then some intervention with insecticides may be required. The basis for this decision should be population levels of live scales that are deemed sufficient to cause damage directly or through the production of honeydew and sooty mold (soft scales only). Scale

bodies from previous generations often remain on the plant for several months and may be mistaken for live scales, resulting in the application of pesticides at inappropriate times. For effective suppression, most scale species should be in young nymphal stages at the time of application, since pesticides are not very effective against eggs, large nymphs, or adults. No economic injury levels or thresholds are available for scale insect pests in Florida. Thus, the manager must evaluate each situation, considering the intensity and extent of scale populations and how much damage is likely to result. Generally, the intent of spraying for scale insects is to reduce populations with a single application such that no additional sprays are necessary during that season and disruption to biological control is minimized.

Treatment, when warranted, should focus on selection of an appropriate material (see Table 1), but equally important, should be applied with thorough coverage in mind. Since scale insects are completely or largely immobile; direct contact is essential.

Spray volume, ground speed, nozzling, and location of the pest populations should all be emphasized to get maximum target coverage. If only a few trees are involved, then spot treatment with a handheld sprayer or other focused application equipment will provide best results. Generally, spray applications designed for contact with pests on the outer canopy are not effective at suppressing scales, especially if the scales are numerous in the interior of the tree. The follow-up to pesticidal applications for scale insects should involve evaluation of live scale on the appropriate parts of the tree. Dead scale will not be visibly different from live scale at first. Hatching crawlers will also create the impression that the spray was not effective. Complete elimination of scale insects following a pesticidal spray is neither practical nor necessary, and in fact may be counterproductive.

Soft scales are generally not pests needing treatment. Following mild winters and when populations build within specific groves, treatment, where needed, should be based on scouting for crawlers and young nymphs during the generation that develops in April–May. Applications at other times are ineffective.

Citrus snow scale likewise is a local problem requiring occasional treatment in specific groves or portions of groves. Evidence for the need to treat includes high populations of crawlers showing on patches of bark that have been brushed clean during the previous week, and the association of visible snow scale populations with bark splitting, particularly on young trees that are rapidly increasing in trunk girth.

Spot treat wood of heavily infested trees to runoff with a handheld sprayer.

Mealybugs

Citrus mealybugs (*Planococcus citri*) are normally under good biological control by a complex of natural enemies in citrus. However, intensive spraying for psyllid control may have disrupted this control. Their waxy covering, sedentary lifestyle, and preference for feeding in concealed locations make them very difficult to kill with insecticides. Only the most toxic materials have appreciable efficacy against mealybugs, materials that also pose risks to the environment and are most likely to disrupt biological control of other pests. Consequently, treatment is warranted only in cases of severe infestations, or when the fruit itself is attacked. Systemic materials give superior control while minimizing impacts on beneficials but may not act quickly enough to prevent damage when high populations are established.

Whiteflies

The most important whiteflies in Florida are citrus whitefly (*Dialeurodes citri*), the cloudy-winged whitefly (*Singhiella* (= *Dialeurodes*). *citrifolii*), the woolly whitefly (*Aleurothrixus floccosus*), and citrus blackfly (*A. woglumi*). Whiteflies are dependent on new growth for their development and reproduction; consequently, they are active in citrus only during periods of flush. Large populations of these insects can deposit considerable volumes of honeydew, leading to sooty mold accumulation. These insects are constantly present in most groves in very low numbers and are normally under good biological control by various specialist parasitoids and generalist predators. Populations are rarely high enough to warrant treatment unless biological control has been disrupted. Serious infestations of whiteflies are an indication that management practices should be reviewed.

Aphids

The most common aphids in Florida citrus are the green citrus aphid (*Aphis spiraecola*) and the cotton or melon aphid (*A. gossypii*). The green citrus aphid is responsible for curling of young flush due to feeding injury. This aphid and the melon aphid attack many different plant species and migrate into citrus mostly in spring. The brown citrus aphid (*Toxoptera citricida*) is the most important vector of citrus tristeza virus (CTV) responsible for quick decline of trees on sour orange rootstock which often die suddenly with fruit still attached. Brown citrus aphid has a narrow host range restricted largely to Rutaceae, particularly citrus, and have now become rare in Florida, possibly due

to intensive spraying for psyllids. However, melon aphid is also a vector of CTV and is also dark in color, but mottled, distinguishing it morphologically from brown citrus aphid. Aphids are dependent on the availability of newly expanding leaves for their development and reproduction, so these insects may be problems during periods of new citrus growth, primarily on young trees in spring and fall. Aphids are largely controlled by many generalist natural enemies such as ladybeetles, hoverflies, and lacewings, that normally maintain their populations, and those of other flush-feeding insects, below levels that warrant treatment in producing groves. Excessive honeydew accumulation on leaves will result in the growth of sooty mold fungus that blocks light and reduces photosynthetic activity. However, mature groves sustain little damage and should not need treatment. Treatment is warranted only in young groves (< 3 yrs old) if a large portion (i.e. > 50%) of expanding terminals are infested. Surveys for aphids should be conducted early in flushing cycles when most terminals are still in the feather stage. Systemic materials, such as Admire, applied to the soil will give good control with minimal impact on beneficial species, but the time required for uptake of these materials by the tree restricts their usefulness as preventive, rather than responsive, treatments.

Recommended Chemical Controls

READ THE LABEL. Some product labels specify rates per acre, while others specify rates per volume delivered (e.g. per 100 gallons). Refer to label for details on how product should be mixed for desired targets.

See Table 1.

Rates for pesticides are given as the maximum amount required to treat mature citrus trees unless otherwise noted. To treat smaller trees with commercial application equipment including handguns, mix the per-acre rate for mature trees in 250 gallons of water. Calibrate and arrange nozzles to deliver thorough distribution and treat as many acres as this volume of spray allows.

Table 1. Recommended chemical controls for scale insects¹, mealybugs, whiteflies, and aphids.

IRAC MOA ¹	Pesticide Trade name	Rate/Acre ²	Comments	Pests controlled
1B	Chlorpyrifos			
	Lorsban 4E	5 pt	Restricted Use Pesticide. Highly toxic to bees, do not apply during bloom.	Aphids Mealybugs , Psyllids, orange-dog, katydids, grasshoppers, thrips
1B	Dimethoate			
	various products	see label		Aphids, scales , psyllids
4A	Clothianidin (soil drench)			
	Belay 50 WDG	3.2-6.4 oz	For use on non-bearing trees only, do not apply within 1 year of fruit harvest. Do not exceed 12.8 oz/A (0.4 lb AI/A) of Belay 50 WDG per acre per year. Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period.	Aphids , psyllids, citrus leafminer
	Belay Insecticide	3-12 fl oz	Refer to the Section 18 Emergency Exemption label issued by the Florida Department of Agriculture and Consumer Services for application directions of this product to bearing citrus trees. For bearing trees, do not apply more than 12 fl oz per acre per application and do not apply more than 24 fl oz per acre in a 12 month period.	
4A	Imidacloprid			
	Various products, 2F, 4F and 4.6F	see label	Limit of 0.5 lb AI/A per growing season regardless of application type (soil and/or foliar) and trade name of imidacloprid product used.	Aphids, Mealybugs Scale Insects Whiteflies , Psyllids, Citrus leafminer (soil only)
	Foliar Application	Half to full rate	Do not apply during bloom or within 10 days of bloom or when bees are actively foraging.	
	Soil Application	Half to full rate	8 fl oz of Admire Pro 4.6F) per acre per 12 months when applied to soil. Do not exceed 0.5 lb/AI per application. See SLN for additional information.	
4A	Thiamethoxam			
	Actara (foliar application)	4.0-5.5 oz	Do not exceed a total of 11.0 oz/A (0.172 lb AI/A) of Actara or 0.172 lb a.i. of thiamethoxam-containing products per acre per growing season. Do not apply during pre-bloom or during bloom when bees are actively foraging.	Aphids, Mealybugs Scale Insects Whiteflies , Psyllids
	Platinum 75 SG (soil drench)	1.83-3.67 oz	Do not exceed a total of 3.67 oz/A (0.172 lb AI/A) of Platinum 75 SG or 0.172 lb a.i. of thiamethoxam-containing products per acre per growing season. Do not apply during pre-bloom or during bloom when bees are actively foraging.	Aphids, Mealybugs Scale Insects Whiteflies , Psyllids, Citrus leafminer
16	Buprofesin			
	Applaud	1-2 fl oz	Apply for scale insects when crawler emergence is heavy.	Mealybugs Scale insects Whiteflies

IRAC MOA ¹	Pesticide Trade name	Rate/ Acre ²	Comments	Pests controlled
23	Spirotetramat		Only controls psyllid nymphs, not adults. Limit of 0.32 lb ai per acre per 12 months. Minimum interval of 21 days between applications	
	Movento 240	10 fl oz + 3% v/v	Do not make more than one application during primary citrus bloom period. Recommended to be applied in 2% horticultural mineral oil	Aphids, Mealybugs Scale Insects Whiteflies, Citrus rust mites, psyllids
	Movento MPC	16 fl oz + 3% v/v	Do not apply within 10 days prior to bloom, during bloom, or until petal fall is complete. Recommended to be applied in 2% horticultural mineral oil	
29	Floconamid			
	Beleaf 50SG	2.8 oz	Apply when populations begin to build	Aphids
UN³	Horticultural Mineral Oil			
	97+% (FC 435-66, FC 455-88, or 470 oil)	5 gal	Do not apply when temperatures exceed 94°F. 470 weight oil has not been evaluated for effects on fruit coloring or ripening. These oils are more likely to be phytotoxic than lighter oils.	Aphid, Scales leafminer, citrus rust- mite, aphids, scales

¹ Mode of action class for citrus pesticides from the Insecticide Resistance Action Committee (IRAC) Mode of Action Classification V.8.4 (2018).

² Lower rates may be used on smaller trees. Do not use less than the minimum label rate.