

2018–2019 Florida Citrus Production Guide: Postbloom Fruit Drop¹

N. A. Peres and M. M. Dewdney²

Postbloom fruit drop (PFD) must be controlled on processing and fresh market fruit. PFD, caused by the fungus *Colletotrichum acutatum*, affects all species and cultivars of citrus, but severity on a given cultivar varies according to the time of bloom in relation to rainfall. Navel and Valencia oranges have experienced the most severe damage in Florida since they tend to have extended or multiple blooms.

Most spores of this fungus are produced directly on the surface of infected petals. Spores are splash-dispersed by rain to healthy flowers where they infect within 24 hours and produce symptoms in 4–5 days. The fungus survives between bloom periods as resistant structures on the surface of leaves, buttons, and twigs. Flowers are susceptible from the button stage (with white tissue present) until they are open.

Groves with persistent calyxes (buttons) from the previous year should be closely examined once the bloom begins. If infected flowers are present on scattered early bloom, fungicide application recommendations should be followed once sufficient bloom is present for the fungicide application to be economical. Groves with a history of PFD should be checked twice weekly during the bloom period. Ground and aerial applications are effective for control of PFD. Low

volume application equipment can be used provided they give good coverage of the flowers, but ensure that you are following the minimum volume levels on the label as it varies among products. The removal of declining trees such as those with HLB, blight or phytophthora, where off-season blooms may provide a site for fungal spore buildup, and a reduction in overhead irrigation during bloom should reduce disease severity.

A fungicide application decision system, called the PFD-FAD system, has been developed and is available at <http://pfd.ifas.ufl.edu/>. The system takes into consideration the disease history in the grove, the cultivar susceptibility, the weather conditions (rainfall and leaf wetness duration), and the time of the last fungicide application. The PFD-FAD system is easy to use but should be checked daily and fungicides applied within 48 hours after the recommendation. It is also available in Spanish and Portuguese.

More recently, the Citrus Advisory System (CAS; Figure 1) was developed to facilitate even further growers decision on the need for fungicide applications. The system uses real-time weather data from Florida Automated Weather Network (FAWN) stations throughout the state to determine if risk conditions for PFD development are low (green), moderate (yellow), or high (red). Specific fungicide

1. This document is PP-45, one of a series of the Plant Pathology Department, UF/IFAS Extension. Original publication date December 1995. Revised September 2013, April 2016, and May 2018. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

2. N. A. Peres, professor, Plant Pathology Department, UF/IFAS Gulf Coast Research and Education Center; and M. M. Dewdney, associate professor, Plant Pathology Department, UF/IFAS Citrus REC; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. It is not a guarantee or warranty of the products named, and does not signify that they are approved to the exclusion of others of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

spray recommendations are given according to the disease risk conditions. CAS is available at <http://agroclimate.org/tools/cas>. If desired, notifications can be sent via SMS or e-mail for an alert to check the model because an infection event has occurred.




Figure 1. Citrus PFD Advisory System available at <http://agroclimate.org/tools/cas>.

Preventive fungicide programs are difficult to implement especially in groves with large number of declining trees, which might bloom for an extended period since the number of fungicides available as well as the number of applications for each fungicide is limited. As a reminder, groves with a history of PFD should be checked twice weekly during the bloom period. If symptomatic flowers are found on scattered bloom, PFD-FAD or CAS recommendations should be followed once sufficient bloom is present to justify a fungicide application.

Of the products recommended for control of PFD, the strobilurin-containing fungicides Abound, Amistar Top, Gem, Headline, Priaxor, and Pristine are effective but do not have a long residual effect. Ferbam is less effective and should not be used alone but can be combined with low rates of other products to maximize protection and reduce the risk of resistance development. No resistance has been detected to date. The strobilurin-containing fungicides should not be used alone more than once per season, but can be used more than once if combined with Ferbam.

Recommended Chemical Controls

READ THE LABEL.

Rates for pesticides in are given as the maximum amount required to treat mature citrus trees unless otherwise noted. To treat smaller trees with commercial application equipment including handguns, mix the per acre rate for mature trees in 125 gallons of water. Calibrate and arrange nozzles to deliver thorough distribution and treat as many acres as this volume of spray allows.

Table 1. Recommended chemical controls for Postbloom Fruit Drop.

Pesticide	FRAC MOA ²	Mature Trees Rate/Acre ¹
Ferbam Granuflo	M03	5–6 lb. Maximum 3 ferbam applications a year and do not apply more than 6 lb ai/acre in a single application.
Abound ³	11	12.0–15.5 fl oz. Do not apply more than 92.3 fl oz/acre/season for all uses.
Abound ³ + Ferbam	11, M03	12.0 fl oz + 5 lb. Do not apply more than 92.3 fl oz/acre/season of Abound for all uses. Maximum 3 ferbam applications a year and do not apply more than 6 lb ai/acre in a single application.
Amistar Top (formerly Quadris Top) ^{3,4}	11 + 3	15.4 fl oz. Do not apply more than 61.5 fl oz/acre/year. Do not apply more than 0.5 lb ai/acre/season difenoconazole. Do not apply more than 1.5 lb ai/acre/season azoxystrobin.
Gem 500 SC ³	11	1.9–3.8 fl oz. Do not apply more than 15.2 fl oz/acre/season for all uses. Do not apply within 7 days of harvest.
Gem ³ + Ferbam	11, M03	1.9 fl oz + 5 lb. Do not apply more than 15.2 fl oz/acre/season of Gem for all uses. Do not apply within 7 days of harvest. Maximum 3 ferbam applications a year and do not apply more than 6 lb ai/acre in a single application.
Headline SC ³	11	12–15 fl oz. Do not apply more than 54 fl oz/acre/season for all uses.
Headline ³ + Ferbam	11, M03	12 fl oz + 5 lb. Do not apply more than 54 fl oz/acre/season of Headline for all uses. Maximum 3 ferbam applications a year and do not apply more than 6 lb ai/acre in a single application.
Pristine ^{3,4}	11 + 7	16–18.5 oz. Do not apply more than 74 oz/acre/season for all uses.
Priaxor ^{3,4}	11 + 7	9–11 fl oz. Do not apply more than 44 fl oz/acre/year.

¹ Lower rates can be used on smaller trees. Do not use less than the minimum label rate.

² Mode of action class for citrus pesticides according to the Fungicide Resistance Action Committee (FRAC) 2018. Refer to ENY624, Pesticide Resistance Management, in the *2018–2019 Florida Citrus Pest Management Guide* for more details.

³ Do not use more than 4 applications of strobilurin fungicides/season. Do not make more than 2 sequential applications of strobilurin fungicides.

⁴ Do not make more than 4 applications of Pristine, Amistar Top, or Priaxor/season. Do not make more than 2 sequential applications of Pristine, Amistar Top, or Priaxor before alternating to a non-strobilurin, SDHI, or DMI.