



UNIVERSITY OF
FLORIDA

EENY-129

EXTENSION

Institute of Food and Agricultural Sciences

Sweetpotato Whitefly B Biotype of Silverleaf Whitefly, *Bemisia tabaci* (Gennadius) or *Bemisia argentifolii* Bellows & Perring (Insecta: Homoptera: Aleyrodidae)¹

Heather J. McAuslane²

Introduction

The sweetpotato whitefly, *Bemisia tabaci* (Gennadius), has been recorded in the United States since the late 1800s. Before 1986, it was only an occasional pest of cultivated crops. However, in 1986 in Florida, the insect we called *B. tabaci* became an extreme economic pest. It was found attacking crops that it had not infested previously, such as poinsettia, and was resistant to many formerly effective insecticides. It transmitted new plant-pathogenic viruses that had never affected cultivated crops and induced plant physiological disorders, such as tomato irregular ripening and squash silverleaf disorder. This whitefly was thought to be a new biotype of *B. tabaci* (designated variously as B biotype, Florida biotype, or poinsettia biotype), possibly introduced from the Middle East. It quickly moved into other southern states with intensive agricultural and horticultural industries (Texas, Arizona, and California), and displaced the original *B. tabaci* (A biotype), which can no longer be found in the United States. The B biotype was described in 1994 as a new species,

Bemisia argentifolii Bellows & Perring. Its common name is the silverleaf whitefly, because of its unique ability to induced silverleaf disorder in squash. The designation of this insect as a new species is not universally accepted and I will refer to *B. tabaci* B biotype (= *B. argentifolii*) as *Bemisia*.

Distribution

Bemisia is primarily a pest of cultivated plants in tropical and warm temperate regions of the world. It is found throughout the southern United States and can overwinter outdoors as far north as South Carolina. It is found infesting greenhouses in more northern latitudes in the United States and Canada. It is widely distributed throughout the Caribbean Islands, Central and South America, and Mexico. It is present throughout much of southern Europe, Africa, India, and has recently moved into Australia.

1. This document is EENY-129, one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: April 2000. Revised: November 2002. This document is also available on Featured Creatures Website at <http://creatures.ifas.ufl.edu>. Please visit the EDIS Website at <http://edis.ifas.ufl.edu>. Additional information on these organisms, including many color photographs, is available at the Entomology and Nematology Department website at <http://entnemdept.ifas.ufl.edu/> and the Division of Plant Industry website at <http://www.doacs.state.fl.us/~pi/>.
2. Heather J. McAuslane, associate professor, Entomology and Nematology Department, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.

The Institute of Food and Agricultural Sciences is an equal opportunity/affirmative action employer authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, sex, age, handicap, or national origin. For information on obtaining other extension publications, contact your county Cooperative Extension Service office. Florida Cooperative Extension Service/Institute of Food and Agricultural Sciences/University of Florida/Christine Taylor Waddill, Dean.



Figure 1. 'Florida Lanai' cherry tomato showing internal symptoms of *Bemisia*-induced tomato irregular ripening disorder (bottom) and control fruit from uninfested plant (top). (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Shahab Hanif-khan, University of Florida



Figure 2. 'Zucchini Elite' showing symptoms of *Bemisia*-induced squash silverleaf disorder. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Yasmin Cardoza, University of Florida

Description and Life History

Infestations of whiteflies, especially in greenhouses, may occasionally be a mixture of *Bemisia* and *Trialeurodes vaporariorum* (Westwood), the greenhouse whitefly. They can be differentiated at the adult stage based on the position in which the wings are held over the body while alive (see example in Whitefly Knowledgebase), close to the body and tent-like in *Bemisia*, and more loosely in *T. vaporariorum*. Another way to differentiate adult *Bemisia* and *T. vaporariorum* is to examine the compound eyes using a microscope. The upper and

lower compound eyes of *T. vaporariorum* are completely divided whereas they are joined by one ommatidium in *Bemisia*. Adult *Bemisia* are soft and whitish-yellow when they first emerge from their nymphal exuvia. Within a few hours, their two pairs of wings become iridescent white due to the deposition of a powdery wax. The body remains light yellow with a light dusting of wax. The body of the female measures 0.96 mm from the tip of the vertex (head) to the tip of the abdomen, while the male is somewhat smaller at 0.82 mm.



Figure 3. Adult *Bemisia argentifolii* Bellows & Perring or *B. tabaci* (Gennadius), B strain. Credits: Scott Bauer, USDA

Adult whiteflies emerge through a T-shaped slit in the integument of the last nymphal instar. The remaining white, transparent "shell" is called the exuvia. If the exuvia has a round hole in it rather than a slit, an adult parasitoid emerged. Adults that emerge may simply fly up the same plant or over to another plant. These are called trivial flights. Some individuals however, are primed for short-distance migration of up to several kilometers. Migrating individuals usually develop on plants that are senescing. These migrations can often be massive and can lead to severe infestation of newly planted crops.

As far as we know, the sexes do not rely on pheromones to locate one another. Mating can occur as soon as the whiteflies have expanded and hardened their wings, usually within a few hours. *Bemisia* has arrhenotokous parthenogenetic reproduction where virgin females can only lay haploid eggs which give rise to males. Mated females can produce haploid and diploid (female) eggs. There is usually a pre-oviposition period of less than a day to a few




Figure 4. Exuvia of *Bemisia* with characteristic T-shaped whitefly adult emergence slit (left), and exuvia of *Bemisia* with round emergence hole through which an adult parasitoid emerged (right). (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Jane C. Medley, University of Florida

days, depending on the temperature. The female lands on a plant, using visual stimuli such as plant color, and walks or flies to the lower surface of a leaf. She tests the suitability of the plant by probing the leaf with her piercing-sucking mouthparts and ingesting a small amount of sap. Chemoreceptors on the tip of her mouthparts and within her mouthparts sense the leaf's chemical composition. If the host plant is deemed acceptable, she will insert her mouthpart stylets into the phloem from where she sucks plant sap. While she is feeding she may lay eggs, often in a semi-circular arrangement as she swivels her body around her feeding site. Female longevity can range from 10 to 24 days during which time she can lay between 66 and 300 eggs, depending on host plant and temperature.

Whitefly eggs are oval in shape and somewhat tapered towards the distal end. The broader end has a short stalk, 0.024 mm, that is inserted by the ovipositing female into the leaf. The egg obtains moisture through this stalk. The egg is approximately 0.21 mm in length and 0.096 mm in width. The egg is pearly white when first laid but darkens over time. The distal end of the egg becomes dark brown just before the first nymphal instar ecloses. At 25° C, the eggs will hatch in six to seven days.

The first nymphal instar is capable of limited movement and is called the crawler. It is oval in shape and measures approximately 0.27 mm in length and 0.14 mm in width. The dorsal surface of the crawler



Figure 5. Newly laid eggs of *Bemisia* are pale yellow while those about to hatch are dark brown. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: James Castner, University of Florida

is convex while the ventral surface, appressed to the leaf surface, is flat. The crawler has three pairs of well-developed four-segmented legs, three-segmented antennae, and small eyes. It is whitish-green in color and has two yellow spots, the mycetomes, visible in the abdomen through the integument (skin). The mycetomes house several species of endosymbiotic bacteria that may play an important role in whitefly nutrition. The crawlers usually move only a few centimeters in search of a feeding site but can move to another leaf on the same plant. They initiate feeding on the lower surface of a leaf, also feeding in the phloem. After they have begun feeding, they will molt to the second nymphal instar, usually two to three days after eclosion from the egg.

The second, third and fourth nymphal instars are immobile with atrophied legs and antennae, and small eyes. The nymphs secrete a waxy material at the margins of their body that helps adhere them to the leaf surface. Nymphs are flattened and oval in shape, greenish-yellow in color, and range from 0.365 mm (second instar) to 0.662 mm (fourth instar) in length. The mycetomes are yellow. The second and third nymphal instars each last about two to three days.

The red-eyed nymphal stage is sometimes called the "pupal stage". There is no molt between the fourth nymphal instar and the red-eyed nymphal stage but there are morphological differences. The fourth and red-eyed nymphal stages combined lasts for five to




Figure 6. Various nymphal instars of *Bemisia*. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: James Castner, University of Florida

six days. The stage gets its name from the prominent red eyes that are much larger than the eyes of earlier nymphal instars. The red-eyed nymphal stage is also less flattened and more convex in shape. This stage is more yellow than the fourth instar and the mycetomes are less visible. The red-eyed nymphal stages of *Bemisia* and *T. vaporariorum* can be distinguished easily in a mixed infestation. The sides of *Bemisia* red-eyed nymphs are boat-shaped or wedged (see example in Whitefly Knowledgebase) while those of *T. vaporariorum* are perpendicular to the leaf surface.



Figure 7. Red-eyed nymphal or "pupal" stage of *Bemisia*. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: James Castner, University of Florida

Hosts

Bemisia is widely polyphagous, feeding on over 500 species of plants in 74 families. Its hosts include

vegetable, field, and ornamental crops. Of the important vegetable crops grown in Florida, *Bemisia* is a major pest of tomato, peppers, squash, cucumber, beans, eggplant, watermelon, and cabbage. The Florida-grown field crops of potato, peanut, soybean and cotton are heavily attacked by *Bemisia*. The ornamental host plants of *Bemisia* are too numerous to list, but include poinsettia, hibiscus, and chrysanthemum.

Damage

Bemisia can cause economic damage to plants in several ways. Heavy infestations of adults and their progeny can cause seedling death, or reduction in vigor and yield of older plants, due simply to sap removal. When adult and immature whiteflies feed, they excrete honeydew, a sticky excretory waste that is composed largely of plant sugars. The honeydew can stick cotton lint together, making it more difficult to gin and therefore reducing its value. Sooty mold grows on honeydew-covered substrates, obscuring the leaf and reducing photosynthesis, and reducing fruit quality grade.



Figure 8. Sooty mold developing on soybean leaves covered with *Bemisia* honeydew. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: James Castner, University of Florida

Feeding by immature *Bemisia*, but not adults, has been associated with several developmental physiological disorders of plants. Tomato irregular ripening was first noted in Florida tomatoes in 1987. Tomatoes that develop on plants that are heavily infested with whiteflies may incompletely develop external color, resulting in streaking. Even if fruits appear normal externally, the internal tissue may be

white, hard, and unripe. It has been estimated that, in 1989, Florida tomato growers lost \$25 million to tomato irregular ripening. Squash silverleaf disorder is another developmental disorder caused by feeding of immature whiteflies, also first noted in Florida in 1987. This disorder affects many *Cucurbita* species, including the squashes and pumpkins of *C. pepo*, *C. moschata*, and *C. mixta*. Feeding by immature whiteflies causes newly developing leaves, but not the leaves on which they are feeding, to take on a silvery appearance due to the separation of the upper epidermis from the underlying cell layer. The resultant air space reflects light, causing the silvery color. Fruits that develop on silvered plants may be bleached, and are of lower quality grade. Other physiological disorders caused by *Bemisia* include lettuce leaf yellowing and stem blanching, carrot light root, pepper streak, *Brassica* white stem, and chlorosis of new foliage of many plants.




Figure 9. 'Florida Lanai' cherry tomato showing external symptoms of *Bemisia*-induced tomato irregular ripening disorder (left) and control fruit from uninfested plant (right) (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Shahab Hanif-khan, University of Florida

Bemisia vectors several serious plant-pathogenic geminiviruses in the United States. In Florida, our main concerns are the tomato yellow leaf curl virus (TYLCV), tomato mottle virus (TMoV), and bean golden mosaic virus (BGMV). Advice for home gardeners (see Florida Pest Alert for more details) to manage TYLCV includes destroying and disposing of symptomatic tomato plants so that they cannot become a source of inoculum for healthy plants, and managing *Bemisia* populations with insecticides, if necessary.



Figure 10. Poinsettia showing chlorosis of new foliage that developed during heavy feeding by immature *Bemisia*. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring).



Figure 11. Tomato foliage showing characteristic yellowing and leaf curling associated with infection by *Bemisia*-vectored tomato yellow leaf curl virus. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring).

Cultural Control

Several cultural controls can reduce *Bemisia* populations and lessen their impact on crops. One important control method is sanitation. During the growing season, geminivirus-infected plants can be rogued out and destroyed. After harvest, crop residue (tomato plants, in particular) should be removed so that there is no plant material for large numbers of whiteflies to carry over the crop-free period, usually the height of summer.

Israeli researchers have had success with the use of barriers to keep viruliferous *Bemisia* from invading greenhouses. Greenhouses are screened

with very fine mesh plastic screen. Ventilation must be increased however, to reduce the likelihood of infection by plant pathogens. Whitefly infestations have also been reduced with the use of UV-absorbing greenhouse plastic films. Whiteflies do not enter greenhouses or areas covered with this type of plastic as frequently as they do greenhouses covered in non-UV-absorbing material.

Colored plastic mulches may be effective in reducing whitefly populations and geminivirus incidence. In Florida, tomato plants mulched with yellow or aluminum plastic mulches yielded more and had less tomato mottle virus infection than those planted on white or black plastic mulches. In Costa Rica, living mulches (e.g., perennial peanut and cilantro) may reduce somewhat the spread of geminivirus within tomato fields.

Host plant resistance offers limited hope for whitefly management. Tomato and bean breeders are attempting to develop varieties with resistance to geminiviruses, but these varieties are still some time from commercialization. No varieties of host plants have been found to be highly resistant to whiteflies themselves, however, some plant factors are not preferred by whiteflies. For example, smooth-leaved varieties of cotton and soybean are less preferred by ovipositing female *Bemisia* than are hairy-leaved varieties. This holds true for many plant species. Glossy (less waxy) crucifers, such as broccoli and collard, are less acceptable for oviposition than are varieties with a normal wax layer.

Biological Control

In Florida, *Bemisia* rarely reaches outbreak population levels in natural environments or in agricultural situations where no insecticides are applied. In these situations, natural biological control keeps *Bemisia* below economically injurious levels. Much mortality is caused by minute parasitic wasps (parasitoids) in the aphelinid family. Female parasitoids lay their eggs inside the whitefly nymph or between the whitefly and the leaf surface, depending on the genus of wasp. The immature parasitoids develop within the whitefly host, eventually consuming the entire host, except the integument. The immature parasitoid pupates within

the integument of the host and the adult parasitoid emerges through a round hole. The most common parasitoids attacking *Bemisia* in Florida are in two genera, *Encarsia* and *Eretmocerus*. *Encarsia pergandiella* (Howard) and *Encarsia transvena* (Timberlake) are common throughout the state, while *Encarsia nigricephala* Dozier is common in north central Florida. Several species of *Eretmocerus* are also common throughout the state. *Eretmocerus* species cause mortality to whiteflies by host-feeding, in addition to parasitism. Females stab the immature whitefly repeatedly with their ovipositor and then turn around and feed from the wound, obtaining valuable protein with which to provision their eggs.



Figure 12. Parasitoid in the *Eretmocerus* genus parasitizing a *Bemisia* nymph (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits : Heather McAuslane, University of Florida



Figure 13. Pupa of *Encarsia pergandiella* within body of *Bemisia* nymph. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring).



Figure 14. Black pupal case of *Encarsia transvena* within body of *Bemisia* nymph. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: James Castner, University of Florida



Figure 15. Pupa of *Encarsia nigricephala* (left) and *Eretmocerus* sp. (right) within body of *Bemisia* nymphs. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: James Castner, University of Florida

Bemisia populations can also be reduced by predation. Many species of insects, mites, and spiders feed on immature and adult whiteflies. Some well-known generalist predators that prey on whiteflies include *Chrysoperla* species larvae (lacewings), *Orius* species (minute pirate bugs), and *Geocoris* species (big-eyed bugs). Several coccinellid species are specialist whitefly predators, such as *Delphastus catalinae* (LeConte) and *Nephapsis oculatus* (Blatchely).

Under certain climatic conditions, usually wet and warm, *Bemisia* populations can be drastically



Figure 16. Female of an *Eretmocerus* species host-feeding on *Bemisia* nymph. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Heather McAuslane, University of Florida



Figure 17. Lacewing larva (*Chrysoperla* sp.) feeding on *Bemisia* nymphs. (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Jack Dykinga, USDA

reduced by natural epizootics of entomopathogens. *Beauveria bassiana* and *Verticillium lecanii*, two fungal pathogens with efficacy against *Bemisia* and other insects, have been commercialized and are available for field and greenhouse application against *Bemisia*.

In protected plant culture, such as greenhouses, it is not possible to rely on natural biological control. Several effective species of parasitoids, such as *Eretmocerus eremicus*, and predators, such as lacewings and the coccinellid *D. catalinae*, are available commercially for control of *Bemisia*.



Figure 18. Adult minute pirate bug (*Orius* sp.) feeding on *Bemisia* nymphs (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Jack Dykinga, USDA

Chemical Control

Bemisia is difficult to control with insecticides for two reasons; adults and immatures infest the lower surfaces of leaves which are difficult to reach with insecticide sprays and they have developed resistance to many insecticide chemistries. Homeowners would be best advised to treat populations before they reach outbreak levels and to spray immature whiteflies with a mixture of insecticidal soap or insecticidal oil in water (see label for directions). Care should be taken to cover the lower surface of the leaves. Other chemical control recommendations can be found at the following sites:

[Insect Management Guide for Vegetables](#)

[Insect Management Guide for Field Crops \(includes Cotton, Peanuts and Soybean\)](#)

[Insect Management Guide for Turf and Ornamentals](#)



Figure 19. Adult bigeyed bug (*Geocoris* sp.) feeding on *Bemisia* nymphs (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Jack Dykinga, USDA




Figure 20. Adult coccinellid predator of *Bemisia* nymphs, *Delphastus catalinae* (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* Bellows & Perring). Credits: Kim Hoelmer, USDA




Figure 21. *Bemisia* nymph infected with the entomopathogenic fungus, *Paecilomyces fumosoroseus* (*Bemisia* = sweetpotato whitefly B biotype, *Bemisia tabaci* (Gennadius), or silverleaf whitefly, *Bemisia argentifolii* (Bellows & Perring). Credits: Heather McAuslane, University of Florida

Selected References

Bellows, T. S., Jr., Perring, T. M., Gill, R. J., and D. H. Headrick. 1994. Description of a species of *Bemisia* (Homoptera: Aleyrodidae). *Ann. Entomol. Soc. Am.* 87: 195-206.

Brown, J. K., D. R. Frohlich, and R. C. Rosell. 1995. The sweetpotato or silverleaf whiteflies: biotypes of *Bemisia tabaci* or a species complex? *Annu. Rev. Entomol.* 40: 511-534.

Csizinszky, A. A., D. J. Schuster, and J. B. Kring. 1997. Evaluation of color mulches and oil sprays for yield and for the control of silverleaf whitefly, *Bemisia argentifolii* (Bellows and Perring) on tomatoes. *Crop Prot.* 16: 475- 481.

Evans, G. A. and A. B. Hamon. (5 November 2002). Whitefly taxonomic and ecological Web site: an on-line interactive catalog of the whiteflies (Hemiptera: Aleyrodidae) of the world and their parasites and predators.

(
http://www.fsca-dpi.org/homoptera_hemiptera/Whitefly/whitefly_catalog.htm).

Fasulo, T. R. et al. (1995). Whitefly Knowledgebase. UF/IFAS.
<http://whiteflies.ifas.ufl.edu/wfly0002.htm> (4 April 2000).

Lopez-Avila, A. 1986. Taxonomy and biology, pp. 3 - 11. In M. J. W. Cock (ed.), *Bemisia tabaci* - A Literature Survey on the Cotton Whitefly with an Annotated Bibliography. CAB International Institute of Biological Control, Ascot, U.K.

Gerling, D. (ed.). 1990. Whiteflies: their Bionomics, Pest Status and Management. Intercept, Ltd., Andover, UK.

Gerling, D. and R. T. Mayer (eds.). 1996. *Bemisia*: 1995. Taxonomy, Biology, Damage, Control and Management. Intercept, Ltd., Andover, UK.

Hamon, A. B. and V. Salguero. 1987. *Bemisia tabaci*, sweetpotato whitefly, in Florida (Homoptera: Aleyrodidae: Aleyrodinae). Entomology Circular No. 292, Fla. Dept. Agric. and Consumer Serv., Division of Plant Industry.

Koehler, P.G., D.E.Short and T.R. Fasulo. (1998). Pests In and Around the Home. UF/IFAS. SW-126. CD-ROM.

Liu, T.-X., and P. A. Stansly. 1999. Searching and feeding behavior of *Nephaspis oculatus* and *Delphastus catalinae* (Coleoptera: Coccinellidae), predators of *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Environ. Entomol.* 28: 901-906.

Maynard, D. N. and D. J. Cantliffe. 1990. Squash silverleaf and tomato ripening: new vegetable disorders in Florida. Vegetable Crops Fact Sheet VC-37, IFAS, Univ. of Fla.

McAuslane, H. J., F. A. Johnson, D. L. Colvin, and B. Sojack. 1995. Influence of foliar pubescence on incidence and parasitism of *Bemisia argentifolii* (Homoptera: Aleyrodidae) on soybean and peanut. *Environ. Entomol.* 24: 1135-1143.

Momol, T. (4 March 1999). Tomato yellow leaf curl virus on tomato in north Florida and south Georgia. UF/IFAS Pest Alert.
<http://extlab7.entnem.ufl.edu/pestalert/tmm-0304.htm> (4 April 2000)

Momol, T. and J. Funderburk (20 August 1999). Tomato yellow leaf curl virus on tomato: update from north Florida. UF/IFAS Pest Alert.

<http://extlab7.entnem.ufl.edu/PestAlert/tmm-0820.htm> (4 April 2000)

Tsai, J. H., and K. Wang. 1996. Development and reproduction of *Bemisia argentifolii* (Homoptera: Aleyrodidae) on five host plants. *Environ. Entomol.* 25: 810-816.