



Cotton Growth and Development¹

D.L. Wright and R.K. Sprenkel²

Introduction

Management of cotton requires an understanding of the growth habit and responses of the plant to the environment. This publication provides basic information for understanding how a cotton plant develops and some of the factors that can alter the pattern of development.

Seasonal Development of the Cotton Plant

The cotton plant is a perennial in many parts of the tropics and subtropics where it may reach a height of 15 to 20 feet. In Florida and across the commercial cotton belt of the U.S., cotton is grown as an annual and attains a plant height of 2 to 5 feet or more if moisture and plant nutrients (primarily nitrogen) are not limiting. Air temperatures in the 90° to 95°F range are considered near optimum for growth. Very little growth takes place below 60°F, and temperatures above 100°F for several days may be detrimental, especially if soil moisture is low. However, cotton is usually considered drought tolerant because of its extensive root system. An average daily growth rate for the roots of 1/2 inch may occur until first flower when root growth begins to

level off and then declines starting about 90 days after planting.

Cotton fibers (lint) are produced on the seed inside the boll. The seed is also the key to obtaining good stands of cotton plants. Seeds consist of two cotyledons and an embryo. Prior to ginning and delinting, the top layer of the seed coat has two types of fiber, long lint fiber and short linters. After ginning and acid delinting, seed are treated with fungicide prior to planting. Fungicides aid in stand establishment since acid delinting leaves cracks in the seed coat, giving disease organisms an entry point into the germinating seed. Germination begins within a few hours after moisture is taken up. The cotyledons eventually form the first green leaves as well as containing stored food which supplies energy for germination and early development.

Approximately 4 to 10 days after planting, cotyledonary or seed leaves are fully expanded (Table 1). These leaves are on node number 0 and are borne on opposite sides of the main stem. The nodes above the seed leaves occur in a spiral arrangement around the stem and bear a single true leaf. At the base of each main stem leaf, in the angle between the leaf and the stem, there are two, sometimes three

-
1. This document is SS-AGR-238, one of a series of the Agronomy Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date February 2005. Reviewed March 2008. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.
 2. D. L. Wright, professor, Agronomy Department, North Florida Research and Education Center, Quincy FL; R. K. Sprenkel, professor, Entomology and Nematology Department, and Associate Director, North Florida Research and Education Center, Quincy FL; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean

axillary buds. These buds give rise to vegetative branches on the lower nodes (nodes 2 through 5 or 6). At nodes 6 or 7, and above, generally on lateral branches, are fruiting branches, which bear the floral buds, or cotton squares. If a cotton plant does not produce squares by node 9, a problem exists and its cause(s) should be determined and corrected if possible.

The time required for development from pinhead square to white bloom is approximately 23 days (Table 1). Pollination occurs on the first day the flower is open (white bloom stage) and generally early in the day. The flower turns pink (or red) the day following pollination. The interval between corresponding nodes on successive fruiting branches (vertical flowering interval) is 2 to 3 days. For example, under optimum conditions, a cotton plant having a white flower on the first lateral node of a fruiting branch will produce another white flower on the first lateral node of the next higher branch 2 to 3 days later. The interval between successive flowers on the same fruiting branch (horizontal fruiting interval) is 5 to 6 days. Following fertilization, the hollow fibers begin to lengthen and will reach their final staple length in approximately 3 weeks. For the next several weeks, the walls of the fibers thicken through the deposition of successive layers of cellulose. A graphic depiction is shown in Figure 1 (Landivar and Benedict, 1996).

Cotton is indeterminate, meaning that flowering will continue until stopped by frost, drought, insect attack or some other cause. Shedding of squares, flowers or young bolls is common. Under good conditions, only 35 to 40% of the squares normally produce mature bolls. Once bolls are 12 days old or older, they usually will not shed unless the plant suffers severe stress (temperature, moisture, insect or disease). The time required for development from the pink flower stage to the open boll stage is approximately 55 days. Cloudy weather and below-optimum temperatures increase the boll maturation period. Late in the growing season, 65-70 days are required for development from the pink flower to the open boll stage.

Normally, 100 to 120 bolls are required to produce a pound of seed cotton (160 to 170 for a

pound of lint). However, varieties that produce relatively small bolls may require more bolls to produce a pound of seed cotton. Also, bolls developing later in the season are smaller and therefore more are required to produce a pound of cotton lint. Figure 2 (Landivar and Benedict, 1996) shows a diagrammatic sketch of a cotton plant with mainstem nodes and fruiting nodes and general plant architecture.




Figure 1. Phases of cotton production in relation to time course of plant height increase and time of first square, first bloom and cutout for short-season production. Credits: Landivar and Benedict, 1996.




Figure 2. Diagrammatic sketch of cotton plant showing mainstem nodes (M) fruiting nodes (O), and general plant architecture for NAWF=5. Credits: Landivar and Benedict, 1996.

Fruit Shed

Cotton has the potential for setting about 90% of the crop in the first three weeks of blooming (Table 2). However, fruit shed usually causes the setting

period to be considerably longer. Fruit shed in cotton is a physiological process. Squares and small bolls shed because an abscission zone forms between the fruiting branch and the peduncle (boll stem). During the abscission process, enzymes loosen the connection between the cells allowing the weight of the square or boll to break the peduncle. The weakening of cells at the abscission zone is controlled by the balance of plant hormones, ethylene and abscissic acid (ABA) which promote abscission and indole acetic acid (IAA) which inhibits abscission. Because of the process involved, several days are required between the stimulus causing shed and the actual loss of fruit. Insect feeding on small squares prior to bloom may also contribute to fruit shed. Large squares, blooms and medium to large size bolls are most resistant to shed, possibly due to their high proportion of IAA (which inhibits abscission) to ethylene and ABA (which promote abscission). Conversely, small to medium-sized squares and small-sized bolls have a high proportion of ethylene and ABA to IAA and are therefore more likely to shed their bolls.

Factors That May Cause Shed

Cotton sheds fruit for a variety of reasons. Some of the more important causes for abscission which have been identified and studied are listed below.

Reduced Photosynthate Supply

Photosynthates are sugars produced through photosynthesis and used in plant growth (leaves, squares, bolls, etc.). The amount of sugars in a plant may be reduced if the supply is reduced or if the demand for the sugars increases. The supply is reduced with low light, older leaves, water and nutrient stress, foliage damage from insects and environmental events, and extreme temperatures. The demand increases with the presence of immature bolls, rank plant growth and high day and night temperatures.

Light

Sunlight is required by cotton plants to produce photosynthate. Full sunlight is required for maximum photosynthesis. During cloudy or overcast weather, photosynthesis is greatly reduced. Furthermore, the

higher temperatures of the summer increase the need for sugars which increase the amount of shed. During cloudy weather young bolls are the main fruit size to be shed.

Even with full sunlight, rank growth cotton may experience considerable self-induced fruit shed. This is because in this type of cotton, once fruit gets to the bloom or small boll stage, the leaves feeding sugar to these fruit (the leaf at the base of the fruit or one adjacent to it) are already shaded by new foliage growth at a higher level in the canopy. Loss of these fruit causes the cotton to put more sugars into leaves, stems, nodes, etc., thus perpetuating the problem.

Temperature Extremes

Cold temperatures reduce photosynthesis and sugar production resulting in shed. Generally, cotton is more tolerant of high temperatures. However, if cotton is unable to cool itself below 90°F through evaporation, shed will occur. Evaporative cooling becomes difficult if there is insufficient soil moisture and/or if the humidity is extremely high.

Another cause of boll shed is due to high nighttime temperatures when pollen sterility may occur. This type of shed occurs 17-19 days following night temperatures that remain at 85°F or above.

Soil Moisture

Excess and insufficient soil moisture are both known to cause fruit shed. In the case of excess soil moisture (to the point of saturation), oxygen levels in the soil are decreased causing stomates to close, which reduces photosynthesis and evaporative cooling, causing increased fruit shed.

An insufficient amount of soil moisture has several effects on a plant that can lead to shed. First is the inability of the plant to regulate its temperature through evaporative cooling. This occurs when a plant cannot obtain moisture from the soil. Secondly, prolonged low soil moisture levels prematurely age leaves causing a reduction in photosynthate supply and shed.

Moisture in Bloom

Another adverse effect of moisture on shed occurs when open blooms contain water (as might occur with an early-morning rainfall). Water causes pollen to rupture, thereby preventing pollination. Unpollinated flowers are shed.

Nitrogen

As with several other factors mentioned above, both insufficient and excess nitrogen can lead to fruit shed. The effect of nitrogen deficiency on fruit set is two-fold. First, nitrogen deficient plants stop developing new nodes and squares and enter premature cutout (the point at which the cotton plant stops producing additional fruiting forms). Secondly, nitrogen deficiency slightly increases shed of young bolls presumably due to slowing the formation of photosynthate.

Excess nitrogen is currently thought to increase fruit shed by favoring rank growth, which leads to shading (see section on "Light"), reduced photosynthesis and shed.

Diseases

Some causes of vascular wilts, such as *Fusarium* and *Verticillium*, increase fruit shed by preventing the plant from moving water and sugars to the fruit. Additionally, some strains of *Verticillium* induce the production of the abscission formation hormones ethylene and ABA. Early squares have been shed from *Fusarium* infections in the bloom and as many as 30% more squares have shed without fungicide applications during bloom as those that have had fungicide application.

Early Cotton

There are several reasons to set a crop of cotton as quickly as possible and avoid relying on a late or top crop. These reasons include:

- A cotton plant has a greater number of blooms during the initial weeks of flowering than later in the fruiting period (Table 2).
- A cotton plant also sets a higher percent of blooms during the first weeks of flowering (Table 2). When taken together, these two

factors result in a potential of 88% of the crop being made in the first three weeks of flowering.

- Bolls set during the first 3 weeks of fruiting usually are the largest and contain the highest quality fiber. Late-set bolls are frequently smaller and may contain finer and less mature fiber.
- A delay in setting fruit encourages plants to grow taller. This may lead to lodging and makes pest control more difficult.
- Pest populations tend to increase as the season progresses. Protecting squares and young bolls late in the growing season is more difficult (and expensive) than protecting an early crop.

Growing Degree Days

One of the keys to cotton growth and development is temperature. Without a temperature that is sufficient for physiological processes to take place, adequate light, nutrients and water would be of little use to a cotton plant. Efforts by researchers have shown that the cotton plant develops on an orderly schedule that is controlled largely by temperature and that the minimum temperature at which a cotton plant grows is approximately 60°F. From this knowledge came the concept of DD-60s or growing degree-day summations (Table 3). Degree-days for cotton are calculated as follows:

$$((\text{Daily high temperature} + \text{Daily low temperature}) \div 2) - 60^{\circ}\text{F} = \text{Degree-Days}$$

Although the growing degree-day concept is applicable to most situations, some factors such as cultivar or geographic location may cause poor approximations of actual plant growth. Likewise, problems may be encountered if plants are under water or nutrient stress or have been damaged by insects, weather or chemicals. Table 3 gives the generally accepted DD-60s for cotton in the southeast.

Summary

A considerable body of knowledge on the development of the cotton plant has been generated through research. This publication was written to

provide an introduction to the understanding of how a cotton plant develops. For more complex developmental problems and interactions not covered in this publication, consult a specialist with the Cooperative Extension Service.

Table 1. Typical growth and development of a cotton plant.

Event	Time Required (Days)	Average Time Required (Days)
Planting to emergence	4-14	7
Planting to 1st square	35-45	39
Planting to 1st Bloom	55-70	62
Pinhead square to white flower	20-35	23
White flower to pink flower	1	1
Pink flower to open boll	50-60	55

Table 2. Development of fruiting in cotton.

Week of Blooming	% of Total Blooms	% Blooms Set	% of Crop
1	8	94	21
2	24	78	43
3	29	43	4
4	27	21	9
5	10	13	2
6	2	11	1

Table 3. DD-60s required for cotton development.

Event	DD-60s from Planting
Emergence (stand establishment)	45-130
Appearance of first square	440-530
Appearance of first flowers	780-900
Peak Blooming	1350-1500
First open boll	1650-1850
Defoliation	1900-2600

Reference

Landivar, J. A., and J. H. Benedict. 1996.
Monitoring system for the management of cotton growth and fruiting.
<http://agfacts.tamu.edu/~jbenedic/monitor.html>.