

# ENY-854 Quick Reference Guide to Citrus Insecticides and Miticides **UF** UNIVERSITY of FLORIDA

M.E. Rogers, P. A. Stansly, L. L. Stelinski and J. D. Yates



## Products recommended in the Florida Citrus Pest Management Guide and their effects on selected pests and their natural enemies.

| Pesticide active ingredient                 | Target pest                 |                  |                  |                  |                  |                    |                  |                  | Effects on natural enemies |
|---|-----------------------------|------------------|------------------|------------------|------------------|--------------------|------------------|------------------|----------------------------|
|   | Mode of Action <sup>1</sup> | Psyllid          | Leafminer        | Rust Mites       | Spider Mites     | Root Weevil Adults | Scale Insects    | Mealybugs        |                            |
| Abamectin + oil                             | 6                           | ++               | +++ <sub>R</sub> | +++ <sub>R</sub> | +                | +(oil)             | +(oil)           | +(oil)           | medium                     |
| Acetamiprid                                 | 4                           | -                | +++ <sub>R</sub> | -                | -                | ?                  | +                | ++               | medium                     |
| Aldicarb                                    | 1A                          | +++ <sub>R</sub> | -                | +++ <sub>R</sub> | +++              | -                  | -                | -                | low                        |
| Carbaryl                                    | 1A                          | +++ <sub>R</sub> | -                | +                | -                | +++ <sub>R</sub>   | +++ <sub>R</sub> | +                | high                       |
| Chlorpyrifos                                | 1B                          | +++ <sub>R</sub> | +                | +                | -                | +                  | +++ <sub>R</sub> | +++ <sub>R</sub> | high                       |
| Diflubenzuron                               | 15                          | ++               | +++ <sub>R</sub> | +++ <sub>R</sub> | -                | +++ <sub>R</sub>   | -                | -                | low                        |
| Dimethoate                                  | 1B                          | +++              | -                | -                | -                | ?                  | +++ <sub>R</sub> | +                | high                       |
| Fenbutatin oxide                            | 12                          | -                | -                | +++ <sub>R</sub> | +++ <sub>R</sub> | -                  | -                | -                | low                        |
| Fenpropathrin                               | 3                           | +++ <sub>R</sub> | -                | +                | +                | +++ <sub>R</sub>   | -                | +                | high                       |
| Imidacloprid (soil application, nonbearing) | 4                           | +++ <sub>R</sub> | +++ <sub>R</sub> | -                | -                | +                  | ++               | +                | low                        |
| Imidacloprid (foliar application)           | 4                           | +++ <sub>R</sub> | +                | -                | -                | -                  | ++               | +                | medium                     |
| Petroleum oil                               | NR                          | +                | ++ <sub>R</sub>  | ++ <sub>R</sub>  | ++               | +(eggs)            | ++ <sub>R</sub>  | +                | low                        |
| Phosmet                                     | 1B                          | +++ <sub>R</sub> | -                | +                | ?                | +++ <sub>R</sub>   | ?                | ?                | medium/high                |
| Pyridaben                                   | 21                          | -                | ?                | +++ <sub>R</sub> | +++ <sub>R</sub> | -                  | -                | -                | high                       |
| Spinosad                                    | 5                           | -                | +++ <sub>R</sub> | -                | -                | -                  | -                | -                | low                        |
| Spinetoram                                  | 5                           | +++ <sub>R</sub> | +++ <sub>R</sub> | -                | ?                | ?                  | ?                | ?                | low                        |
| Spirodiclofen                               | 23                          | -                | -                | +++ <sub>R</sub> | +++ <sub>R</sub> | ?                  | -                | -                | low                        |
| Spirotetramat                               | 23                          | +++ <sub>R</sub> | -                | +++ <sub>R</sub> | ?                | ?                  | ++               | ?                | low                        |
| Sulfur                                      | NR                          | -                | -                | +++ <sub>R</sub> | +++              | -                  | ?                | ?                | high (short term)          |
| Zeta-cypermethrin                           | 3                           | +++ <sub>R</sub> | -                | -                | ?                | +++                | ?                | ?                | high                       |

<sup>1</sup>Mode of action class for citrus pesticides from the Insecticide Resistance Action Committee; NR = no resistance potential

(R) = product recommended for control of pest in Florida Citrus Pest Management Guide

(+++)= good control of pest

(++)= short-term control of pest

(+)= low levels of pest suppression

(-)= no observed control of pest

(?)= insufficient data available

**Revised December 2008**



### **Imidacloprid soil drench rates for solid plantings of nonbearing citrus**

| Tree Height  | Rate Product/A* | Applications per season | Ounces per tree | Trees per ounce |
|--|-----------------|-------------------------|-----------------|-----------------|
| <b>Imidacloprid 2F (Admire and various generic products)</b> |                 |                         |                 |                 |
| 2 ft – 4 ft  | 8 fl oz         | 4                       | 0.057 fl oz     | 17.5 trees      |
| 4 ft – 6 ft  | 16 fl oz        | 2                       | 0.114 fl oz     | 8.77 trees      |
| <b>Imidacloprid 4.6F (Admire PRO)</b>                        |                 |                         |                 |                 |
| 2 ft – 4 ft  | 3.5 fl oz       | 4                       | 0.025 fl oz     | 40 trees        |
| 4 ft – 6 ft  | 7 fl oz         | 2                       | 0.05 fl oz      | 20 trees        |
| <i>*Rates based on 140 trees per acre</i>                    |                 |                         |                 |                 |

### **Restricted Entry Interval (REI) and Pre-Harvest Interval (PHI) for products listed on front page**

| Active Ingredient   | Product brand names*            | Restricted entry interval (REI) | Pre-harvest interval (PHI) |
|---|---------------------------------|---------------------------------|----------------------------|
| Abamectin   | Agri-mek 0.15EC                 | 12 hours                        | 7 days                     |
| Acetamiprid   | Assail 70 WP                    | 12 hours                        | 7 days                     |
| Aldicarb  | Temik 15G                       | 48 hours                        | 0; 30 days lemons          |
| Carbaryl  | Sevin XLR Plus                  | 12 hours                        | 5 days                     |
| Chlorpyrifos  | Lorsban 4 E                     | 5 days                          | 21 days                    |
| Diflubenzuron   | Micromite 80 WGS                | 12 hours                        | 21 days                    |
| Dimethoate  | Dimethoate 4 E                  | 48 hours                        | 15 days                    |
| Fenbutatin oxide  | Vendex 50 WP                    | 48 hours                        | 7 days                     |
| Fenpropathrin   | Danitol 2.4 EC                  | 24 hours                        | 1 day                      |
| Imidacloprid  | Admire Pro 4.6 F, Provado 1.6 F | 12 hours                        | 0                          |
| Phosmet   | Imidan 70 W                     | 24 hours                        | 7 days                     |
| Pyridaben   | Nexter miticide                 | 12 hours                        | 7 days                     |
| Spinosad  | Spintor 2 SC                    | 4 hours                         | 1 day                      |
| Spinetoram  | Delegate WG                     | 4 hours                         | 1 day                      |
| Spirodiclofen   | Envidor 2 SC                    | 12 hours                        | 7 days                     |
| Spirotetramat   | Movento 240 SC                  | 24 hours                        | 1 day                      |
| Sulfur  | various                         | 12 hours                        | 0                          |
| Zeta-cypermethrin   | Mustang 1.5 EW, Mustang Max EC  | 12 hours                        | 1 day                      |
| <i>*Additional generic brand names may be available for a given active ingredient. Use pesticides safely. Read and follow directions on the manufacturer's label.</i> |                                 |                                 |                            |

Additional citrus pest management information can be found in the Florida Citrus Pest Management Guide available online at <http://www.crec.ifas.ufl.edu/extension/pest/index.htm>