



Identifying Nutrient Deficiencies in Ornamental Plants¹

Amy L. Shober and Geoffrey C. Denny²

Introduction and Purpose

Healthy plant growth and reproduction requires 17 nutrients. Of these, carbon (C), oxygen (O) and hydrogen (H) are derived mainly from the atmosphere and water. Soil minerals and/or soil organic matter are the main source of the remaining essential nutrients. When essential nutrients in plant tissue are present at levels below the needs of a specific plant, that plant will often exhibit deficiency symptoms. If left untreated, nutrient deficiencies may reduce plant growth, flowering, fruiting or aesthetic appeal. Under these circumstances, fertilizers are often applied to landscapes and gardens to provide essential plant nutrients and correct plant deficiency symptoms. The purpose of this publication is to provide guidance for the visual diagnosis of major nutrient deficiency symptoms in ornamental plants.

Essential Plant Nutrients

Essential plant nutrients are divided into two main groups: 1) macronutrients and 2) micronutrients. Macronutrients are needed in greater quantities for growth and are generally found in higher concentrations in plant tissue than the

micronutrients. This also means that plant growth is generally more limited by the availability of macronutrients than by micronutrients. Nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), and sulfur (S) are plant macronutrients. Iron (Fe), zinc (Zn), manganese (Mn), copper (Cu), molybdenum (Mo), boron (B), nickel (Ni) and chlorine (Cl) are plant micronutrients. For more information on the essential plant nutrients and their function in plants see EDIS publication SL 200 *Plant Nutrients for Citrus Trees*: <http://edis.ifas.ufl.edu/ss419>.

Identifying Plant Nutrient Deficiencies

Plants may exhibit nutrient deficiency symptoms if soil levels of one or more essential nutrients are lower than the needs of a specific plant or if soil conditions prevent one or more plant nutrients from being in a form that plant roots can absorb. Even though these nutrient deficiencies often present visual symptoms, it may be difficult to identify specific plant nutrient deficiencies without plant tissue testing. This is because many deficiencies will present symptoms that look very similar to one another. Also,

1. This document is SL 318, one of a series of the Soil and Water Science Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date March 2010. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.

2. Amy L. Shober, assistant professor, Department of Soil and Water Science, Center for Landscape Conservation and Ecology, Gulf Coast Research and Education Center (REC)--Balm; Geoffrey C. Denny, assistant professor, Department of Environmental Horticulture, Center for Landscape Conservation and Ecology, Gulf Coast REC--Balm; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

it is quite easy to mistake a nutrient deficiency symptom for other abiotic problems, such as pesticide drift, or a biotic plant disease. As a rule of thumb, biotic problems tend to appear with asymmetrical patterns, while nutrient deficiency symptoms tend to be distributed symmetrically over the plant tissue surface. Finally, if plants are under additional stress (e.g., water, insect or disease) or if more than one nutrient is deficient, visual symptoms may be misleading.

The first step to visual identification of a nutrient deficiency is to determine where on the plant the symptoms are appearing. This will be dictated by whether the nutrient is mobile or immobile in the plant. A nutrient is considered mobile if the plant is able to move it from one part of the plant to another as it is needed. In contrast, immobile plant nutrients cannot move to different parts of the plant. Therefore, deficiency symptoms of mobile nutrients will appear first in older parts of the plant (Figure 1), while symptoms for immobile nutrients will be seen first in new growth (Figure 2). For example, symptoms of N, P, K and Mg deficiencies will manifest in the old growth because they are mobile, while symptoms of Ca, Cu, B, Fe, Mn and Zn deficiencies will appear in the new growth because they are immobile.



Figure 1. Nutrient deficiency symptoms appearing on old growth. In this case, old leaves are exhibiting chlorosis or yellowing. (Credit: University of Florida – IFAS Plant Nutrient Deficiency Database)

The next step to visual identification of a nutrient deficiency is to determine the characteristics of the symptoms. For example, leaves may appear chlorotic



Figure 2. Nutrient deficiency symptoms appearing on new plant growth. (Credit: University of Florida – IFAS Plant Nutrient Deficiency Database)

or yellow in color (Figure 1) or there may be necrotic or dead tissue (Figure 3). Symptoms may also present on the leaf tips (Figure 3) or be interveinal (Figure 4). The dichotomous key can be used to aid in the visual diagnosis of common nutrient deficiencies in plants when symptoms present first in old leaves (Figure 5) or new leaves (Figure 6).



Figure 3. Necrotic (dead) tissue on the leaf tip. (Credit: Geoffrey Denny, UF-IFAS)

Once a nutrient deficiency symptom has been identified, corrective actions, such as applying fertilizers or lime, can be taken. Before fertilizers are applied to the soil, a soil test should be completed to determine the nutrient status and pH of the soil. For more information on soil sampling and testing, see EDIS publication SL281 *Soil Sampling and Testing for the Home Landscape or Vegetable Garden*: <http://edis.ifas.ufl.edu/ss494>. This is important because when soil pH is too acidic, elements such as K, Ca and Mg may exist in a chemical form that plant roots cannot absorb even if these elements are present in sufficient amounts. Similarly, Fe, Mn, Zn and B



Figure 4. A leaf exhibiting interveinal chlorosis. (Credit: University of Florida-IFAS Plant Nutrient Deficiency Database)

exist in forms that are unavailable to plants at alkaline pH. For more information on soil pH and nutrient availability see EDIS publication SL 256 *Soil pH and the Home Landscape or Garden*: <http://edis.ifas.ufl.edu/ss480>. Under these circumstances, a simple soil test can prevent the application of excess fertilizers and thereby reduce the risk of plant or environmental degradation. Also, it can identify situations where foliar fertilizer applications may be more effective than soil applications. For more information about soil sampling and testing see EDIS publication SL 281 *Soil Sampling and Testing for the Home Landscape or Vegetable Garden*: <http://edis.ifas.ufl.edu/ss494>.




Figure 5. Guide to visual diagnosis of common essential plant nutrient deficiencies for symptoms that appear first in old leaves. (Credit: Geoffrey Denny, UF-IFAS)




Figure 6. Guide to visual diagnosis of common essential plant nutrient deficiencies for symptoms that appear first in new leaves. (Credit: Geoffrey Denny, UF-IFAS)