

Eye Gnats, Grass Flies, Eye Flies, Fruit Flies *Liohippelates* spp. (Insecta: Diptera: Chloropidae)¹

Erika Machtinger and Phillip E. Kaufman²

Introduction

Members of the genus *Liohippelates*, formerly *Hippelates*, are very small true flies and a common occurrence in much of North and South America (Sabrosky 1980, Kumm 1935). These non-biting pests are attracted to fluids secreted by the eyes, nose, and ears on both humans and animals. Some species are attracted to discharge from open wounds and excrement (Goddard 2007). Because of their propensity for hovering around the eyes, this genus has been referred to commonly as “eye gnats,” but are also known as grass flies, eye flies, and fruit flies.

Extremely large aggregations of eye gnats are common in areas that have loose sandy soils, especially in the southern United States. These high concentrations of flies are a great nuisance to humans and animals in rural towns as well as agricultural, recreational, and tourist areas. While they do not bite, *Liohippelates* spp. have been implicated in the transmission of several diseases to humans and livestock including human acute conjunctivitis (pink eye). Due to the increase of transmission of pink eye and the reduction in labor efficiency caused by extreme numbers of eye gnats in the Coachella Valley, California, a control project was initiated by the Bureau of Entomology in the 1920s to focus on the life history and possible control measures for this pest. Hall (1932) subsequently implicated *Liohippelates pusio* (Loew) as one of the limiting factors in the development of the Coachella Valley.

The majority of the flies encountered in the southeastern region of the United States are *L. pusio* and *L. bishoppi* (Sabrosky), and as a result the biological information below will focus broadly on these two species.



Figure 1. Lateral view of an adult *Liohippelates* sp., from a horse farm in north central Florida.

Synonymy

Lioppelates pusio was first named *Hippelates pusio* by Loew in 1872, and *L. bishoppi* was described by Sabrosky as *H. bishoppi* in 1941. In 1929, Duda described the genus *Liohippelates*, which now contains the most important

1. This document is EENY-485 (IN884), one of a series of the Entomology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date April 2011. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. Erika Machtinger, graduate student, and Phillip E. Kaufman, Entomology and Nematology Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

common species of eye gnats in the southeastern United States including what was *H. pusio* and *H. bishoppi*.

Liohippelates pusio (Loew)

Hippelates pusio Loew, 1872

Hippelates splendens Adams, 1904

Hippelates lituratus Becker, 1912

Liohippelates pusio; Duda

Liohippelates bishoppi (Sabrosky)

Hippelates bishoppi Sabrosky, 1941

Liohippelates bishoppi; Duda

Distribution

Both *Liohippelates* spp. and closely related *Hippelates* spp. occur throughout much of North America (Sabrosky 1941) and 270 species have been described (Sabrosky 1987). Bingham (1941) determined during his investigations of *Liohippelates* in the southeastern U.S. that *L. pusio* was the most common pest of this genera. Later that year, Sabrosky (1941) found and described a new species, *L. bishoppi*, misidentified by Bingham as *L. pusio*, which made up a moderate portion of the southeastern *Liohippelates*.

Liohippelates bishoppi is found from February through September, and *L. pusio* year round. *Liohippelates pusio* can be found from Washington to North Dakota, south to Pennsylvania, as well as California south to Mexico and east to the southeastern United States and Bermuda (GNI 2010). *Liohippelates bishoppi* is found from Saskatchewan to Quebec, south to Colorado, Texas and Florida. Other *Liohippelates* species found in the southeastern and eastern U.S. are *L. bicolor* (Coquillett) and *L. pallipes* (Loew), whereas *Hippelates* species found in the same area include *H. nobilis* Loew, *H. plebejus* Loew, and *H. dorsalis* Loew (Sabrosky 1941).

Description

Adults: Adult *Liohippelates* are true flies (bearing only two wings) and are very small, approximately 1.5 to 2 mm long, and shiny black or grayish in color with clear wings. Some species have reddish or yellowish heads. Legs tend to be reddish-yellow to brown with brown bands (Hall 1932).

Eggs: The eggs of *Liohippelates* flies are a pearlescent white and approximately 0.5 mm long. They are half as wide (0.25 mm) as long, resembling a banana with one side curved



Figure 2. Lateral view of an adult *Liohippelates*, unknown species.

and one nearly straight (Hall 1932). Eggs are deposited below the surface of loose soils.

Larvae: The larvae hatch in seven to 11 days and feed on organic matter (Kahn 2008). The larvae average 3 mm in length and are a whitish color (Hermes and Burgess 1930). Larvae are pointed towards the anterior end and rounded towards the posterior. The mouth hook is darker than the rest of the body and curved gently downward.





Figure 3. Primary *Liohippelates* breeding sites in Florida include loose soils with organic matter and moisture, as is found on many livestock facilities throughout the state.

Pupae: The pupae are approximately 2.25 mm in length, depending on the media in which larvae developed and grew (Hall 1932), and a reddish-brown color, turning darker as they age. *Liohippelates* pupate in or around sand mixed with organic matter, such as manure, hay, and remaining harvested crop debris.

Life Cycle

Liohippelates spp. are holometabolous, meaning they have four distinct life stages: egg, larvae, pupa and adult. The life cycle of *Liohippelates* can range from 11 days to three months depending on development conditions such as temperature and moisture (Hall 1932). Development from egg emergence to the adult is completed in approximately three weeks during the summer in Florida (Bigham 1941) and multiple generations can occur each year. Breeding sites are primarily those with freshly disturbed soil mixed with organic matter, such as cut grass and hay, and moisture. Disturbances can be caused by digging, plowing, harrowing, or even by livestock activities (Bigham 1941, Mulla 1962).



Eye gnat. A, Adult. B, Eggs. C, Larva. D, Pupa.

Figure 4. Life cycle of *Liohippelates* species

Medical and Veterinary Importance

Liohippelates are not host specific and will feed on the fluid from wounds, eyes, nose, and other areas on humans as well as livestock and domestic pets in a variety of situations. While they do not bite, due to their persistent feeding

behavior this group of insects is responsible for extreme annoyance to people and animals when they occur in large numbers. In livestock, it has been suggested that high numbers of *Liohippelates* can cause losses in condition, such




Figure 5. Congregation of adult *Liohippelates* spp. around the eye of a horse.

as weight, due to continuous and unrelieved attack (Hinkle et al. 2001).

Many species of *Liohippelates* have been implicated in the mechanical transmission of several diseases and conditions in humans and livestock. Mechanical transmission of pathogens causing disease from *Liohippelates* to humans or livestock occurs when a fly comes in contact with a pathogen, such as a bacterium or virus, and subsequently feeds or rests on a human or animal, transmitting the pathogen from one location to another by contact. As feeding occurs on the blood, mucous, and other fluids around natural orifices of mammals, pathogens can be carried easily into the body.

Human acute conjunctivitis, also known as “pink eye” or “sore eyes,” is caused by one or more bacteria that can be mechanically transmitted by *Liohippelates*. Pink eye causes pain, swelling, changes in vision and extreme sensitivity to light (MIDEHA 2010). Incidents of conjunctivitis have been shown to increase during *Liohippelates* outbreaks (Dow and Hines 1957, Greenberg 1973). In California, Hall (1926)

correlated pink eye to high populations of *Lioppelates flavipes* (Loew).

Liohippелates flavipes and *L. pallipes* have been shown to mechanically transmit the spirochete *Treporema pertenuе*, the causative agent of yaws, in Jamaica and parts of South America (Kumm 1935, Kumm and Turner 1936, Saunders et al. 1936). Yaws is a skin infection that causes ulceration and, in severe cases, can affect bone and cartilage. Yaws occurs mainly in warm and humid regions of the world, including much of South America, and primarily affects



Figure 6. This person has yaws with 8 month-old juxta-articular nodules on the left elbow.

children under the age of 15. Yaws is especially common in poor communities and those in crowded conditions (DNZ 2010).

Another form of acute conjunctivitis caused by the bacterium *Haemophilus influenzae* biotype *aegyptius* has been linked to a few species of *Liohippелates* (Tendella et al. 1994). This bacterium has been linked to the cause of Brazilian purpuric fever, which causes sepsis in children (Harrison et al. 1989, The Brazilian Purpuric Fever Study Group 1992). Sepsis is a serious condition caused by extremely high numbers of foreign microbes in the bloodstream that results in hyperactivity of the body's natural immune response, negatively affecting other body functions. Additionally, *Liohippелates* is capable of mechanically

transmitting streptococcal infections to human skin (Francy et al. 1988). While they do not bite or bloodfeed, *Liohippелates* possess sponging mouthparts with “spines” that have the potential to cause ocular lesions in animals (Hinkle et al. 2001). Two species of *Hippelates* found in Florida, *H. plebejus* and *H. nobilis*, are larger species with rasping mouthparts capable of producing a biting sensation when feeding on sensitive skin (Bigham 1941).

Liohippелates has been implicated in mechanical transmission of pathogens between animals as well. In cattle, *Liohippелates* can cause acute bovine mastitis (Sanders 1940) and vesicular stomatitis (Taplin et al. 1967). Acute bovine mastitis is inflammation of the mammary glands, primarily caused by bacterial infection, which can be extremely painful and compromise milk quality. Mastitis may also cause death in extreme cases (Bradley 2002). Treatment for mastitis includes the use of antibiotics, which has possible implications in public health as resistant bacterial strains may not respond to treatment (Bradley 2002) and milk from treated cattle cannot be sold for human consumption until the required withholding time has elapsed.

Vesicular stomatitis is a reemerging viral disease that presents as lesions on the tongue, coronary bands, and other body regions of cattle as well as horses, sheep, goats, and pigs (USDA 2007). This disease is remarkably similar in presentation to the deadly foot-and-mouth disease, which was eradicated from the United States in 1929 (USDA 2007), and infection can transfer to people who handle infected animals.

Management

Currently, there is no effective area-wide control. Various insecticides, fogs, and soil treatments have been tested, but due to the incredible numbers of flies produced in the soil, area-wide treatment is nearly impossible. Insecticides applied on a community level for mosquito control may reduce attack, but development sites quickly repopulate as the insecticide dissipates (Kahn 2008).

Physical barriers, such as screened porches and windows on houses and fly masks or fly sheets (which resemble a fine cloth mesh, largely impenetrable by even the smallest pests) on livestock, can provide some relief from annoyance and protection from possible pathogen transmission. Sanitation methods to eliminate potential breeding sites, such as reducing manure and organic matter in soil and reducing loose soil matter, may reduce breeding at the location. However, often reducing *Liohippелates* and *Hippelates* breeding sites is nearly impossible in livestock facilities



Figure 7. Congregation of adult *Liohippelates* around the puncture wound of a horse.

as manure, moisture, and soil disturbance are inseparable from animals.

Repellents containing diethyl toluamide (DEET) provide temporary protection from eye gnats (Hall and Gerhardt 2009), and many commercial livestock insecticides that contain pyrethrins, piperonyl butoxide, or other common ingredients may provide some relief to animals.

[Florida Insect Management Guide for external parasites around animal facilities](#)

[Florida Insect Management Guide for filth-breeding flies](#)

Selected References

- Bigham JT. 1941. *Hippelates* (eye gnats) investigations in the southeastern states. *Journal of Economic Entomology* 34: 439-444.
- (BPFSG) The Brazilian Purpuric Fever Study Group. 1992. Brazilian purpuric fever identified in a new region of Brazil. *Journal of Infectious Disease* 163: 516-519.
- Bradley A. 2002. Bovine mastitis: an evolving disease. *The Veterinary Journal* 164: 116-128.
- (DNZ) DermNet NZ. (January 2010). Yaws. *DermNet NZ: the dermatology resource*. <http://dermnetnz.org/bacterial/yaws.html> (28 February 2011).
- Dow RP, Hines JD. 1957. Conjunctivitis in southwest Georgia. *Public Health Reports* 72: 441-448.
- Dow RP, Bigham JT, Sabrosky CW. 1951. Sequel to “*Hippelates* (eye gnat) investigations in the southeastern states” by John T. Bigham. *Proceedings of the Entomological Society of Washington* 53: 263-271.
- Francy DB, Moore LG, Smith GC, Jakob WL, Taylor SA, Calisher CH. 1988. Epizootic vesicular stomatitis in Colorado, 1982: Isolation of virus from insects collected along the northern Colorado Rocky Mountain Front Range. *Journal of Medical Entomology* 25: 343-347.
- (GNI) Global Names Index. (2010). *Global Names Index - Index of Scientific Names*. http://gni.globalnames.org/name_strings (28 February 2011).
- Goddard J. 2007. Non-biting Flies. pp. 191-200. *In* Physicians Guide to Arthropods of Medical Importance, 5th edition. CRC Press, Boca Raton, FL. 480 pp.
- Greenberg B. 1973. Flies and Disease. Vol. 2. Biology and Disease Transmission. Princeton University Press, Princeton, NJ. 856 pp.
- Hall RD, Gerhardt RR. 2009. Flies (Diptera). *In* Mullen Gr, Durden LA (editors), *Medical and Veterinary Entomology*, Second Edition. Elsevier, Burlington, MA. 637 pp.
- Hall Jr. DG. 1932. Some studies on the breeding media, development, and stages of the eye gnat *Hippelates pusio* Loew (Diptera: Chloropidae). *American Journal of Epidemiology* 16: 854-864.
- Harrison IH, Da Silva GA, Pitman M, Fleming DW, Vranjac A, Broome CV. 1989. Epidemiology and clinical spectrum of Brazilian purpuric fever. *Journal of Clinical Microbiology* 27: 599-604.
- Herms WB, Burgess, RW. 1930. A description of the immature stages of *Hippelates pusio* Loew and a brief account of its life history. *Journal of Economic Entomology* 23: 600-603.
- Hinkle NC, Scholl PJ, Mock DE and Warner WB. 2001. Research and extension needs for integrated pest

- management for arthropods of veterinary importance. pp. 261-262. In Geden CJ, Hogsette JA (editors), Proceedings of a workshop in Lincoln, Nebraska. Second Edition. 328 pp.
- (ITIS) (2010). *Liohippelates*. Integrated Taxonomic Information System. <http://www.itis.gov/> (28 February 2011).
 - Kahn C (ed.). (2008). Eye gnats. *The Merck Veterinary Manual*. <http://www.merckvetmanual.com/mvm/index.jsp?cfile=htm/bc/71717.htm&word=hippelates> (28 February 2011).
 - Kumm HW. 1935. The natural infection of *Hippelates pallipes* Loew with the spirochete of yaws. *Royal Society of Tropical Medicine and Hygiene* 29: 265-272.
 - Kumm HW, Turner TB. 1936. The transmission of yaws from man to rabbits by an insect vector, *Hippelates pallipes* Loew. *American Journal of Tropical Medicine* 16: 1-16
 - Looper M, Stokes SR, Waldner DN, and Jordan ER. (2001). Feeding waste milk to dairy calves. *New Mexico State University College of Agricultural, Consumer and Environmental Sciences*. http://aces.nmsu.edu/pubs/_d/d-208.html (28 February 2011).
 - (MIDEHA) Infectious Disease and Environmental Health Administration. (2010). Conjunctivitis ("Pink Eye"). *Maryland Department of Health & Mental Hygiene*. <http://ideha.dhmh.maryland.gov/pdf/conjunctivitis.pdf> (3 May 2011).
 - Mulla MS. 1962. The breeding niches of *Hippelates* gnats. *Annals of the Entomological Society of America* 55: 389-393.
 - Sabrosky CW. 1941. The *Hippelates* flies or eye gnats: preliminary notes. *Canadian Entomologist* 73: 23-27.
 - Sabrosky CW. 1987. Chloropidae. pp. 1049-1067. In McAlpine JF, et al. (editors), *Manual of Nearctic Diptera*. Volume 2. Research Branch Agriculture. Canada Monograph 28: 675-1332.
 - Sanders DA. 1940. *Hippelates* flies as vectors of bovine mastitis (preliminary report). *Journal of the American Veterinary Medical Association* 97: 306-308.
 - Saunders GM, Kumm HW, Rerrie JI. 1936. The relationship of certain environmental factors to the distribution of yaws in Jamaica. *American Journal of Hygiene* 23: 558-579.
 - Taplin D, Zaias N, Rebell G. 1967. Infection by *Hippelates* flies. *Lancet* 2: 472.
 - Tondella MLC, Paganelli CH, Bortoloho IM, Tankano OA, Trino K and Brandileone MCC. 1994. Isolamento de *Harmophilus aegyptius* associado a febre purpúrica Brasileira de cloropídeos (Diptera) dos generos *Hippelates* e *Liohippelates*. *Revista do Instituto de Medicina Tropical de São Paulo* 36: 105-109.
 - (USDA) United States Department of Agriculture. (2007). Vesicular stomatitis. http://www.aphis.usda.gov/publications/animal_health/content/printable_version/fs_vesicular_stomatitis_07.pdf (6 May 2011).