



Healthy Living: Staying Regular¹

Linda B. Bobroff and Luisa Oliver-Cordero²

What causes constipation in older adults?

- · Low fiber intake
- Certain medications
- Limited mobility
- Dehydration
- Chronic laxative abuse
- Ignoring the urge to defecate
- Decreased motility in the colon
- Decreased rectal sensation

Why is staying regular a concern for older adults?

Having a bowel movement fewer than three times a week is defined as constipation. This condition is very common in persons over age 65. Free-living older adults are less likely to be constipated than are persons living in nursing homes. Over time, chronic constipation can cause serious and painful conditions such as hemorrhoids and diverticular disease.

Diverticula are small outpouchings in the gut. When they get inflamed they can be quite painful.



Figure 1. Prunes Credits: Photo by *las*, and used here under Creative Commons license CC BY-NC-ND 2.0. http://flic.kr/p/8K4C9z

Which medications increase risk for constipation?

Several types of medications can cause constipation in some people. Ask your pharmacist for information about the side effects of each of the medications that you take. Here are some medications commonly associated with constipation:

• Antacids with aluminum or calcium

- 1. La versión en español de este documento es *Vida Saludable: Manteniéndose Regular* (FCS8570-Span). Originally developed with funding from the Florida Department of Elder Affairs in partnership with state, county, and local agencies, this document is FCS8570, one of a series of the Department of Faily, Youth and Community Sciences, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. First published: November 1999. Revised for December 2010 by Linda B. Bobroff. Revised with minor revisions August 2011. Visit the EDIS website at http://edis.ifas.ufl.edu.
- 2. Linda B. Bobroff, PhD, RD, LD/N, professor; Luisa Oliver-Cordero, BS, RD, LD/N, former nutrition educator/trainer; Department of Family, Youth and Community Sciences; Institute of Food and Agricultural Sciences; University of Florida; Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A&M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer-Chancy, Interim Dean

- Anticholinergics
- Antidepressants
- Antihistamines
- Calcium channel blockers
- Iron supplements (high doses)
- Diuretics
- Non-steroidal anti-inflammatories (NSAIDs)

What can I do to prevent constipation?

- Eat about 25–30 grams of fiber a day.
- Be as active as possible every day.
- Drink 6–8 cups or more of water and non-alcoholic drinks each day.
- Get into a routine for having a bowel movement; perhaps after breakfast.
- Take a fiber supplement if you are not able to eat enough dietary fiber each day. Talk to your doctor before using fiber supplements.

What are some examples of high-fiber foods?

Food	Fiber (grams)
Baked beans, 1/3 cup*	6–7
Bran cereal, 1 ounce	5–13
Wheat, shredded, 2 biscuits	5
Bran muffin, 1 medium	2–4
Fresh fruits, 1 medium	2–4
Broccoli, ½ cup (All vegetables contain fiber.)	3
Nuts, ¼ cup	2–3
Whole wheat bread, 1 slice	2
Brown rice, ½ cup	2
*Use canned beans for convenience.	

How can I add fiber to my diet?

Many tasty foods contain fiber! You can eat more fruits, vegetables, nuts, and seeds as snacks and in meals. Make substitutions like using whole grains instead of refined grains in baking.

Here are some more ideas:

- Choose high-fiber breakfast cereals.
- Add fresh, frozen, or canned fruits to low-fat milk or juice, and mix in the blender.
- Add extra vegetables to your favorite casserole or soup.
- Eat brown rice instead of white rice.
- Eat 8–10 almonds, pecans, cashews, or walnuts for a snack
- Select high-fiber breads, cereals, and pastas.
- Purée vegetables in the blender and add to soups, quickbread batter, or other dishes.
- Use beans and nuts more often as protein sources instead of meat.
- Enjoy a piece of fruit for a snack at least once every day.
- Sprinkle chopped nuts on your hot or cold cereal.

Where can I get additional, reliable information?

The Family and Consumer Sciences (FCS) agent at your county Extension office may have more written information and classes for you to attend. In Florida, you can find your Extension office at this website: http://solutionsforyourlife.ufl.edu/map.

A registered dietitian (RD) also can provide reliable information to you.

Visit the following Internet sites for more information:

- htttp://www.seniors.gov
- http://nutrition.gov
- http://solutionsforyourlife.ufl.edu
- http://www.choosemyplate.gov
- Constipation Myths, a University of Florida IFAS
 Extension publication http://edis.ifas.ufl.edu/pdffiles/fs/fs15500.pdf