

Carbon Sequestration and Storage by Gainesville's Urban Forest¹

Francisco Escobedo, Jennifer A. Seitz, and Wayne Zipperer²

Climate change is a world-wide issue, and it may seem as if only actions by national governments can work effectively against it. In fact, individuals and small communities, too, can make wise choices and impacts. Communities can mitigate climate change through reducing fossil fuel consumption and good management of its urban forest. Urban trees can reduce concentrations of atmospheric carbon dioxide by storing carbon in their roots, stems, and branches. Urban forests can also help reduce carbon dioxide emissions from fossil-fuel-based power plants because their shade and wind protection reduces energy consumption for heating and cooling buildings. By estimating the amount of carbon removed by trees, we can determine the role of urban forests in mitigating climate change and also assign an economic value to the amount of carbon sequestered by an urban forest.

Using the USDA Forest Service's Urban Forest Effects (UFORE) model (<http://www.ufore.org>), researchers at UF/IFAS's School of Forest Resources and Conservation and the USDA Forest Service estimated carbon storage and sequestration by urban trees in Gainesville, Florida (Escobedo and others 2010). Carbon storage is the estimated total amount of woody biomass held in a tree's stem and branches over its life. Carbon sequestration is the estimated amount of carbon a tree's stem and branches take up during one year of growth. In 2006, the researchers collected data from 93 one-tenth-acre field plots located randomly across the city of Gainesville. The model uses tree species

equations to estimate tree dry weight from stem diameter. Approximately 50% of a tree's dry weight biomass is carbon; the researchers estimated the dry weight and then divided by two in order to arrive at an estimate of the amount of carbon stored in Gainesville's trees. In estimating carbon sequestration, researchers accounted for average annual growth for different types of trees in different size classes and in different conditions (Nowak and others 2002). Since carbon value is traded in carbon offset markets in units of carbon dioxide, carbon estimates were then converted to carbon dioxide (CO₂) equivalents. Values were multiplied by \$4 per metric ton (mt) CO₂ equivalent or the August 2008 market value (<http://www.ecosystemmarketplace.com/>).

Table 1. Comparison of average carbon stored and sequestered per tree by diameter at breast height (DBH) size classes in Gainesville, Florida.

DBH Class ¹ (cm)	Per Tree C Storage (kg)*	Per Tree C Sequestered (kg/year)
1 – 15	22	2
16 – 30	192	9
31 – 45	658	17
46 – 60	1349	9
61 – 76	2812	33
77+	8858	111

From: Escobedo and others. 2010. *1 kg = 2.2 lbs

1. This document is FOR210, one of a series of the School of Forest Resources and Conservation Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date March 2009. Revised February 2012. Visit the EDIS [website at http://edis.ifas.ufl.edu](http://edis.ifas.ufl.edu).
2. Francisco Escobedo, assistant professor, Jennifer A. Seitz, Extension associate, School of Forest Resources and Conservation, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611. Wayne Zipperer, research scientist, USDA Forest Service.

Table 1 compares the average amount of carbon stored and sequestered per tree by different size classes. Large trees greater than 77 centimeters in diameter at breast height (dbh) sequester the most carbon in Gainesville. Overall healthier and larger trees sequester more carbon annually than do younger trees with a small dbh because of the limited growth and size. Eventually, if small trees remain healthy and continue to grow, they will accumulate more carbon as their biomass increases. Trees in poor condition sequester less carbon than do healthy trees, and dead trees actually emit carbon as they decompose.




Figure 1. Net carbon sequestered per land use area and the associated value in Gainesville's urban forest. Credits: Escobedo and others. 2010.

Since carbon is valued in terms of CO₂ equivalents, C results were converted to CO₂ by multiplying by 3.67. Figure 1 depicts a comparison of the economic value and CO₂ stored by trees located in different land-use areas in Gainesville. Trees in residential and forested areas store 76% of all CO₂, which is more than trees in commercial or industrial areas simply because more of Gainesville is devoted to residential use and to forests than is devoted to commercial and industrial uses. However, by analyzing on a per-hectare basis, 1 hectare of forested land use can sequester 8 mt of CO₂ per year, while residential and industrial areas in 1 year sequester 4 and 1 mt CO₂ per hectare, respectively.

As trees grow, they store more CO₂ by assimilating it in their woody tissue. Conversely, as they die and decompose, they release much of the stored CO₂ back into the atmosphere. Thus, CO₂ sequestration can be negative if there are more dead than live trees (if more CO₂ is being emitted by decomposing trees than is being sequestered by healthy, growing trees). Dead and decomposing trees in Gainesville emitted approximately 3,500 mt CO₂ in the study year. Accounting for this loss, CO₂ sequestration by trees is about 9,740 mt CO₂ per year with a value of \$38,954. Overall, trees in Gainesville capture, or offset, about 3% of all the annual human-related CO₂ emissions in the city (Escobedo and others 2010).




Figure 2. The ten tree species that sequester the most carbon dioxide (CO₂) in Gainesville, Florida (% of all CO₂ sequestered by Gainesville's urban tree population). Credits: Escobedo and others 2010.

The 10 tree species in the city that sequester the most CO₂ are illustrated in Figure 2. Of all the species sampled, laurel oak (*Quercus laurifolia*) sequesters the most carbon dioxide or about 16% of the total CO₂ sequestered. This is because laurel oaks are more abundant in Gainesville than other trees, including live oaks and tupelos. Live oaks are much less abundant than laurel oaks but because of their large size and greater life span they can sequester a greater portion of CO₂ than other trees. Most of the trees in Figure 2 also are on the list of the ten most common trees found in Gainesville. For more information about common trees in Gainesville, read "Gainesville's Urban Forest Structure, Composition, Size, and Density" <http://edis.ifas.ufl.edu/FR276>.

Conclusion

Cities are a major source of carbon dioxide emissions. As demonstrated by this fact sheet, urban and natural trees can help mitigate the effects of climate change somewhat by sequestering CO₂ but can provide multiple other environmental, social, and economic benefits (Dobbs and others 2010). However, urban trees can only sequester a small portion of all carbon dioxide emitted from cities. In addition, decomposing trees and mulch, tree maintenance activities that use fossil-fuel-burning tools (e.g., chainsaws, leaf blowers, mowers, and cars and trucks), and improperly placed trees that cause shading in winter can also result in emissions of CO₂ (Dobbs and others 2011). It is important for communities to reduce fossil fuel emissions and manage for and preserve large, healthy trees to maximize the amount of CO₂ sequestered by an urban forest.

Literature Cited

Dobbs, C., F. Escobedo, and W. Zipperer. 2011. A framework for developing urban forest ecosystem services and goods indicators. *Landscape and Urban Planning*, 99:196-206.

Escobedo, F, S. Varela, M. Zhao, J. Wagner, and W. Zipperer. 2010. Analyzing the efficacy of subtropical urban forest in offsetting carbon emissions from cities. *Environmental Science and Policy*. 13:362-372.

Nowak, D.J. and D.E. Crane. 2002. Carbon storage and sequestration by urban trees in the USA. *Environmental Pollution* 116:381-389.