

# Bacterial Spot of Pepper<sup>1</sup>

Camille McAvoy, Pamela Roberts, and Jeffrey Jones<sup>2</sup>

## Introduction

Pepper fruits in the genus *Capsicum* are produced world-wide as a fresh vegetable, dried for spices, or used in commercial products such as dietary supplements or natural coloring in cosmetic products. Florida is the second largest producer of sweet bell pepper (*Capsicum annuum*), providing 32% of what is grown in the United States. In 2019, the acreage of bell pepper production in Florida was 12,000 acres and valued at nearly \$235 million USD ([nass.usda.gov](https://nass.usda.gov)). In addition to sweet bell, a wide variety of specialty pepper types are also produced in Florida: Thai pepper (*C. frutescens*), habanero and Scotch bonnet (*C. chinense*), ají or Peruvian hot pepper (*C. baccatum*), jalapeño, serrano, and cherry (*C. annuum*). A limiting disease problem on all peppers is bacterial spot caused by three species of *Xanthomonas* (*X. euvesicatoria*, *X. perforans*, and *X. gardneri*) (Jones et al. 2004). *Xanthomonas euvesicatoria* is the predominant species occurring on pepper in the United States, while the other two, *X. perforans* and *X. gardneri*, are considered to be of minor importance on pepper. Bacterial spot causes severe defoliation, resulting in reduced plant vigor, exposure of fruit to sunburn, and reduced yield. Infected fruit are unmarketable due to lesions on fruit surface. Total yield losses in severely affected crops can occur.

## Symptoms

On the plant, all aboveground plant tissue is susceptible to infection by the bacterial pathogen. On leaves, the first symptoms are water-soaked spots visible on the lower leaf surface within 3 to 4 days of infection. Within another 2–3 days, the lesions become tan to brown and appear slightly greasy on the upper leaf surface (Figure 1). Leaf lesions expand and coalesce to form large areas of yellow and brown tissue with a characteristic greasy appearance (Figure 2). Leaf margins may become scorched due to bacterial invasion of hydathodes. Infected leaves often drop prematurely, even when only moderate damage from bacterial spot is incurred (Figure 3). Premature defoliation leading to thinning of the plant canopy increases incidence of sun scald on fruits. Lesions on fruit appear initially as small, raised blemishes that are a slightly lighter green than the normal fruit color. Fruit lesions enlarge, turn brown to black, and appear as raised warts or scabs (Figure 4). Under humid conditions, other microbes, such as those causing bacterial soft rot, may enter the fruit lesions and cause massive fruit decay.

1. This document is PP362, one of a series of the Plant Pathology Department, UF/IFAS Extension. Original publication date February 2021. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Camille McAvoy, assistant research scientist, PhD; Pamela Roberts, professor, Plant Pathology Department, UF/IFAS Southwest Florida Research and Education Center; and Jeffrey Jones, professor, Plant Pathology Department; UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition. Use all chemicals safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 1. Initial symptoms of bacterial spot on pepper leaves exhibiting water-soaking (the dark-green halo surrounding the tan lesion).

Credits: P. Roberts, UF/IFAS



Figure 2. As lesions age, the lesions expand and coalesce to form large areas of yellow and brown tissue.

Credits: P. Roberts, UF/IFAS



Figure 3. Severely defoliated plants with extensive loss of lower leaves due to bacterial spot.

Credits: P. Roberts, UF/IFAS



Figure 4. Bacterial spot lesions on fruit appear as raised warts or scabs and are dark colored.

Credits: P. Roberts, UF/IFAS

## Causal Organism and Host Resistance

Bacterial spot in pepper in Florida and worldwide is primarily caused by *Xanthomonas euvesicatoria*. *Xanthomonas perforans* is most associated with bacterial spot on tomato, but while generally considered nonpathogenic on pepper, *X. perforans* was found causing disease on pepper in Florida and Alabama (Schwartz et al. 2015; Newberry et al. 2019). The *X. euvesicatoria* strains affecting pepper can be divided into eleven races (0 through 10) based on the *Bs1*, *Bs2*, *Bs3*, and *Bs4c* resistance genes in pepper. A summary of the pepper genes can be found in Table 1. There are six dominant resistance genes (*Bs1*, *Bs2*, *Bs3*, *Bs4C*, *Bs7*, and *BsT*) and two recessive resistance genes (*bs5* and *bs6*). Resistance genes in pepper have been incorporated into new cultivars using traditional breeding methods. Currently available commercial bacterial spot-resistant cultivars contain various combinations of these resistance genes. While some bell pepper varieties have resistance to all known races of *X. euvesicatoria*, others do not. Many of the specialty and chile pepper varieties do not have gene resistance packages. For more information on pepper varieties and resistance for Florida production, please see the EDIS chapter on *Pepper Production* in the *Vegetable Production Handbook of Florida* at <https://edis.ifas.ufl.edu/cv130>.

Table 1. Dominant and recessive genes introgressed into pepper that confer resistance to one or more races of *Xanthomonas euvesicatoria*.

Gene nomenclature	Races of <i>X. euvesicatoria</i>	Response
<i>Bs1</i>	0, 2, and 5	Hypersensitive
<i>Bs2</i>	0 through 3, 7, and 8	Hypersensitive
<i>Bs3</i>	0, 1, 4, 7, and 9	Hypersensitive
<i>Bs4C</i>	0, 1, 3, 4, and 6	Hypersensitive
<i>Bs7</i>		Hypersensitive
<i>BsT</i>		Hypersensitive
<i>bs5</i>	0 through 10	Non-hypersensitive
<i>bs6</i>	0 through 10	Non-hypersensitive

## Disease Cycle and Epidemiology

The bacteria are seedborne, and outbreaks of the disease can occur within transplant production. When associated with the seed, plant cotyledons may become infected upon contact with seed coat or upon emerging from the soil (Ritchie 2000). *Xanthomonas euvesicatoria* can survive on volunteer (tomato/pepper) plants in the field as well as in soil when crop residue is present but not yet decomposed (Potnis et al. 2015). The bacteria can also survive epiphytically on nonhost species within the cropping system. *X. euvesicatoria* can survive on other solanaceous crops, including tomato. In tomato, *X. euvesicatoria* is just one of four species that cause bacterial spot. More information on bacterial spot of tomato can be found at <https://edis.ifas.ufl.edu/pp353>. The bacteria are spread from plant to plant by either water droplets, such as from rainfall or heavy dew, or aerosols containing bacteria. Once these droplets or aerosols encounter host tissues, *X. euvesicatoria* enters through wounds and various natural openings, including stomata and lenticels. It is not uncommon to observe severely diseased varieties next to relatively unaffected plants in the field, even in the absence of resistance genes in both. Outbreaks are possible whenever warm weather occurs with frequent episodes of wind-driven rain as well as night temperatures above 65°F. These environmental conditions favor the likelihood of bacterial spot developing in pepper plantings. In southern Florida, the most severe outbreaks are observed in the months from August through November, coinciding with the rainy season.

## Disease Management

Bacterial spot management requires an integrated approach using several tactics including cultural, sanitation, and chemical control measures.

## Cultural and Sanitation Practices

Host resistance is a critical component of an integrated pest management (IPM) strategy to control bacterial spot on pepper. If known bacterial spot infections have occurred in the past, start by choosing a variety that contains resistance genes. Some newer varieties offer resistance genes to all known races, while many of the older varieties do not contain all available resistance genes. Careful consideration should be given to the choice of variety. Consult the *Vegetable Production Handbook of Florida* (<https://edis.ifas.ufl.edu/cv130>) for an updated list of pepper varieties. Because bacteria transmission is possible in pepper seed, choose certified disease-free seed when possible. Scouting is a critical component of IPM, and plants in fields and transplant houses should be inspected frequently. Inspect all transplants for disease, and use only healthy transplants. In fields where peppers or tomatoes have been produced previously, all volunteers and host crop debris should be destroyed before beginning field preparation for a new season. It is recommended to rotate fields to avoid volunteer and host crop debris. Water is critical in the spread and development of bacterial spot. Although becoming less common, overhead irrigation should be avoided whenever possible. Workers and farm equipment should be kept out of fields when conditions are wet as the bacteria can be readily spread from plant to plant through contact with wet foliage. Disinfect hand tools and other equipment as a precaution to prevent bacterial spread. Other *Xanthomonas* species (such as *Xanthomonas axonopodis* and *Xanthomonas musacearum*) are capable of dispersing via dirty tools and hands (An et al. 2020). Cull piles should be located as far away from field operations as possible.

## Chemical

The effectiveness of copper bactericides is limited because of the widespread occurrence of copper tolerance among strains of *X. euvesicatoria* in Florida (Marco and Stall 1983; Stall et al. 1986). However, because copper tolerance is qualitative and not quantitative, copper is still used in most operations, including organic production. Avoid unnecessary use of magnesium as foliar or soil applications because research has shown it may increase severity of bacterial spot on pepper (Jones et al. 1983). When using any copper bactericides, it is recommended that the bactericide be combined with a dithiocarbamate fungicide, such as mancozeb. While the exact mode of action is still unknown for this fungicide, the addition of a dithiocarbamate increases copper solubility. This may account for the increased bactericidal activity. Numerous biopesticides are also

labeled for bacterial spot control on pepper, and a current list may be found in *Chapter 19. Biopesticides and Alternative Disease and Pest Management Products in the Vegetable Production Handbook* (<https://edis.ifas.ufl.edu/cv295>). The efficacy and consistency of many of these products typically is limited. Best results are observed when biopesticides are used to augment the standard copper-mancozeb applications. Biopesticides do not represent a viable replacement for the use of copper-mancozeb applications, especially for the control of copper-tolerant isolates of *X. euvesicatoria*. It is advised to include many products with varying modes of action into a spray program because it can reduce the possibility of a population becoming resistant to any one mode of action.

## Use of Actigard in Chile Peppers

Acibenzolar-S-Methyl (ASM), or Actigard® (US Trade Name), is a systemic acquired resistance inducer, or SAR. It has been used successfully in conjunction with other chemistries in tomato production to manage bacterial spot. However, in pepper production, it is only currently registered for use on hot or non-bell-type peppers. Caution is urged when deciding to incorporate ASM into a bacterial leaf spot management plan for non-bell-type peppers. The label instructions should be followed carefully to ensure the best performance of the product. Used alone, ASM is an inefficient method of management of bacterial leaf spot in peppers (Roberts et al. 2008); however, its use with additional methods, such as biologicals and even host resistance, may show an additive effect in controlling bacterial spot on chile peppers. If using to control *Xanthomonas* in hot or non-bell-type peppers, it is strongly suggested that another chemistry be tank mixed to improve effectiveness.

## Literature Cited

- An, S.-Q., N. Potnis, M. Down, F.-J. Vorhölter, Y.-Q. He, A. Becker, D. Teper, Y. Li, N. Wang, L. Bleris, and J.-L. Tang. 2020. “Mechanistic Insights into Host Adaptation, Virulence, and Epidemiology of the Phytopathogen *Xanthomonas*.” *FEMS Microbiol. Rev.* 44 (1): 1–32. <https://doi.org/10.1093/femsre/fuz024>
- Jones, J. B., S. S. Woltz, and J. P. Jones. 1983. “Effect of Foliar and Soil Magnesium Application on Bacterial Leaf Spot of Peppers.” *Plant Dis.* 67:623–624. <https://doi.org/10.1094/PD-67-623>
- Kurowski, C., K. Conn, and P. Himmel. 2019. “Guideline for Identification of Pepper Bacterial Leaf Spot Races Using Differential Hosts.” CPPSI Collaboration for Plant Pathogen Strain Identification. <https://cppsi.ucdavis.edu/sites/g/files/dgvnsk8206/files/inline-files/225903-2.pdf>
- Marco, G. M., and R. E. Stall. 1983. “Control of Bacterial Spot of Pepper Initiated by Strains of *Xanthomonas campestris* pv. *vesicatoria* That Differ in Sensitivity to Copper.” *Plant Dis.* 67 (7): 779–781. <https://doi.org/10.1094/PD-67-779>
- Newberry, E. A., R. Bhandari, G. V. Minsavage, S. Timilsina, M. O. Jibrin, J. Kemble, E. J. Sikora, J. B. Jones and N. Pontis. 2019. “Independent Evolution with the Gene Flux Originating from Multiple *Xanthomonas* Species Explains Genomic Heterogeneity in *Xanthomonas perforans*.” *Appl. Environ. Microbiol.* 85 (20): e000885-19. <https://doi.org/10.1128/AEM.00885-19>
- Potnis, N., S. Timilsina, A. Strayer, D. Shantharaj, J. D. Barak, M. L. Paret, G. E. Vallad, and J. B. Jones. 2015. “Bacterial Spot of Tomato and Pepper: Diverse *Xanthomonas* Species with a Wide Variety of Virulence Factors Posing a Worldwide Challenge.” *Mol Plant Pathol.* 16 (9): 907–920. <https://doi.org/10.1111/mpp.12244>
- Ryan, R. P., F. J. Vorhölter, N. Potnis, J. B. Jones, M. A. Van Sluys, A. J. Bogdanove, and J. M. Dow. 2011. “Pathogenomics of *Xanthomonas*: Understanding Bacterium-Plant Interactions.” *Nature Reviews Microbiology* 9 (5): 344–355. <https://doi.org/10.1038/nrmicro2558>
- Schwartz, A. R., N. Potnis, S. Timilsina, M. Wilson, J. Patané, J. Martin Jr., G. V. Minsavage, D. Dahlbeck, A. Akhunova, N. Almeida, G. E. Vallad, J. D. Barak, F. F. White, S. A. Miller, D. Ritchie, E. Goss, R. S. Bart, J. C. Setubal, J. B. Jones, and B. J. Staskawicz. 2015. “Phylogenomics of *Xanthomonas* Field Strains Infecting Pepper and Tomato Reveals Diversity in Effector Repertoires and Identifies Determinants of Host Specificity.” *Front. Microbiol.* 6:535. <https://doi.org/10.3389/fmicb.2015.00535>
- Stall, R. E., D. C. Loschke, and J. B. Jones. 1986. “Linkage of Copper Resistance and Avirulence Loci on a Self-Transmissible Plasmid in *Xanthomonas campestris* pv. *vesicatoria*.” *Phytopath.* 76 (2): 240–243. <https://doi.org/10.1094/Phyto-76-240>