

Hydrilla leafcutter moth (unofficial common name); *Parapoynx diminutalis* Snellen (Insecta: Lepidoptera: Crambidae)¹

Julie Baniszewski, Emma N.I. Weeks, and James P. Cuda²

Introduction

Parapoynx diminutalis Snellen is an adventive Asian moth with an aquatic larval stage. The moth is found associated with a variety of water bodies including river backwaters, lakes, and ponds (Habeck 1996). The aquatic larvae commonly attack hydrilla, *Hydrilla verticillata* (L.f. Royle), and other aquatic plants (Buckingham and Bennett 1989, 1996).

The moth was identified in 1971 in India and Pakistan during scouting trips to attempt to determine potential biological control agents for hydrilla. Despite having potential for hydrilla destruction, the moth was declared to be a generalist feeder and unsuitable for release into U.S. water bodies for hydrilla control (Baloch et al. 1980). However, the moth was later found in Florida in 1976 by United States Department of Agriculture technicians who were testing herbicides for hydrilla control. The larvae (caterpillars) found on hydrilla were observed to be eating the invasive weed. The pathway, method, or time of the moth's arrival remains unknown (Del Fosse et al. 1976).

Synonymy

According to the global Pyraloidea database (Nuss et al. 2003–2013) and Shibuya (1928) the following junior synonyms have been used for *Parapoynx diminutalis*:

Parapoynx dicentra Meyrick, 1885

Oligostigma pallida Butler, 1886

Nymphula diminutalis Meyrick, 1894

Nymphula uxoralis Strand, 1919

Distribution

Parapoynx diminutalis is native to parts of Asia, Africa, and Australia (Buckingham and Bennett 1996). In its adventive range, *Parapoynx diminutalis* has been found in Panama (notably in the Panama Canal, which was infested with hydrilla), Honduras, and Florida. In commercial greenhouses, the moth has been observed colonizing aquatic plants in England and Denmark (Agassiz 1978, 1981). *Parapoynx diminutalis* was first seen in Florida in Fort Lauderdale in 1976 but progressively appeared in more northern counties, eventually reaching Alachua and Putnam counties by 1979 (Balciunas and Habeck 1981). In the early 1980s, hydrilla surveys in other southeastern states revealed that the moth's range did not extend beyond Florida (Balciunas and Minno 1985). Even in northern Florida, the cooler water temperatures caused populations to be reduced in late winter and early spring. Milder climates such as those found in Panama may enable populations to thrive throughout the year (Buckingham and Bennett 1996).

1. This document is EENY583, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date January 2014. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

2. Julie Baniszewski, graduate student; Emma N.I. Weeks, assistant research scientist; and James P. Cuda associate professor, Entomology and Nematology Department, UF/IFAS Extension, Gainesville, FL 32611.

More recent studies indicate that the moth also is established in Louisiana, where 37 moths were collected from 1984–1992 (Brou Jr. 1993).

Description

Eggs: Eggs are smooth and bright yellow when laid (Figure 1); they turn white, and then become transparent as they develop. The eggs are generally deposited on plant leaves or stems just below the water surface in masses of various sizes (Buckingham and Bennett 1996).



Figure 1. Eggs of *Parapoynx diminutalis* Snellen, within one day of being laid (left). Egg mass sizes vary and are often laid on plant tissue (right).

Credits: Lyle J. Buss, University of Florida.

Larvae: Larvae can be distinguished from those of other aquatic species by the presence of branched gills (Habeck 1996) and brown spots on the head and the tip of the thorax. The larval stage consists of seven instars; all seven are off-white with yellow-brown legs (Habeck and Balciunas 2005). The larvae are mobile and feed on hydrilla leaves. The first instar is white yet nearly transparent and has 1 mm long setae (hairs) (Figure 2). Instars 2 through 7 are white, later instars begin to turn yellow as they approach pupation (Figure 3). In later instars, the length increases and the




Figure 2. *Parapoynx diminutalis* Snellen, first instar larva eating hydrilla. First instar larvae are transparent, allowing consumed hydrilla to be visible in the gut.

Credits: Lyle J. Buss, University of Florida.

external gills develop (Figure 4). Instars 3 through 7 use plant tissue to construct a silk case, and often retreat into the case between feeding events (Buckingham and Bennett 1996).



Figure 3. An early instar of *Parapoynx diminutalis* Snellen (left). Larvae are mobile and retreat into a cocoon between feedings. Cocoons are constructed of plant materials and attached to a hydrilla stem (right). Credits: Julie Baniszewski, University of Florida.



Figure 4. A late instar of *Parapoynx diminutalis* Snellen, feeding on hydrilla (left). Instars 2 through 7 are white, later instars begin to turn yellow closer to pupation (right). Branched gills are visible and help identify this species in the larval stage.

Credits: Lyle J. Buss, University of Florida.

Pupae: Pupae have three tubercles (or nodules) along each side and two setae (or hairs) on the head. Female pupae can be distinguished from males by their larger size and by their antennae. Female antennae are shorter, extending only to the wing tips, whereas male antennae are longer and extend past the wing tips (Buckingham and Bennett 1996). Late seventh instar larvae (or pre-pupae) enclose themselves in a white cocoon, which is attached to a submerged plant stem (Figure 5). After 1–2 days in the cocoon, the white pre-pupae have developed into yellow pupae inside the cocoon. The eyes turn red, then brown, and the wings become visible as pupation progresses (Buckingham and Bennett 1996).




Figure 5. A cocoon constructed and occupied by a *Parapoynx diminutalis* Snellen larva. *Parapoynx diminutalis* Snellen use plant stems, leaves and other materials to construct their cocoons and attach to submerged stems or plant material. Credits: Julie Baniszewski, University of Florida.

Adults: Moth adults are white with brown or tan markings or bands on the wings and tan bands on the body (Figure 6). Females typically differ from males by their longer wingspans, more pointed forewings, larger abdomens, and shorter antennae, and they lack the noticeable white setae displayed by males on the tip of the abdomen (Buckingham and Bennett 1996).




Figure 6. An adult *Parapoynx diminutalis* Snellen, female moth (left) and male moth (right). Females have longer wingspans, more pointed forewings, and larger abdomens. Males have longer antennae and more distinct white setae (hairs) at the tip of the abdomen. Credits: Lyle J. Buss, University of Florida.

Life Cycle and Biology

Parapoynx diminutalis undergoes complete metamorphosis from an aquatic caterpillar to a moth. Life stages include the egg, seven larval instars (the seventh instar includes a pre-pupa stage), the pupa, and finally adult emergence (Buckingham and Bennett 1996). The life cycle of *Parapoynx diminutalis* ranges from 25 to 41 days for development and about five days for the adult life span. Females lay on average about 200 eggs, but can lay just a few to over 500.

The eggs require 4–6 days to develop before first instars hatch. Adults typically emerge from pupae after dusk and are quick to fly to avoid potential predators. The adults drink water using a reduced proboscis, but they do not appear to feed (Buckingham and Bennett 1996). *Parapoynx diminutalis* mating has not been studied in detail but has been observed occasionally and seems to occur at around three hours after dusk. Although the maximum time *in copula* is unknown, pairs of moths *in copula* facing in opposite directions were noted to rest for at least 30 minutes. After mating, there is a one-day pre-oviposition period. Females then oviposit soon after dusk just below the water surface on leaves or stems. First instars have been shown to hatch both below and above the water surface, although it has been observed that the females typically oviposit below the water surface (Buckingham and Bennett 1996).

Hosts

Larvae are commonly found on the aquatic weed hydrilla, *Hydrilla verticillata*. The initial discovery of the moth on hydrilla led to an interest in the moth as a possible biological control agent of this invasive weed. In the field, larvae and pupae have been found in small numbers on coontail (*Ceratophyllum demersum* L.), southern naiad (*Najas guadalupensis* (Sprengel) Magnus), and Illinois pondweed (*Potamogeton illinoensis* Morong) (Buckingham and Bennett 1996). Furthermore, in laboratory studies, while *Parapoynx diminutalis* larvae preferred hydrilla, they could also complete development on various other plants including coontail, southern naiad, fanwort (*Cabomba caroliniana* Gray), Brazilian waterweed (*Egeria densa* Planchon), and Eurasian water-milfoil (*Myriophyllum spicatum* L.) (Buckingham and Bennett 1989).

Damage

Plant damage is inflicted by the larvae, which not only eat the leaf and stem tissue, but use these materials to prepare their pupal cocoon as well (Figure 7). The main food source for *Parapoynx diminutalis* is the aquatic weed hydrilla (Buckingham and Bennett 1996). In most natural situations in the U.S., hydrilla is invasive and an undesirable weed because it develops surface mats and disrupts natural ecosystems (Haller and Sutton 1975; Hetrick and Langeland 2012). There have been few studies to quantify the effect of feeding moth larvae on hydrilla biomass. Feeding of the moth larvae on hydrilla in Florida was thought to have a positive effect on hydrilla-invaded water bodies (Del Fosse et al. 1976) by reducing the need for herbicide applications to control hydrilla mats. However, naturalized populations of this moth are too sporadic to have a significant effect

on hydrilla density. For example, in northern Florida populations build up during the summer months and can cause extensive defoliation of hydrilla, but in the winter, populations decline rapidly with cooler water temperatures (Buckingham and Bennett 1996).




Figure 7. Damage inflicted on hydrilla leaf by a first instar of *Parapoynx diminutalis* Snellen.

Credits: Lyle J. Buss, University of Florida.

Importance as a Biological Control Agent

The moth was identified in Pakistan and India during scouting trips to locate potential biological control agents for hydrilla (Baloch et al. 1980). At this point, the researchers observed the moth damaging hydrilla and believed that the moth could be an effective control agent due to its destructive capabilities.

An important characteristic of a biocontrol agent is host specificity. When laying eggs, female moths are not highly selective, which makes other plants susceptible to consumption by developing larvae. Furthermore, the moth is limited by winter temperatures, and populations decline during the cooler months to a level that is almost undetectable. Sensitivity to cooler climates and lack of host specificity makes the moth a poor biological control agent of hydrilla (Habeck and Balciunas 1976; Buckingham and Bennett 1989).

Monitoring/Management

Monitoring for adult moths can be done using ultraviolet (UV) black lights or incandescent light bulbs, which are both attractive to the moth (Buckingham and Bennett 1996).

Hydrilla is invasive, and the actions of the moth rarely require management and are usually considered to be desirable. However, in certain situations where the presence of

hydrilla is needed, such as in research with other biocontrol agents, management of the moth larvae may be necessary to prevent consumption of the plant material. In these situations, a strain of the biorational insecticide *Bacillus thuringiensis* has been tested for controlling *Parapoynx diminutalis*. *Bacillus thuringiensis* subspecies *kurstaki*, commonly known as Btk, is specific to lepidopteran pests. Btk produces proteins that are toxic to larvae; the proteins bind to the midgut when consumed and kill the larvae (Bauce et al. 2006; Van Driesche et al. 2008). A commercially available Btk product has been shown to cause 80% mortality of *Parapoynx diminutalis* larvae in about four days (Buckingham and Bennett 1996).

Selected References

- Agassiz D. 1978. Five introduced species, including one new to science, of China mark moths (Lepidoptera: Pyralidae) new to Britain. *Entomologist's Gazette* 29: 117-127.
- Agassiz D. 1981. Further introduced China mark moths (Lepidoptera: Pyralidae) new to Britain. *Entomologist's Gazette* 32: 21-26.
- Balciunas JK, Habeck DH. 1981. Recent range extension of a hydrilla damaging moth, *Parapoynx diminutalis* (Lepidoptera: Pyralidae). *Florida Entomologist* 64: 195-196.
- Balciunas JK, Minno MC. 1985. Insects damaging hydrilla in the USA. *Journal of Aquatic Plant Management* 23: 77-83.
- Baloch GM, Sana-Ullah, Ghani MA. 1980. Some promising insects for the biological control of *Hydrilla verticillata* in Pakistan. *Tropical Pest Management* 26: 194-200.
- Bauce E, Kumbasli M, Van Fankenhuyzen K, Carisey N. 2006. Interactions among white spruce tannins, *Bacillus thuringiensis* subsp. *kurstaki*, and spruce budworm (Lepidoptera: Tortricidae), on larval survival, growth and development. *Journal of Economic Entomology* 99: 2038-2047.
- Brou Jr VA. 1993. Range extension of the moth *Parapoynx diminutalis* Snellen (Lepidoptera: Pyralidae). *Southern Lepidopterists' News* 15: 33-34.
- Buckingham GR, Bennett CA. 1989. Laboratory host range of *Parapoynx diminutalis* (Lepidoptera: Pyralidae), an Asian aquatic moth adventive in Florida and Panama on *Hydrilla verticillata* (Hydrocharitaceae). *Environmental Entomology* 18: 526-530.
- Buckingham GR, Bennett CA. 1996. Laboratory biology of an immigrant Asian moth, *Parapoynx diminutalis* (Lepidoptera: Pyralidae), on *Hydrilla verticillata* (Hydrocharitaceae). *Florida Entomologist* 79: 353-363.

- Del Fosse ES, Perkins BD, Steward KK. 1976. A new record for *Parapoynx diminutalis* (Lepidoptera: Pyralidae), a possible biological control agent for *Hydrilla verticillata*. Florida Entomologist 59: 1-30.
- Haag KH, Buckingham GR. 1991. Effects of herbicides and microbial insecticides on the insects of aquatic plants. Aquatic Plant Management 29: 55-57.
- Habeck DH. 1996. Australian moths for hydrilla control. Vicksburg, Miss: U.S. Army Engineer Waterways Experiment Station. Technical Report. A96: 10.
- Habeck DH, Balciunas JK. 2005. Larvae of Nymphulinae (Lepidoptera: Pyralidae) associated with *Hydrilla verticillata* (Hydrocharitaceae) in North Queensland. Australian Journal of Entomology 44: 354-363.
- Haller WT, Sutton DL. 1975. Community structure and competition between hydrilla and vallisneria. Hyacinth Control Journal 13: 48-50.
- Hetrick SA, Langeland KA. 2012. [Hydrilla management in Florida lakes](#). University of Florida, IFAS Extension. (9 December 2013).
- Kegley SE, Hill BR, Orme S, Choi AH. 2010. [PAN pesticide database](#). Pesticide Action Network, North America (San Francisco, CA. 2010). (9 December 2013).
- Nuss M, Landry B, Vegliante F, Tränkner A, Mally R, Hayden J, Segner A, Li H, Schouten R, Solis MA, Trofimova T, De Prins J, Speidel W. 2003-2013. [Global information system on Pyraloidea](#). (10 January 2014).
- Shibuya J. 1928. The systematic study on the Formosan Pyralidae. Journal of the Faculty of Agriculture, Hokkaido Imperial University, Sapporo, Japan 22: 1-300.
- Van Driesche R, Hoddle M, Center T. 2008. Control of pests and weeds by natural enemies: an introduction to biological control. Blackwell Publishing. Malden, MA.