

# Biological Control of Mosquitoes with Mermithids<sup>1</sup>

E. G. Platzer<sup>2</sup>

**Abstract:** Mermithid nematodes parasitizing mosquitoes have substantial potential for vector control. Studies on the physiological ecology of *Romanomermis culicivora* have defined some of the general requirements of mermithid nematodes and produced general guidelines for the experimental release of mermithids in biological control. Experimental field studies have established the biological control potential of *R. culicivora*, but further development and utilization of this parasite will require a substantial commitment of scientific man-years and funds. **Key words:** Integrated Pest Management, physiological ecology, *Romanomermis culicivora*.

The Phylum Nematoda has five orders with 14 families of obligate insect parasites, but only the Mermithidae have been found in natural populations of mosquitoes (29, 30). Mosquitoes are also utilized by filarioids as intermediate hosts, but these nematodes have no promise for biological control and will not be considered further. There are many reports of mermithids in mosquito populations but, because the nematode identifications were based on larval stages, they are often incomplete or inaccurate. However, there appear to be 15 acceptable species in eight genera (29,30). These mermithids can be divided into two groups: those that complete the parasitic phase of life primarily in the larval stages of the mosquito host and those that complete most of the parasitic growth phase in the adult mosquito (Table 1). In the first group there are four genera and ten species. Of the 133 natural and laboratory infected host species listed, 53% are *Aedes* sp., 20% are *Anopheles* sp., 19% are *Culex* sp., and the remainder of the hosts are species of *Armigeres*, *Culiseta*, *Deinocerites*, *Mansonia*, *Orthopodomyia*, *Psorophora*, *Toxorhynchites*, *Tripteroides*, *Uranotaenia*, or *Wyeomyia* (17,29,30). It may appear obvious from this list that mermithids have had their greatest success as parasites of culicine mosquitoes, particularly those in the genus *Aedes*. However, the list probably more accurately reflects the geographic distribution of entomologists interested in biological control organisms and, therefore, such an assumption is probably premature. A similar assumption, and subsequent explanation can be made for hosts of mermithids

maturing in adult mosquitoes. There, of 14 host species 86% are *Aedes* sp.

## HOST SPECIFICITY

In most cases, we have little information on host specificity other than that the hosts are generally mosquitoes. Some exceptions are *Hydromermis churchilliensis*, which naturally infects chaoborid midges as well as *Aedes* sp., and *Romanomermis culicivora*, which also can infect blackfly larvae under abnormal conditions (30). These examples appear to be extraordinary exceptions to the specificity for mosquito larvae. Specificity within mosquitoes has received little attention. *Perutilimermis culicis* demonstrates species specificity and parasitized only *Aedes sollicitans*; *Strelkovimermis peterseni* demonstrates generic specificity and parasitized only *Anopheles* sp. (13).

The most thorough examination of host specificity has been carried out mainly by Dr. Petersen with *R. culicivora* (17,30). More than 82 species of mosquitoes in 13 genera have been infected in the laboratory or found infected under natural conditions. *R. culicivora* was unable to complete its development in five mosquito species: *Aedes triseriatus*, *Anopheles sinensis*, *Culex territans*, *Mansonia uniformis*, and *Psorophora ferox*.

## ENVIRONMENTAL LIMITATIONS

Most investigations of environmental limitations of mermithids as biological control agents have used *R. culicivora*, but the observations are probably valid for other temperate zone mermithids.

**Temperature:** Petersen (13) reported that *R. culicivora* was infective when air temperatures were above 15 C. Subsequently, Brown and Platzer (1) and Gallo-way and Brust (8) found that some infec-

Received for publication 12 December 1980.

<sup>1</sup>Symposium paper presented at the annual meeting of the Society of Nematologists, New Orleans, Louisiana, August 1980.

<sup>2</sup>Department of Nematology, University of California, Riverside, CA 92521.

Table 1. List of mermithids, their hosts, and sites of collection.\*

| Mermithid   | Mosquito host   | Locality                 |
|---|---|--------------------------|
| Mermithids maturing primarily in larval stages of the host: |   |                          |
| <i>Hydromermis churchilliensis</i>                          | <i>Aedes</i> —3 sp.   | Manitoba                 |
| <i>Octomyomermis muspratti</i>                              | <i>Aedes</i> —8 sp.<br><i>Anopheles</i> —2 sp.<br><i>Culex</i> —5 sp.   | Zambia                   |
| <i>Octomyomermis troglodytis</i>                            | <i>Aedes sierrensis</i>   | California               |
| <i>Romanomermis culicivorax</i>                             | <i>Aedes</i> —29 sp.<br><i>Anopheles</i> —3 sp.<br><i>Armigeres subalbatus</i><br><i>Culex</i> —18 sp.<br><i>Culiseta</i> —4 sp.<br><i>Deinocerites pseudus</i><br><i>Mansonia</i> —2 sp.<br><i>Orthopodomyia signifera</i><br><i>Psorophora</i> —7 sp.<br><i>Toxorhynchites rutilus</i><br><i>Tripteroides bambusa</i><br><i>Uranotaenia</i> —3 sp.<br><i>Wyeomyia smithii</i> | Florida,<br>Louisiana    |
| <i>Romanomermis communensis</i>                             | <i>Aedes</i> —4 sp.   | Manitoba                 |
| <i>Romanomermis hermaphrodita</i>                           | <i>Aedes</i> —2 sp.   | Manitoba                 |
| <i>Romanomermis iyengari</i>                                | <i>Anopheles subpictus</i>  | India                    |
| <i>Romanomermis kiktoreak</i>                               | <i>Aedes</i> —5 sp.   | Northwest Territories    |
| <i>Romanomermis nielseni</i>                                | <i>Aedes</i> —5 sp.<br><i>Culex</i> —2 sp.  | Wyoming                  |
| <i>Strelkovimermis peterseni</i>                            | <i>Anopheles</i> —9 sp.   | Louisiana                |
| Mermithids maturing primarily in adult stages of the host:  |   |                          |
| <i>Culicimermis schakhovii</i>                              | <i>Aedes</i> —6 sp.   | USSR                     |
| <i>Culicimermis</i> sp.                                     | <i>Aedes</i> —4 sp.   | Manitoba                 |
| <i>Empidomermis cozii</i>                                   | <i>Anopheles funestus</i>   | West Africa              |
| <i>Paramermis canadensis</i>                                | <i>Aedes</i> —2 sp.   | British Columbia         |
| <i>Perutillimermis culicis</i>                              | <i>Aedes sollicitans</i>  | Louisiana,<br>New Jersey |

\*Derived from Petersen and Chapman (17) and Poinar (29,30)

tions occurred at water temperatures of 12 and 10 C, respectively. However, the optimum temperature was in the 21–33 C range. In addition, the optimal temperature for development of the parasitic stage of *R. culicivorax* was in the 20–32 C range. Petersen (14) found that *Romanomermis nielseni* postparasites developed and completed oviposition within 7 wk at 17 C. Increasing the temperature to 23 C suppressed oviposition but not postparasite development. In contrast, *R. culicivorax* required 23 wk to complete oviposition at 17 C, but only 3–7 wk at 23 C. This information provided a unique physiological comparison between the two species. Temperature is an important consideration in the use of *R. culicivorax* as a biological control agent; its use in colder temperature zones is precluded (8).

*pH*: Petersen (15) found that *R. culicivorax* was fully infective from pH 5.4 to 7.9 (4.8 to 8.5 were transitory pH exposures). Hence, pH of most natural waters should not be a limiting factor.

*Salts*: Petersen and Willis (20) reported that mild salinity (0.04M NaCl) inhibited the infectivity of *R. culicivorax*. This was confirmed by Brown and Platzer (2) who also reported the hierarchy of ion toxicity on a molar basis for *R. culicivorax* as follows: cations, sodium < potassium < calcium; and anions, chloride < carbonate = sulfate < nitrate < nitrite < phosphate. Therefore, it appeared that *R. culicivorax* would not be an effective biocontrol agent in feedlot runoffs, fertilizer plant wastewater, and brackish water situations. Recently, Petersen (16) has found that *Octomyomermis muspratti* is tolerant of diluted

seawater (3,000–4,000  $\mu\text{mhos/cm}$ ) and water from tree holes (10,000  $\mu\text{mhos/cm}$ ; high in organic matter). These findings suggest that *O. muspratti* has great potential for mosquito biocontrol in polluted waters.

*Oxygen:* In investigations to determine why polluted water compromises the infectivity of *R. culicivora*x, Brown and Platzer (3) reported the effects of lowered oxygen availability on the infective nematode. Transient exposure to low  $\text{O}_2$  tension increased the survival, and thereby the infectivity, of the preparasites of *R. culicivora*x. This effect was explained as resulting from induced quiescence under low  $\text{O}_2$  tensions and hence greater viability under the test conditions. More recently, I have found that the preparasites of *R. culicivora*x stop moving within 8 h in water high in organic content but low in  $\text{O}_2$  (6). Therefore, lowered oxygen tensions in polluted waters may be responsible for the inability of *R. culicivora*x to infect mosquitoes under such conditions.

#### INTEGRATED PEST MANAGEMENT

*Chemicals:* Integrated Pest Management, the multipronged approach to pest control, encourages assessment of pesticide effects on mermithid infections. The first assessment was that of Mitchell et al. (11) in Taiwan, who found that the usual levels of Abate, Dieldrin, and Gama HCH did not adversely affect *R. culicivora*x infections. Finney et al. (7) reported that Altosid 5E, an insect growth regulator, at concentrations typically used for control of mosquito larvae did not interfere with any phase of the parasite's development. However, host mortality was considerably increased when Altosid 5E and *R. culicivora*x were used in combination on *Aedes aegypti* in laboratory experiments. This finding suggests a unique chemical-biological control approach in mosquito control: "If the chemical don't get 'em, the nemas will!" Platzer and Brown (25) reported that a variety of copper-based organic algicides and copper sulfate didn't compromise the infectivity of *R. culicivora*x as long as the concentrations were in the ranges usually used for algae and weed control. In summary, it appears that *R. culicivora*x is tolerant of a variety of chemical pest control measures used in

aquatic ecosystems and combination treatments are possible.

*Invertebrate predators:* Most research on environmental limitations has been directed towards evaluation of abiotic factors. Mitchell et al. (11) were the first to recognize that ostracods preyed on mermithid preparasites. Similarly, copepods and young gammarids will attack copepod preparasites (27), and the dynamics of copepod predation on mermithids has been reported in detail (28). Both satiated and starved adult copepods (*Cyclops vernalis*) attack preparasites rapidly in small volumes of water (less than 2 ml). This attack or predation rate is dependent on the volume of water, thus indicating that searching or hunting behavior controls predation rate. In laboratory studies with larger volumes of water (one liter), it was shown that mermithid preparasite populations were reduced significantly by copepod densities of 20–100/liter. Although field studies have not been carried out, I suspect that copepod density might play a significant role in the success or failure of field applications.

Invertebrate predators also may reduce the potential of establishment of aquatic mermithids (27). Diving beetles and gammarids, dragonfly and damselfly naiads, and small crayfish attack mermithid postparasites rapidly (27). Isopods, however, will attack postparasites only if other food is limited. Such predation may account for poor or no establishment of the mermithid after field application of *R. culicivora*x.

#### FIELD APPLICATIONS

Essentially all field releases of mermithid parasites of mosquitoes have been conducted with *R. culicivora*x, although in one study, Petersen and Willis (22) released *S. peterseni* and after 5 yr obtained more than 80% parasitization. In 1971 Petersen and Willis (21) treated 10 natural sites 20 times with preparasites of *R. culicivora*x and obtained 65, 58, and 33% parasitism of second, third, and fourth instars, respectively, of the *Anopheles* sp.; 94% of second-instar anophelines and 64% of all *Anopheles* were infected at rates of 1,000 preparasites/ $\text{m}^2$  of surface area application. Application rates less than 1,000 preparasites/ $\text{m}^2$  of surface area were less effective, but higher

application rates did not increase the number of infected mosquito larvae. Establishment of the nematode was observed in 7 of the 10 sites.

Petersen et al. (19) treated fallow rice fields in Louisiana infested with *Psorophora confinnis* and *Anopheles quadrimaculatus*. Increasing the application rate from 180 to 1,450 preparasites/yard<sup>2</sup> of surface area increased the parasitism of *P. confinnis* from 10 to 38%. They estimated that 3,900 preparasites/yard<sup>2</sup> would give 95% infection. Sixteen percent of *A. quadrimaculatus* were infected at the rate of 181 preparasites/yard<sup>2</sup> and 61% at 724/yard<sup>2</sup> of surface area. They estimated that 1,300 preparasites/yard<sup>2</sup> of surface area would give 95% infection. Caution should be used in making this type of prediction. Brown et al. (4), for example, found that increasing treatment rates from 1,000 to 25,000 preparasites/m<sup>2</sup> of surface area did not significantly increase the number of *Culex tarsalis* larvae infected.

Petersen and Willis (22) made a total of 30 releases in 21 sites (15 at rates of 1,000 and 15 at rates of 2,000 mermithid preparasites/yard<sup>2</sup> of surface area) to control *Anopheles* larvae. The primary species present was *Anopheles crucians*. At the lower rate, an average of 76% of the hosts were infected and parasitism averaged 60, 80, 86 and 77% in first through fourth instars, respectively. At 2,000 preparasites/yard<sup>2</sup>, an average of 85% of the larvae were infected. Poor correlation was found between infection rates and vegetation and water depth.

Brown et al. (4) conducted field tests against four species of mosquitoes in three natural (rice fields and ponds) and two artificial (1- and 6-m<sup>2</sup> ponds) habitats at treatment levels ranging from 706 to 25,000 preparasites/m<sup>2</sup> of surface area. Anopheline larvae were more susceptible than culicine. Although both *C. tarsalis* and *Culiseta inornata* were reported in laboratory studies to be good hosts for *R. culicivora*, field studies with these two species were disappointing.

Levy and Miller (10) applied *R. culicivora* at the rate of 3,600 preparasites/m<sup>2</sup> of surface area to control mosquitoes breeding in a grassy field. In 10 potholes and ditches sampled in the field, they found 88–100% infection in *Culex nigripalpus*,

*Aedes taeniorhynchus*, *Psorophora columbiana*, and *Psorophora ciliata*. In some cases all larvae were killed within 24 h due to multiple parasitism.

Levy and Miller (9) released preparasites into two unused sewage settling tanks at 64,000 and 110,000 preparasites/m<sup>2</sup> of surface area and obtained 37 and 54% infection of *C. pipiens quinquefasciatus*, respectively. Although these treatment levels seem high, large numbers of *C. p. quinquefasciatus* larvae were present and host: parasite treatment ratios were approximately 5:1 and 3:1. High concentrations of phosphates, chlorides, and carbonates probably lowered the infectivity of the preparasites, as much higher infection levels would be expected in this species at the dosages applied.

Petersen et al. (18) successfully controlled *Anopheles albimanus* and *Anopheles punctipennis* in Lake Apostepeque, El Salvador, with *R. culicivora*. The mosquito breeding areas, a 2–5 meter band (total area = 10,700 m<sup>2</sup>) around the lake was treated 11 times during 7 wk with 2,400–4,800 preparasites/m<sup>2</sup> at each application. The *Anopheles* population declined from 10 per sample dip prior to the first preparasite application to 0.6 per dip by the end of the release period; a 17-fold reduction in the number of mosquito larvae. This was the first successful large-scale attempt to control mosquitoes with a parasite.

Another technique for application of *R. culicivora* was reported by Petersen and Willis (23) who distributed the eggs and postparasitic stages of *R. culicivora* in 13 habitats known to breed pasture mosquitoes. They reported 52% of *Aedes atlanticus*, 59% of *Aedes tormentor*, 38% of *P. columbiana*, and 51% of *Psorophora howardii* were parasitized after the pastures were flooded. Brown-Westerdahl et al. (5) found that early season application of *R. culicivora* postparasites was effective in providing continuous partial control (weekly mean infection = 60%) of *A. freeborni* and *C. tarsalis* throughout the rice growing season in Northern California. Infections were observed up to 12 m from point of application of 1,500 postparasites. The nematodes overwintered, and infected sentinel and native mosquitoes were found

the following season. *Platzer* and *Eby* (26) also showed that postparasites can produce sufficient eggs to persist for two seasons of mosquito control.

In field releases of mermithid nematodes, it is important that the application be made by trained personnel who thoroughly understand the environmental limitations of the organism. Under such conditions, the full potential for biological control will be achieved with mermithid nematodes.

### ESTABLISHMENT

*Petersen* and *Willis* (26) have conducted the only long-term studies on the ability of *R. culicivora* to become established after application. They showed that this nematode can become established in many semi-permanent and permanent mosquito breeding sites, but that certain problems existed: lack of host populations, periodic lack of breeding water in the semipermanent sites, and inaccessibility due to changes in site access. Periods of little or no water tended to reduce the levels of infection in susceptible hosts, and flushing of the sites by heavy rains reduced nematode populations. It also appeared that sites which produced most mosquitoes also produced most *R. culicivora* infections. *Petersen* and *Willis* (24) noted that recycling occurred in 10 sites 6–29 wk after treatment with preparasites. In 1974 three of the sites treated in 1971 were still producing infected hosts (7–25%). Five of six sites treated in 1971 and 1973 and 7 of 12 sites treated only in 1973 were producing parasitized hosts in 1974.

*Petersen* and *Willis* (23) released *S. peterseni* in Louisiana in 1971. Two years later 88% of the *Anopheles* sp. were parasitized. *Nickle* (12) reported that *R. culicivora* was established and overwintered in Maryland where winter temperatures dropped as low as –17 C.

### SUMMARY

Although substantial potential has been attributed to the effectiveness of mermithids for vector control, more time and research is required to realize this potential. Additional studies on the biology of *R. culicivora* and other mermithids are needed if

we are to fully utilize the biocontrol potential of these nematodes. A greater commitment of scientific and economic resources are required. Biocontrol with *R. culicivora* has been under development for 10 yr; at the current level of interest and funding it will probably require 10 more years to obtain effective use of this or alternate organisms in biological control of vector populations. In addition, we need a strong commitment from agencies to train personnel in the handling and use of such biological control organisms.

### LITERATURE CITED

1. *Brown, B. J., and E. G. Platzer.* 1977. The effects of temperature on the infectivity of *Romanomermis culicivora*. *J. Nematol.* 9:166-172.
2. *Brown, B. J., and E. G. Platzer.* 1978. Salts and the infectivity of *Romanomermis culicivora*. *J. Nematol.* 10:53-64.
3. *Brown, B. J., and E. G. Platzer.* 1978. Oxygen and the infectivity of *Romanomermis culicivora*. *J. Nematol.* 10:110-113.
4. *Brown, B. J., E. G. Platzer, and D. S. Hughes.* 1977. Field trials with the mermithid nematode, *Romanomermis culicivora*, in California. *Mosq. News.* 37:603-608.
5. *Brown-Westerdahl, B., R. K. Washino, and E. G. Platzer.* 1979. Early season application of *Romanomermis culicivora* provides continuous partial control of rice field mosquitoes. *Proc. Calif. Mosq. Vector Cont. Assoc.* 47:55 (Abstr.).
6. *Dhillon, M. S., M. S. Mulla, and E. G. Platzer.* 1980. Evaluation of nematode *Romanomermis culicivora* against cemetery mosquitoes. *Mosq. News.* 40:531-535.
7. *Finney, J. R., R. Gordon, W. J. Condon, and T. N. Rusted.* 1977. Laboratory studies on the feasibility of integrated mosquito control using an insect growth regulator and a mermithid nematode. *Mosq. News* 37:6-11.
8. *Galloway, T. D., and R. A. Brust.* 1977. Effects of temperature and photoperiod on the infection of two mosquito species by the mermithid, *Romanomermis culicivora*. *J. Nematol.* 9:218-221.
9. *Levy, R., and T. W. Miller, Jr.* 1977. Experimental release of a mermithid to control mosquitoes breeding in sewage settling tanks. *Mosq. News* 37:410-414.
10. *Levy, R., and T. W. Miller, Jr.* 1977. Experimental release of *Romanomermis culicivora* (Mermithidae: Nematoda) to control mosquitoes breeding in Southwest Florida. *Mosq. News* 37:483-486.
11. *Mitchell, C. J., P. Chen, and H. C. Chapman.* 1974. Exploratory trials utilizing a mermithid nematode as a control agent for *Culex* mosquitoes in Taiwan. *J. Formosan Med. Assoc.* 73:241-254.
12. *Nickle, W. R.* 1979. Probable establishment and overwintering of a mermithid nematode parasite of mosquitoes in Maryland. *Proc. Helminthol. Soc. Wash.* 46:21-27.

13. Petersen, J. J. 1973. Role of mermithid nematodes in biological control of mosquitoes. *Exp. Parasitol.* 33:239-247.
14. Petersen, J. J. 1976. Comparative biology of the Wyoming and Louisiana populations of *Reesimermis nielsenii*, parasitic nematode of mosquitoes. *J. Nematol.* 8:273-275.
15. Petersen, J. J. 1979. pH as a factor in parasitism of mosquito larvae by the mermithid *Romanomermis culicivorax*. *J. Nematol.* 11:105-106.
16. Petersen, J. J. 1981. Observations on the biology of *Octomyomermis muspratti*, a nematode parasite of mosquitoes. *J. Invert. Pathol.*, in press.
17. Petersen, J. J., and H. C. Chapman. 1979. Checklist of mosquito species tested against the nematode parasite *Romanomermis culicivorax*. *J. Med. Entomol.* 15:468-471.
18. Petersen, J. J., H. C. Chapman, O. R. Willis, and T. Fukuda. 1978. Release of *Romanomermis culicivorax* for the control of *Anopheles albimanus* in El Salvador. II. Application of the nematode. *Am. J. Trop. Med. Hyg.* 27:1268-1273.
19. Petersen, J. J., C. D. Steelman, and O. R. Willis. 1973. Field parasitism of two species of Louisiana rice field mosquitoes by a mermithid nematode. *Mosq. News* 33:573-575.
20. Petersen, J. J., and O. R. Willis. 1970. Some factors affecting parasitism by mermithid nematodes in southern house mosquito larvae. *J. Econ. Entomol.* 63:175-178.
21. Petersen, J. J., and O. R. Willis. 1972. Results of preliminary field application of *Reesimermis nielsenii* (Mermithidae: Nematoda) to control mosquito larvae. *Mosq. News* 32:312-316.
22. Petersen, J. J., and O. R. Willis. 1974. Experimental release of a mermithid nematode to control *Anopheles* mosquitoes in Louisiana. *Mosq. News* 34:316-319.
23. Petersen, J. J., and O. R. Willis. 1974. *Diximermis peterseni* (Nematoda: Mermithidae): A potential biocontrol agent of *Anopheles* mosquito larvae. *J. Invert. Pathol.* 24:20-23.
24. Petersen, J. J., and O. R. Willis. 1975. Establishment and recycling of a mermithid nematode for the control of mosquito larvae. *Mosq. News* 35:526-532.
25. Platzer, E. G., and B. J. Brown. 1976. Physiological ecology of *Reesimermis nielsenii*. *Proc. Int. Colloq. Invert. Pathol.*, Kingston, Canada, pp. 263-267.
26. Platzer, E. G., and J. E. Eby. 1980. Mermithid nematodes in mosquito control: persistence in small experimental ponds. *Proc. Calif. Mosq. Vector Cont. Assoc.* 48:40-41.
27. Platzer, E. G., and L. L. MacKenzie-Graham. 1978. Predators of *Romanomermis culicivorax*. *Proc. Calif. Mosq. Vector Cont. Assoc.* 46:93 (Abstr.).
28. Platzer, E. G., and L. L. MacKenzie-Graham. 1980. *Cyclops vernalis* as a predator of the preparasitic stages of *Romanomermis culicivorax*. *Mosq. News* 40:252-257.
29. Poinar, G. O., Jr. 1975. *Entomogenous Nematodes*. Leiden: E. J. Brill.
30. Poinar, G. O., Jr. 1979. *Nematodes for Biological Control of Insects*. Boca Raton, Florida: CRC Press.