

Bibliography on *Hydrellia griseola* Fallén (Diptera: Ephydriidae) and review of its biology and pest status

Louis S. Hesler

10th MEDLAB, Box 1041
Department of the Army
APO AE 09180

Abstract: A bibliography is presented on *Hydrellia griseola* (Fallén), a cosmopolitan species of shore fly that is an agricultural pest throughout much of its range. The literature review includes 195 references and spans 180 years: from 1813, when *H. griseola* was first described (as *Notophila griseola*), to 1993. My review updates and extends by 110 years a previous review of this species. The host plant associations, bionomics, distribution, pest status, and control of this species are reviewed. The known host plants of larval *H. griseola* are listed. My survey of the literature supports the hypothesis that *H. griseola* is increasing its range as a pest of rice. *H. griseola* can be controlled by cultural or chemical methods, but due to agronomic and environmental constraints on these methods, my review points out the need for more research on and use of the natural enemies of *H. griseola*.

Introduction

Hydrellia Robineau-Desvoidy (1830) (Diptera: Ephydriidae: Notophilinae) is the most species rich genus of shore flies, and one of the most widely distributed genera in its family (Deonier, 1971; Zatwarnicki, 1988). *Hydrellia* contains about 11% of all species of shore flies and is recorded from all continents except Antarctica. Five species (*fascitibia* v. Roser, *griseola* [Fallén], *ischiaca* Loew, *tenebricosa* Collin, and *tibialis* Cresson) are Holarctic, and at least five (*griseola*, *ischiaca*, *philippina* Ferino, *prosternalis* Deeming, and *sasakii* Yuasa and Isitani) are economically important (Zatwarnicki, 1988). Also, *H. pakistanae* Deonier (from southern Asia) and *H. balciunasi* Bock (from Australia) have been introduced into the United States for biological weed control (Buckingham *et al.* 1989, 1991).

H. griseola (the smaller rice leafminer) is the most widely distributed and one of the most economically significant species (Joshi *et al.*, 1986; Zatwarnicki, 1988). The literature on *H. griseola* dates back to 1813, when Fallén originally described this species as *Notophila griseola*. Macquart (1835) later placed it in the genus *Hydrellia*. Since then, a considerable body of literature has accumu-

lated on this species. Joshi *et al.* (1986) developed a bibliography limited to the years 1913-1983 on *Hydrellia* spp. that attack rice. Grigarick (1959a) and Deonier (1971) reviewed the biology of *H. griseola*.

However, an expanded and updated bibliography on *H. griseola* is needed. Ideally, this bibliography should cover the entire body of literature to date. Second, an updated review of the literature on *H. griseola* is also warranted.

Therefore, I present an expanded bibliography on *H. griseola*, including references from 1813 through early 1993. In addition to the bibliography, I summarize the literature, presenting a brief review of the distribution, host plant associations, bionomics, pest status, and control of this species. I discuss unresolved issues regarding the distribution of *H. griseola*, note the increasing recognition of this species as a pest of rice, and point out the need for more research on the natural enemies of *H. griseola*.

Distribution, host range, and bionomics

In the Palearctic, *H. griseola* occurs throughout Eurasia and insular regions of southeast Asia (Barrión and Litsinger, 1981; Zatwarnicki, 1988).

In the western hemisphere, *H. griseola* is found throughout the United States and Canada (Grigarick, 1959a; Deonier, 1971), and in Costa Rica (Cresson, 1918), Uruguay (Parker *et al.*, 1951-52), and Colombia (Weber *et al.*, 1988). The apparent discontinuous distribution in Latin America and South America may simply be due to incomplete information on the geographic distribution of *H. griseola*, or perhaps it may indicate introductions of this species at various sites.

The distribution of *H. griseola* in Africa and Australia is uncertain. Willcocks (1925) originally reported that *H. griseola* occurs in Egypt, but recent reports indicate that the species may actually be *H. prosternalis* (Ismail *et al.*, 1979; Isa, 1989). Halfpapp (1989) reported that *H. griseola* had been found in Queensland (Australia), but Bock (1990) asserted that specimens collected by Halfpapp represented a new sibling species (*i.e.*, *H. mareeba* Bock). More intensive surveys for *H. griseola* on these two continents is needed.

The wide distribution of *H. griseola* reflects its tolerance for a great range of habitats and environmental conditions. Adult activity and development of the immature instars occur at temperatures ranging from about 10° to 40°C (Grigarick, 1959a). Altitudinally, *H. griseola* has been collected from 87 m below sea level (Death Valley, USA) to 2950 m above (Mount Timpanogos, Utah, USA) (Deonier, 1971).

H. griseola occurs in various ephemeral habitats of aquatic, semi-aquatic, and terrestrial ecosystems (Grigarick, 1959a; Deonier, 1964b, 1971; Scheiring and Foote, 1973). This species occurs in natural settings as well as anthropogenic habitats (*e.g.*, rice paddies and other agroecosystems, canals, artificial ponds) (Grigarick, 1959a; Deonier, 1964b, 1971; Scheiring and Foote, 1973). Methods are described for sampling the different life stages of *H. griseola* in various habitats (see Grigarick, 1959a, b; Deonier, 1971; Deonier, 1972b).

H. griseola is a polyphagous plant-feeder in the larval stage (Grigarick, 1959a; Deonier, 1964b; 1971). The majority of larval host plants are members of the family Graminae (Table 1). Larvae are endophytophagous, burrowing into leaves and leaf sheaths (Grigarick, 1959a; Deonier, 1971; Manandhar and Grigarick, 1983). The larvae feed on leaf mesophyll, and this feeding creates mines between the epidermal layers of the leaf. Individual larvae may mine more than one leaf or sheath. Pupariation may occur within leaf mines or off of plants altogether (*e.g.*, at the bottom of ice-covered

pools [Burghel, 1959a]). Often, development is completed on more than one plant species (Deonier, 1971). In Japan, Kuwayama (1955) found that both larval and puparial mines were present on monocotyledonous plants, whereas only puparial mines were found on dicotyledonous plants. Total developmental time ranges from 13 d at 32-35 °C to 94 d at 10 °C (Grigarick, 1959a). Up to 11 generations have been estimated to occur in a single year in California (Grigarick, 1959a).

Adults have a varied diet. They feed on, among other things, fungi, algae, diatoms, nectar, and leaf epidermis (Deonier, 1971; 1972a). They also prey and scavenge upon small insects such as small flies (including other *Hydrellia*), Psilidae, early instar Ephemeroptera and Odonata, Collembola, and Braconidae (Grigarick, 1959a; Deonier, 1971).

Under optimal conditions, adults may be quite long-lived. Grigarick (1959a) was able to maintain adult males for up to 130 d and females up to 139 d. He also found that adults mate as early as 3 d and as late as 70 d after emergence. Females can oviposit as early as 5 d and as late as 93 d after emergence (Grigarick, 1959a). Eggs of *H. griseola* are laid singly on the upper surfaces of leaves and require high humidity (80-100%) for hatching (Grigarick, 1959a).

The overwintering habits of *H. griseola* vary geographically. *H. griseola* overwinters primarily as adults in California (Grigarick, 1959a) and as adults in the Black Sea region of the former USSR (Kas'yanov, 1967). In Hungary, only parthenogenetic females are known to overwinter (Szito, 1976). By contrast, *H. griseola* overwinters in puparia in the Maritime Province of the former USSR and in northern temperate regions of Japan (*i.e.*, Hokkaido district, Honshu) (Kuwuyama, 1956; Klimanova, 1971). In southern Japan (*i.e.*, south Ibaraki Prefecture), all life stages can be found during the winter months (Iwamoto, 1966).

Pest status

H. griseola is a pest of several crops, including rice (*Oryza sativa* L.), wheat (*Triticum aestivum* L.), barley (*Hordeum vulgare* L.), oats (*Avena sativa* L.), timothy (*Phleum pratense* L.), and onions (*Allium* spp.) (see Grigarick, 1959a and references therein). Larvae develop also on maize (*Zea mays* L.) and wild rice (*Zizania aquatica* L.) (Grigarick, 1959a; Schmidt, 1962), but economically significant infestations of *H. griseola* have not been reported on these two crops. Injury to plants results from leaf- and stem-mining by the larvae. Their

Table 1: Principal host plants of *Hydrellia griseola* larvae, based on host plant records in Grigarick (1959a) and Deonier (1971).

| | | |
|-------------------------------------|---|--|
| Alismataceae | Graminae cont. | Graminae cont. |
| <i>Alisma</i> spp. | <i>Calamagrostis</i> sp. | <i>Polypogon</i> sp. |
| <i>Damasonium</i> sp. | <i>Catabrosa</i> sp. | <i>Scleropa</i> sp. |
| <i>Sagittaria</i> spp. | <i>Cynodon</i> sp. | <i>Secale cereale</i> L. |
| Caryophyllaceae | <i>Dactylis glomerata</i> L. | <i>Setaria</i> sp. |
| <i>Lychnis</i> sp. | <i>Digitaria</i> sp. | <i>Triticum sativum</i> L. |
| <i>Stellaria media</i> (L.) | <i>Echinochloa crusgalli</i> (L.) Beauv. | <i>Zea mays</i> L. |
| Chenopodiaceae | <i>Eleusine</i> sp. | <i>Zizania aquatica</i> L. |
| <i>Kochia</i> sp. | <i>Eragrostis</i> sp. | Hydrocharitaceae |
| Compositae | <i>Festuca pratensis</i> Huds. | <i>Hydrocharis morsus-ranae</i> L. |
| <i>Bellis</i> sp. | <i>Gaudinia</i> sp. | <i>Stratiotes aloides</i> ¹ |
| Cruciferae | <i>Glyceria</i> sp. | Labiatae |
| <i>Nasturtium officinale</i> R. Br. | <i>Hierochloe</i> sp. | <i>Lamium purpureum</i> L. |
| Cyperaceae | <i>Holcus lanatus</i> L. | Leguminosae |
| <i>Carex</i> sp. | <i>Hordeum vulgare</i> L. | <i>Trifolium</i> sp. |
| <i>Cyperus</i> sp. | <i>Lagarus</i> sp. | Lemnaceae |
| <i>Scirpus</i> sp. | <i>Lolium perenne</i> L. | <i>Lemna minor</i> L. |
| Graminae | <i>Muhlenbergia</i> sp. | Liliaceae |
| <i>Agropyron</i> spp. | <i>Oryza sativa</i> L. | <i>Allium cepa</i> L., <i>A. porrum</i> L. |
| <i>Agrostis</i> sp. | <i>Panicum</i> sp. | Polygonaceae |
| <i>Alopocerus</i> sp. | <i>Paspalum</i> sp. | <i>Polygonum</i> sp. |
| <i>Anthoxanthum odoratum</i> L. | <i>Phalaris arundinacea</i> L. | Scrophulariaceae |
| <i>Apera</i> sp. | <i>Phleum pratense</i> L. | <i>Veronica</i> sp. |
| <i>Avena sativa</i> L. | <i>Phragmites</i> sp. | Typhaceae |
| <i>Brachypodium</i> sp. | <i>Poa annua</i> L., <i>P. pratensis</i> L., <i>P. trivialis</i> L. | <i>Typha latifolia</i> L. |
| <i>Bromus</i> sp. | | |

¹Undetermined author.

feeding reduces photosynthesis, and causes leaf abscission and sometimes seedling death (Manandhar and Grigarick, 1983).

Early notoriety of *H. griseola* was due to its economic significance in small grain crops. Lilljeborg (1861) first reported *H. griseola* as a pest of barley, oats, and timothy in Sweden. Crop losses near 50% have been attributed to *H. griseola* in barley, oats, and wheat in Europe (Balachowsky and Mesnil, 1935).

During the 20th century, however, *H. griseola* has become increasingly recognized as a pest of rice. The first reports of *H. griseola* attacking rice came from California (de Ong, 1922). Later, Nepveu and D'Aguillar (1951) reported *H. griseola* as a pest of rice in France. Serious outbreaks of *H. griseola* in rice occurred in California in 1953 (Lange *et al.*, 1953) and in Japan in 1956 (Kuwayama, 1956). The 1953 outbreak in California resulted in a 10-20% crop loss worth \$16 million (US); an additional \$1.2 million were spent on insecticides for control

(Jensen, 1954). The 1956 infestation in Japan caused a loss of 39,000 kl of rice over a 75,000-ha area (Kuwayama, 1956).

The economic significance of *H. griseola* varies geographically depending on the crop. For instance, *H. griseola* is a pest of rice throughout most of its range. In contrast, its pest status on other crops is limited geographically to Europe and Asia, as the literature is devoid of any reference to serious infestations by *H. griseola* in North and South America in crops other than rice. Grigarick (1959a) did not find *H. griseola* infesting wheat and barley in California, but he was able to rear specimens from these two cereals, as well as oats, under laboratory conditions. Economically significant infestations of onions by *H. griseola* are apparently limited to European regions of the former USSR (Isaev, 1931).

Much of the recent literature on *H. griseola* is related to its occurrence on rice. Since 1980, 30 of the 38 reports about *H. griseola* have dealt exclu-

sively with its economic significance in rice, and two additional reports addressed the pest status of *H. griseola* on rice and at least one other crop. Moreover, the geographic range in which *H. griseola* is recognized as a pest of rice has been increasing. Since 1980, several regions have reported their first accounts of economic loss in rice due to *H. griseola*, including The Philippines (Barrión and Litsinger, 1981), the southern US (Way *et al.*, 1983), India (Chakravarthy, 1987), Colombia (Weber *et al.*, 1988), and perhaps Australia (Halfpapp, 1989; but see Bock, 1990).

H. griseola is likely to receive increased attention as a pest, particularly in rice-growing regions of the US, because of changes in management practices. For instance, the practice of continuous, deep-water culture for weed control may become more prevalent in California rice production due to 1) increasing restrictions on herbicide use in rice fields (Williams *et al.*, 1990), and 2) the adoption of organic (*i.e.*, non-chemical) management practices by some rice growers (Altieri *et al.*, 1983; Hesler *et al.*, 1993). Deep-water culture causes the leaves of rice plants to lie prostrate on the water surface, favoring increased incidence and severity of attack by *H. griseola* (Lange *et al.*, 1953). Maintenance of a shallow water depth (*e.g.*, 7-10 cm) during the late seedling and early tillering stages reduces the potential for damage by *H. griseola* (Grigarick and Washino, 1983; Way *et al.*, 1983; see also Szilvassy and Petrasovits, 1960; Szito, 1985). However, shallow water levels can greatly increase the risk of weed problems (Williams *et al.*, 1990). Because weeds are a serious threat annually to rice production in California (whereas outbreaks of *H. griseola* occur sporadically), growers are likely to continue the use of deep-water culture despite an increased likelihood of problems with *H. griseola*. New means of managing *H. griseola* that are compatible with deep-water culture are clearly needed.

Control

Several means of preventing and controlling infestations of *H. griseola* are available. Many researchers have advocated the integration of cultural, chemical, and biological control methods (Buryi, 1980; Grigarick and Washino, 1983; Weber *et al.*, 1988).

Serious infestations of *H. griseola* can often be prevented by the attention to various cultural factors, such as planting date, field location, and water management (discussed above for rice). For in-

stance, a delay in planting date that minimizes the time in which young plants are exposed to cool, humid weather can significantly reduce the extent of damage caused by *H. griseola* (Hering, 1951; Grigarick and Washino, 1983). In small grains, economically damaging infestations of *H. griseola* can also be prevented by avoiding plantings in low, damp areas near standing or running water (Hering, 1951). In Japan, damage to rice can be limited by avoiding synchrony between the transplanting of seedlings and the period of oviposition by *H. griseola* (Kuwuyama, 1955; Kidokoro *et al.*, 1982).

Insecticides can control infestations of *H. griseola*. These are usually targeted against the adults and newly hatched larvae (*e.g.*, Lange *et al.*, 1953; Kuwuyama, 1955; Grigarick, 1959a; Buryi, 1980), and applications, therefore, require proper timing.

The development of tolerant or resistant varieties of oats (Stavrakis, 1965), spring wheat (Zhu, 1981), and rice (Weber *et al.*, 1988) holds promise for limiting damage by *H. griseola*. Resistance in spring wheat was associated with dense fine hairs on the basal parts of leaves, a thick waxy cuticle, and leaf sheaths that were close to the stem (Zhu, 1981). Mechanisms of resistance in oats and rice were not specified.

Several kinds of natural enemies attack *H. griseola*. Many species of Hymenoptera are known to parasitize *H. griseola*, principally during the larval and pupal stages (for particular species, see Deonier, 1971; Kamijo, 1978; Halfpapp, 1989). In rice, the percentage parasitism in the first generation is usually low, but can approach 80% in later generations (Grigarick, 1959a; Buryi, 1980). A fungus, *Stigmatomyces hydrelliae* Thaxter (Ascomycetes: Laboulbeniales), also attacks *H. griseola* larvae and adults (see Rossi and Cesari Rossi, 1979). Wolf spiders (Araneae: Lycosidae) prey upon adult *H. griseola* in rice fields (Grigarick, 1959a). More research is needed to effectively use parasites and predators to prevent or control outbreaks of this pest, especially in light of the environmental and agronomic constraints with chemical and cultural methods that are discussed above.

Bibliography

The following bibliography contains 195 references on *H. griseola*. Because of the worldwide distribution of this pest, the bibliography collectively contains works written in more than 10 languages. References published in Chinese (C),

Hungarian (H), Japanese (J), Korean (K), or Russian (R), and two in Italian (I), have been listed in English (actual language indicated by abbreviation in parentheses following reference). F = French. Lower case abbreviations are used to indicate the languages in which summaries (abstracts) are written: a = Arabic, e = English, g = German, p = Polish. Articles known to contain synonyms of *H. griseola* are indicated by listing the name(s) in brackets following the reference.

Bibliographical entries were obtained from the following bound abstract sources: Entomology Abstracts (1969, vol. 11 through 1991, vol. 12), Review of Applied Entomology (ser. A, 1955-1990), Science Citation Index (1950-1990). Entries were also obtained from computerized search systems for Agricola (1970-1991), CAB Abstracts (1990-1991), Agris (1986-1988), and Zoological Record (1978-1991). Additional references were derived from citations in some of the reviewed articles.

Literature on *Hydrellia griseola* (Fallén)

- Aldrich, J. M.** 1905. A catalogue of North American Diptera (or two-winged flies). Smithson. Miscell. Collec., vol. 46 (1444). [*H. hypoleuca*, *H. obscuriceps*, *H. scapularis*]
- Ancev, E.** 1974. Neke poznate ili slabo poznate stetne muhe (Diptera) na psenici u Skopskom. Zastite Bilja 25: 271-277.
- Anonymous.** 1987. Rice production. In W.F. Lewis (ed.), California-Arizona Agricultural Digest, Farm Press Publications, Inc., Clarksdale, Mississippi.
- Balachowsky, A., and L. Mesnil.** 1935. Les insectes nuisibles aux plantes cultivees. Tome Premier, Paris.
- Barrion, A. T., and J. A. Litsinger.** 1981. Occurrence of smaller rice leafminer *Hydrellia griseola* (Fallén) (Diptera: Ephydriidae) in the Philippines. Intl. Rice Res. Newsletter 6(6): 19.
- Barrion, A. T., and J. A. Litsinger.** 1986. Ephydrid flies (Diptera: Ephydriidae) of rice in the Philippines. Intl. Rice Res. Newsletter 11(4): 29-30.
- Becker, T.** 1896. Dipterologische Studien IV, Ephydriidae. Berliner Entomol. Zeitschrift 41: 91-267. [*H. griseola*, *H. hypoleuca*, *H. obscuriceps*, *H. scapularis*]
- Becker, T.** 1903. Aegyptische Dipteren Gesammelt und Beschreiben. Mitt. Zool. Sammlung Mus. Natur. Berlin 2(1): 67-195. [*H. griseola*, *H. hypoleuca*, *H. obscuriceps*, *H. scapularis*]
- Becker, T.** 1924. H. Sauter's Formosa Ausbeute, Ephydriidae Diptera). Ent. Mitt. 13: 89-93. [*Psilopa incerta* Becker]
- Becker, T.** 1926. Die fliegen der Palaeartischen Region. Stuttgart, 6(56a). 115 pp.
- Berg, C. O.** 1950. *Hydrellia* (Ephydriidae) and some other acalyprate Diptera reared from *Potamogeton*. Ann. Ent. Soc. America 43: 374-398.
- Bertrand, H.** 1954. Les insectes aquatiques d'Europe (Genres: Larves, Nymphes, Imagos), vol. 1. In P. Lechevalier (ed.), Encyclopedie Entomologique, ser. A, vol. 30. Lechevalier et fils, Paris.
- Bezzi, M.** 1894. I Ditteri del Trentino. Saggio di un elenco delle specie de Ditteri Finora Osservate nel Trentino. Atti Soc. Veneto-Trentino Sci. Nat. (ser. 2) 1: 275-353.
- Bowling, C. C.** 1967. Insect pests of rice in the United States. In Anonymous (ed.), The major insect pests of the rice plant. Johns Hopkins, Baltimore, Md.
- Brischke, C. G. A.** 1883. Breschreibung der Forst-, Garten- und Landwirtschaftlichen Feinde und Frende unter den Insekten. Schriften der Naturforschenden Gesellschaft in Danzig, new ser. 5: 97-125.
- Burgheli, A.** 1959a. New Rumanian species of Dacnusini (Hym., Braconidae) and some ecological observations upon them. Entomol. Monthly Mag. 95: 121-126.
- Burgheli, A.** 1959b. Contributions a la connaissance des Hymenopteres parasitant les jeunes stades d'insectes aquatiques. Bucharest, Universitatea "C. I. Parhon," Analele, Ser. Stuntelor Nat. 22: 143-169.
- Buryi, V.F.** 1980. Pests of rice in the Maritime Province. Zashch. Rast. 11: 32-33. (R)
- Cartwright, W.B., R.A. Blanchard, and C.C. Wilson.** 1933. Notes for 1932 on cereal and forage insects in California. California Dept. Ag. Monthly Bull. 22(2-3): 156-160.
- Casale, A.** 1976. Biological control of *Hydrellia griseola* (Fallén) (Diptera, Ephydriidae) by two new parasites for Italy (Hymenoptera, Braconidae). Riso 25(4): 317-323. (I, e)
- Chakravarthy, A.K.** 1987. Insect pests on main and ratoon rice. Intl. Rice Res. Newsletter 12(4): 35-36.
- Chaillou, C.** 1976. Influence des conditions d'éclairement et de la présence d'un substrat de ponte sur le fonctionnement ovarien des femelles vierges d'*Hydrellia grisiola* Fall. (Diptere Ephydriidae). C. R. Acad. Sci. Paris. 283: 793-796.
- Chaillou, C.** 1978. Influence de la présence du male, de la lumière et du substrat de ponte des œufs chez *Hydrellia griseola* (Diptere, Ephydriidae). Entomol. Exp. Appl. 23(1): 79-90. (e)
- Chiang, H.-C.** 1977. Pest management in the People's Republic of China—monitoring and forecasting insect populations in rice, wheat, cotton and maize. FAO Plant Protec. Bull. 25(1): 1-8.
- Choi, B. S., D. C. Lee, K. R. Yu, and C. K. Park.** 1983. Survey of the rice damage by smaller rice leaf-miner (*Hydrellia griseola* F.) as affected by cultural practices in mid-mountainous area. Res. Rep. Off. Rural Dev. Soil Fert. Crop Protec. Mycology Farm Prod. Util. 25(10): 116-122. (K, e)

- Collins, J. E.** 1928. A new species of *Hydrellia* (Diptera, Ephydriidae) mining the stems of watercress. *Entomol. Monthly Mag.* 64: 128-129.
- Coquillett, D. W.** 1900. Papers from the Harriman Alaska Expedition. IX. Entomological results (3): Diptera. *Proc. Washington Acad. Sci.* 3: 389-464. [*H. scapularis*]
- Coquillett, D. W.** 1904. New North American Diptera. *Proc. Washington Acad. Sci.* 6: 75. [*H. scapularis*]
- Corbetta, G.** 1973. Damages to rice in Italy caused by *Hydrellia griseola* Fall. *Bull. Inf. Rizic France* 145: 11-12. (I, e)
- Cresson, E. T., Jr.** 1918. Costa Rican Diptera collected by Philipp A. Calvert, 1909-1910. Paper III. A report on the Ephydriidae. *American Entomol. Soc. Trans.* 44: 39-68. [*H. hypoleuca*, *H. scapularis*]
- Cresson, E. T., Jr.** 1932. Studies in the dipterous family Ephydriidae. Paper IV. *American Entomol. Soc. Trans.* 58: 1-34.
- Cresson, E. T., Jr.** 1944. Synopses of North American Ephydriidae (Diptera), parts IA and II. *American Entomol. Soc. Trans.* 70: 163-180.
- Cresson, E. T.** 1947. A systematic annotated arrangement of the genera and species of the Neotropical Ephydriidae (Diptera). II. The subfamily Notiphilinae. *American Entomol. Soc. Trans.* 73: 105-124. [*H. hypoleuca*, *H. obscuriceps*]
- Dahl, R. G.** 1959. Studies on Scandinavian Ephydriidae (Diptera Brachycera). *Opuscula Entomol. (Suppl.)* 15: 1-224.
- Dahl, R. G.** 1961. Ephydriidae (Diptera, Brachycera) from Afghanistan. *Entomol. Tidskr.* 82: 87-90.
- Dahl, R. G.** 1964. Revision av Slaktet *Hydrellia* R.-D. (Dipt. Ephydriidae) i Coll. Stenhammar och Zetterstedt. *Opuscula Entomol.* 29: 179-187.
- Dahl, R. G.** 1972. Beitrage zur Kenntnis der Fauna Afghanistans, Ephydriidae, Diptera. *Acta Mus. Moraviae* 56/57: 347-360.
- Dainat, H.** 1971. *Stigmatomyces hydrelliae* Thaxter (Laboulbeniales) parasites de Dipteres Acalipteres dans le sud de la France. *Bull. Soc. Mycol. France* 81(3-4): 337-352.
- Darby, R. E.** 1962. Midges associated with California rice fields, with special reference to their ecology (Diptera: Chironomidae). *Hilgardia* 32: 1-206.
- de Meijere, J. C. H.** 1902. Ueber die Prothorakalstigmen der Dipterenpuppen. *Zoologische Jahrbucher. Abteilungen Ana. Ontog.* 15: 623-692.
- de Meijere, J. C. H.** 1939. Naamlijst van Nederlandsche Diptera. *Tijdschrift voor Entomologie* 82:137-174.
- de Ong, E. R.** 1922. A rice leaf-miner. *J. Econ. Entomol.* 15:432.
- Deonier, D. L.** 1964a. A key to the shore flies of Iowa (Diptera: Ephydriidae). *Iowa St. J. Sci.* 39: 103-126.
- Deonier, D. L.** 1964b. Ecological observations on Iowa shore flies (Diptera, Ephydriidae). *Proc. Iowa Acad. Sci.* 71: 496-510.
- Deonier, D. L.** 1971. A systematic and ecological study of Nearctic *Hydrellia* (Dipt., Ephydriidae). *Smithson. Contrib. Zool.* 68: 1-147.
- Deonier, D. L.** 1972a. Observations on mating, oviposition and food habits of certain shore flies (Diptera: Ephydriidae). *Ohio J. Sci.* 72: 152-166.
- Deonier, D. L.** 1972b. A floating adhesive trap for neustonic insects. *Ann. Ent. Soc. America* 65(1): 269-270.
- Engelhart, V.** 1936. Flies injurious to cereals in the Far East Province. *Plant Proc.* 11: 154.
- Essig, E. O.** 1926. Insects of western North America. Macmillan Co., New York.
- Fallén, C. F.** 1813. Beskrifning Ofver nagra i Sverige funna Vattenflugor (Hydromyzides). *Kongl. Svenska Vet. Akad. Handl.* 34: 240-57.
- Fan, C. D. and H. X. Zhou.** 1983. On the identification of four species of *Hydrellia* flies that are common wheat and rice pests in China (Diptera: Ephydriidae). *Wuyi Sci. J.* 3: 143-144. (C, e)
- Fan, Z., G. Qi, M. Li, R. Yang, and P. Xia.** 1983. Two new species of the genus *Hydrellia* on rice, wheat or barley (Diptera: Ephydriidae). *Entomotax.* 5(1): 7-12. (C, e) [*Hydrellia chinensis*, *H. sinica*]
- Feron, M., and H. Audemard.** 1956. Notes sur *Hydrellia griseola* (Dipt. Ephydriidae), mouche mineuse du riz en France. *Ann. des Epiphyt.* 7(3): 421-430.
- Frey, R.** 1933. En Ammarkningsvard. Insektstandort vid Vusjoki (Ik). *Notulae Entomol.* 13: 81-87.
- Frost, S. W.** 1924. A study of the leafmining Diptera of North America. *Mem. Cornell Univ. Agric. Exp. Station* 78: 1-228.
- Fujisaki, Y.** 1981. Studies on the forecasting method of the occurrence of the smaller rice leaf miner, *Hydrellia griseola* Fallén. I. Survey method. *Annu. Rep. Soc. Plant Protec. N. Japan* 32: 1-2. (J)
- Fujisaki, Y., and K. Abe.** 1985. Overwintering and hatching conditions of the smaller rice leaf miner, *Hydrellia griseola* Fallén. *Annu. Rep. Soc. Plant Protec. N. Japan* 36: 19-21. (J)
- Fujisaki, Y., and M. Ishigaki.** 1981. Studies on the forecasting method of the occurrence of the smaller rice leaf miner, *Hydrellia griseola* Fallén. II. Life table. *Annu. Rep. Soc. Plant Protec. N. Japan* 32: 3-4. (J)
- Fujisaki, Y., M. Ishigaki, and K. Fushimi.** 1980. Studies on the forecasting method of the occurrence of the smaller rice leaf miner, *Hydrellia griseola* Fallén. I. Survey method. *Annu. Rep. Soc. Plant Protec. N. Japan* 31: 110-111. (J)
- Fulmek, L.** 1962. Parasitinsekten det Blattminierer Europas. W. Junk, Haag, Holland.
- Funabasama, K.** 1966. Outbreak factors of rice leaf miner and smaller rice leaf miner in 1965. *Annu. Rep. Soc. Plant Protec. N. Japan* 17: 72. (J)
- Fushimi, K., K. Funabasama, R. Igarashi, and H. Ito.** 1967. Tendency of the occurrence of rice leaf miner at the northern part of Miyagi Prefecture. *Annu. Rep. Soc. Plant Protec. N. Japan* 18: 6. (J)

- Glick, P. A.** 1960. Collecting insects by airplane, with special reference to dispersal of the potato leaf hopper. U.S. Dept. Agric. Tech. Bull. 1222.
- Gobert, E.** 1887. Catalogue des Dipteres de France. Rev. Entomol., Pub. Soc. Francaise Entomol. (Caene), 6(3): 1-87.
- Goetghbuer, M.** 1942. Faunule Dipterologique des Brise-Lames. Bull. Mus. Royale Hist. Nat. Belgique 18(24): 1-10.
- Grensted, L. W.** 1944. Records of Diptera and other insects from Dinas Head, Pembrokeshire. Entomol. Monthly Mag. 80: 201-203.
- Grigarick, A. A.** 1959a. Bionomics of the rice leafminer, *Hydrellia griseola* (Fallén), in California (Diptera: Ephydriidae). Hilgardia 29(1): 1-80.
- Grigarick, A. A.** 1959b. A floating pan trap for insects associated with the water surface. J. Econ. Entomol. 52: 348-349.
- Grigarick, A. A.** 1963a. Rice insect pest control in California. The Rice J. 66(7): 46.
- Grigarick, A. A.** 1963b. Rice plant injury by invertebrate pests. California Agric. 17(8): 6-7.
- Grigarick, A. A.** 1984. General problems with rice invertebrate pests and their control in the United States. Protec. Ecol. 7: 105-114.
- Grigarick, A. A., and W. H. Lange.** 1965. Rice pest investigations. The Rice J. 68(7): 68-70.
- Grigarick, A. A., and R. K. Washino.** 1983. Invertebrates, pp. 49-74. In M.L. Flint (ed.), Integrated pest management for rice. Statewide Integrated Pest Management Project, University of California, Division of Agricultural Science, Davis.
- Grimshaw, P. H.** 1925. The study of flies (Diptera). Naturalist 816: 5-20.
- Gurnberg, K.** 1910. Diptera, Zweifluger. Die Süsswasserfauna Deutschlands de A. Brauer. (2a): 273-82.
- Halfpapp, K. H.** 1989. Rice leafminer *Hydrellia griseola* in Australia. Intl. Rice Res. Newsletter 14(6): 32.
- Hendel, F.** 1926. Blattminenkunde Europas. I. Die Diptereminen. Fritz Wagner, Vienna.
- Hendel, F.** 1933. Neue Acalyptrate Musciden aus der Palaarktischen Region (Dipt.). Deutsche Entomologische Zeitschrift 1933: 39-56.
- Hendrickson, G. O.** 1930. Studies on the insect fauna of Iowa prairies. Iowa St. Coll. J. Sci. 4: 49-179.
- Henning, W.** 1943. Übersicht über die bisber bekannten Metamorphosestadien der Ephydriiden mit Neubeschreibungen nach dem Material der Deutschen Limnologischen Sunda-Expedition. Arb. über Morph. u. Taxonom. Ent. 10(2/3): 105-138.
- Henning, W.** 1958. Die Familien der Diptera Schizophora und Ihre Phyogenetischen Verwandtschaftsbeziehungen. Beiträge zur Entomologie (Berlin) 8: 506-688.
- Hering, E. M.** 1922. Monatliche Anweisungen: Blattminen und Blattminierer. Entomologisches Jahrbuch 31: 36.
- Hering, E. M.** 1925. Minenstudien. VI. Zeitschrift für Morphologie und Ökologie der Tiere 4: 502-539.
- Hering, E. M.** 1926. Die Ökologie der Blattminierenden Insektenlarven. Zool. Bausteine 1: 1-253.
- Hering, E. M.** 1937. Die Blattminen Mittel- und Nord-Europas einschließlich Englands-Bestimmungstabellen aller von Insektenlarven der verschiedenen Ordnungen erzeugten Minen. G. Feller, Neubrandenburg, Germany.
- Hering, E. M.** 1951. Biology of the leaf miners. W. Junk, The Netherlands.
- Hering, E. M.** 1957. Bestimmungstabellen der Blattminen von Europa, vol. 1-3. W. Junk, The Netherlands.
- Hesler, L. S., A. A. Grigarick, M. J. Oraze, and A. T. Palrang.** 1993. Arthropod fauna of conventional and organic rice fields in California. J. Econ. Entomol. 87: 149-158.
- Hirao, J.** 1969. Ecology and control of smaller rice leaf miner attacking transplanted rice. Agr. Hort. 44(6): 677-680.
- Inoue, H.** 1966. Occurrent factor of overwintering type pupa of rice leaf miner. Hokuno 33(4): 17-23. (J)
- Inoue, H.** 1967. Studies on the ecology of the rice leaf miner in Kamikawa Dist. in Hokkaido. Hokuno 34(1): 53-66. (J)
- Isaev, S. I.** 1931. Insects injurious to onions in the Rostov District. Bull. Instit. Controlling Pests Diseases 1: 25-53. (R)
- Ito, H., G. Hino, R. Igarashi, and B. Chiba.** 1970. Occurrence of rice leaf miner in early-planted young plant rice field. Annu. Rep. Soc. Plant Protec. N. Japan 21: 42. (J)
- Iwamoto, S.** 1966. On the seasonal abundance of the smaller rice leaf miner (*Hydrellia griseola* Fallén) in the southern part of the Ibaraki Prefecture. Jpn. J. Appl. Ent. Zool. 10: 110-114. (J)
- Iwamoto, S.** 1976. Some information on the forecast of the smaller rice leaf miner (*Hydrellia griseola* Fallén) occurrence. Bull. Ibataki Agric. Exp. Stn. 17: 97-104. (J)
- Jensen, A., and A. F. Nielsen.** 1961. Kornbadfluen (*Hydrellia griseola*), et skadedyr på korn og graes. Ugeskr. Landm., no. 50, repr. 3.
- Jensen, D. D.** 1954. Proceedings, 232nd Meeting, Pacific Coast Entomological Society. Pan-Pacific Entomol. 30(1): 84-87.
- Johannsen, O. A.** 1935. Aquatic Diptera, Part II. Orthorrhapha-Brachycera and Cyclorrhapha. Mem. Cornell Univ. Agr. Exp. Sta. 177.
- Johnson, C. W.** 1925. Fauna of New England, 15. List of the Diptera or two-winged flies. Occ. Pap. Boston Soc. Nat. Hist. 7: 1-326. [*H. hypoleuca*, *H. scapularis*]
- Jones, B. J.** 1906. Catalogue of Ephydriidae, with bibliography and descriptions of new species. Univ. California Pubs. Tech. Bull. 1(2): 153-198. [*H. hypoleuca*, *H. obscuriceps*, *H. scapularis*]
- Joshi, R. C., E. A. Heinrichs, and S. Uthamasamy.** 1986. A bibliography on *Hydrellia* spp. attacking

- rice. FAO Asia Pac. Pl. Protec. Comm. Quart. News. 29(2): 29-44.
- Kamijo, K.** 1978. Chalcidoid parasites (Hymenoptera) of Agromyzidae in Japan, with description of a new species. *Kontyu* 46(3): 455-469. (J)
- Karnecka, H.** 1985. A review of species of *Hydrellia* (Rob.-Desv.) found in Bohemia (Diptera, Ephydriidae). *Acta Entomol. Bohemoslovaca* 82(4): 315-319. (r)
- Kas'yanov, A. I.** 1967. Diptera as pests of rice. *Zashch. Rast.* 9: 33-34. (R)
- Kas'yanov, A. I.** 1971. Effectiveness of metathione for controlling *Hydrellia griseola* in rice crops. *Khim. Sel'skom. Khoz.* 9(8): 31-32. (R)
- Kas'yanov, A. I.** 1976. Pests of rice crops. *Zashch. Rast.* 8: 28-31. (R)
- Kato, S.** 1955. Morphology of the smaller rice leaf-miner. In S. Kuwayama (ed.), *Investigation on Hydrellia griseola Fallén*, the smaller rice leaf-miner. Spec. Rep. no.3, Soc. Pl. Prot. N. Japan (J)
- Kawall, J. H.** 1867. *Miscellanea Entomologica*. Stettiner Entomol. Zeitschrift 28: 120-121.
- Kidokoro, T., and M. Maeda.** 1985. Effect of larval density on the leaf consumption and mortality of the smaller rice leaf miner, *Hydrellia griseola* Fallén. *Annu. Rep. Soc. Pl. Protec. N. Japan* 36: 17-18. (J)
- Kidokoro, T., Y. Fujisaki, and T. Takano.** 1982. Transplanting time of rice as a factor in the occurrence of outbreaks of the smaller leaf miner, *Hydrellia griseola* (Fallén). *Jpn. J. Appl. Ent. Zool.* 26(4): 306-308. (J, e)
- Kihata, M.** 1962. Investigation on forecast of rice leaf miner in Kamikawa district in Hokkaido. *Annu. Rep. Soc. Plant Protec. N. Japan* 13: 95-96. (J)
- Klimanova, N. K.** 1971. Leaf miners on rice. *Zashch. Rast.* 16(4): 33-34. (R)
- Kloet, G. S., and W.D. Hinks.** 1945. A checklist of British insects. Stockport, England. [*Hydropota griseola*]
- Kortring, A.** 1931. Ueber die Entwicklung der grauen Gerstenminierfliege (*Hydrellia griseola* Fall.) bei verschiedener Ernährung. *Zeitschrift Pfl. Krankh.* 41: 321-333.
- Kowarz, F.** 1894. Verzeichnis der Insekten Bohmen's II. Fliegen. Catalogus Insectorum Faunae Bohemicae. Physiokratisch Gesellschaft, Prague.
- Koyama, T., M. Kikuchi, and J. Hirao.** 1968. On the injuries caused by insect pests and the development of the rice stem borer in the rice plants sown directly on a submerged field. *Bull. Tohoku Natl. Agric. Exp. Stn.* 36: 43-51. (J)
- Kreiter, E. A.** 1927. Contribution to the biology of the barley fly. *Rep. Bur. Appl. Ent.* 3(1): 92-98. (E, g)
- Kreiter, E. A.** 1928. Dipterous larvae occurring on graminaceous plants in Leningrad. *Izu. Otd. Prikl. Eng.* 3(2): 251-264. (R?)
- Kuwayama, S.** 1955. Investigations on *Hydrellia griseola* Fallén, the smaller rice leaf-miner. *Spec. Rep. No. 3*. The Society of Plant Protection of North Japan, Omagari, Japan. (J)
- Lameere, A.** 1907. *Manuel de la Faune de Belgique*, vol. 3. H. Lambertin, Brussels.
- Lampa, S.** 1906. Berattelse till Kungl. Landbruksstyrelsen Angande Verksamheten vid Statens Entomologiska Anstalt under Ar 1905. *Entomol. Tidskr.* 27: 17-64.
- Lange, W. H., and A. A. Grigarick.** 1970. Insect pests, pp. 3-16. In W.H. Lange and M.D. Miller (eds.), *Insects and other pests of rice*. California Agric. Exp. Stn. Circ. 555.
- Lange, W. H., Jr., K. H. Ingrebretsen, and L.D. Davis.** 1953. Rice leaf miner—severe attack controlled by water management, insecticide application. *California Agric.* 7(8): 8-9.
- Leberre, J.R., and G. Stravrakis.** 1965. Etude physiologique du développement d'un Diptère Ephydriidae, *Hydrellia griseola* Fall.: nature chimique des substances digérées par la larve et coefficients d'utilisation digestive. *Compt. Rend.* 260: 4074.
- Leberre, J.R., and G. Stavrakis.** 1966. Etude physiologique du développement d'un Diptère Ephydriidae *Hydrellia griseola* Fall.: évolution biochimique chez la larve et la pupa. *C. R. Acad. Sci. ser. D* 263: 136.
- Leberre, J.R., H. Chevin, and J. P. Moreau** 1961. Longevité, Fécondité et Nutrition de Deux Diptères: *Oscinella frit* L., *Hydrellia griseola* Fall. *Rev. Zoo. Agric.* (Talence) 60: 151-161.
- Lilljeborg, W.** 1861. En flugas harjningar a kornfalten i ostra skane, blekinge och soda delarna af kalmar lan under sommaren 1860. *Tidskr. for landtmanns- och kommunalekonomin*, Uppsala, pp. 205-215.
- Linnaniemi, W. M.** 1913. Zur Kenntnis der Blattminierer, speziell derjenigen Finlands I. *Acta Soc. Fauna Flora Fennica* 37(4): 1-138.
- Loew, H.** 1860. Die Europaeischen Ephydrinidae und die Bisher in Schlesien Beobachteten Arten Derselben. Neue Beiträge zur Kenntniß der Dipteren. Part 7, pp. 1-46. Mittler und Sohn, Berlin.
- Loew, H.** 1862. Monographs of the Diptera of North America. Part I. Smithson. Inst., pp. 129-172.
- Macquart, J.** 1835. *Histoire naturelle des insectes, Dipteres*. Tome deuxième. Pourrat Frères, Paris.
- Majewski, T.** 1972. Rare and new Laboulbeniales from Poland. V. *Acta Myc.* 8(2): 229-237.
- Malloch, J. R.** 1915. Some additional records of Chironomidae for Illinois and notes on other Illinois Diptera. *Bull. Illinois St. Lab. Nat. Hist.* 11: 305-364. [*H. scapularis*]
- Manandhar, D., and A. A. Grigarick.** 1983. Effect of rice leafminer feeding on early growth of the rice plant. *J. Econ. Entomol.* 76: 1022-1027.
- Marchal, P.** 1903. Sur la biologie des *Hydrellia* (Dipt.) degats exercés sur le Cresson par l'*Hydrellia ranunculi* Hal. *Bull. Soc. Entomol. France* 1903: 236-237.

- Mesnil, L.** 1931. Recoltes de Muscides nuisables aux cereales. Rev. Path. Veg. Entomol. Agric. 18(8-9): 226-274.
- Moragues y de Manzanos, F.** 1894. Insectos de Mallorca. Anal. Soc. Espania Hist. Nat. 23: 73-87.
- Morley, C., and E. A. Atmore.** 1915. The Diptera of Norfolk and Suffolk. Norfolk Norwich Nat. Soc. Trans. 10: 162.
- Murai, S.** 1967. A consideration on the seasonal prevalence of adults of *Hydrellia griseola* Fallén in Shonai District, Yamagata Prefecture. J. Yamagata Agr. For. Soc. 24: 29-32. (J, e)
- Nepveu, P., and J. D'Aguilar.** 1951. Un nouveau ravageur du riz in Camargue. Acad. d'Ag. France-Proces. Seance du 4 Juil, pp. 463-465.
- Novoded, M. F.** 1971. Pests on rice. Zashch. Rast. 16(5):12-13. (R)
- Obst, A.** 1980. Control of *Haplodiplosis equestris* and *Hydrellia griseola*. Bayerishces Landwirtschaftliches Wochenblatt. 170(18): 18.
- Osten Sacken, C. R.** 1878. Catalogue of the described Diptera of North America. Smithson. Miscell. Coll. vol. 16, no. 270. [*H. hypoleuca*, *H. obscuriceps*, *H. scapularis*]
- Paik, W. H.** 1967. Insect pests of rice in Korea. In Anonymous (ed.), The major insect pests of the rice plant. Johns Hopkins, Baltimore, Maryland.
- Parker, H. L., P. A. Berry, and A. Silveira Guido.** 1951-1952. Host-parasite and parasite-host lists of insects reared in South American parasite laboratory during the period 1940-1946. Rev. Assoc. Ing. Agron. 92: 15-112.
- Robineau-Desvoidy, A. J. B.** 1830. Essai sur les Myodaires. Mem. Acad. Roy. Sci. Institut. France 2: 1-813.
- Rossi, W., and M. G. Cesari Rossi.** 1979. Su alcune specie di *Stigmatomyces* (Ascomycetes, Laboulbeniales) parassite di Ditteri Italiani. Bull. Museo Civico Storia Naturale Venezia 30: 13-17. (e)
- Satoh, Y., S. Dobashi, and A. Itomi.** 1983. Catch of smaller rice leafminer, *Hydrellia griseola* Fallén by tow net trap and light trap (in Japan). Annu. Rep. Soc. Pl. Protec. N. Japan 34: 9-10. (J)
- Scheiring, J. F., and B.A. Foote.** 1973. Habitat distribution of the shore flies of northeastern Ohio (Diptera: Ephydriidae). Ohio J. Sci. 73: 152-166.
- Scherney, F.** 1966. Zur Biologie der an Gerste minierenden Agromyzidenart, *Agromyza nigrella* Rondani (Dipt.). Z. PflKrank. PflPath. PflSchutz. 73: 141-146.
- Schiner, J. R.** 1864. Fauna Austriaca. Die Fliegen (Diptera). vol. 2, part 1. Carl Gerold's Sohn, Vienna.
- Schmidt, H. A.** 1962. Schaden an Mais durch Dipterenzarven. NachrBl. dtsch. PflSchDienst. 16(3): 58-59.
- Seguy, E.** 1934. Faune de France, 28. Dipteres (Brachycerous). Lechevalier, Paris.
- Seguy, E.** 1950. La biologie des Dipteres. Encyclopedie Entomologique, serie A, 26. Lechevalier, Paris.
- Shurovnekov, B. G.** 1985. Thelytoky and sex ratio disruption in insect populations. Zashch. Rast. 1: 30-31. (R)
- Slosson, A. T.** 1902. Additional list of insects taken in alpine regions of Mt. Washington. Entomol. News 13: 319-321. [*H. obscuriceps*, *H. scapularis*]
- Smith, C. M., J. L. Bagent, S. D. Linscombe, and J. F. Robinson.** 1986. Insect pests of rice in Louisiana. Louisiana Agric. Exp. Stn. Bull. 774. Louisiana State Agricultural Center, Baton Rouge.
- Sorauer, P., and L. Reh.** 1913. Handbuch der Pflanzenkrankheiten. Die Tierischen Feinde, vol. 3. Paul Parey, Berlin.
- Starostin, S. P., V. I. Kondratenko, V. I. Tanskii, P. Tkachenko, and S. L. Tyuterev.** 1987. Principal pests, diseases and weeds of rice and their control. Zashch. Rast. 11: 26-27.
- Stavrakis, G.** 1965. Analyse dans les conditions du laboratoire d'un exemple de resistance varietale d'avoine vis à vis d'un Diptera *Hydrellia griseola* (Ephydriidae), p. 536. In P. Freeman (ed.), Proc. 12th Intl. Congr. Entomol.
- Stavrakis, G.** 1966. Quelques aspects des relations d'*Hydrellia griseola* Fall. et ses plantes-hotes. These Doct. Ing., Fac. Sci. Univ. Paris XI, Orsay.
- Stein, F.** 1867. Eine der Gerste Schädliche Fliege. Berliner Entomologische Zeitschrift 11: 395-397.
- Strobl, P. G.** 1900. Spanische Deapteren. VIII. Wiener Entomol. Zeitschrift 19: 1-10.
- Strobl, P. G.** 1904. Neue Beiträge zur Dipterenfauna der Balkanhalbinsel. Wissenschaftliche Mitteilungen Bosnien Herzegowina 9: 519-581.
- Sturtevant, A. H.** 1926. The seminal receptacles and accessory glands of the Diptera, with special reference to Acalyptratae. J. New York Entomol. Soc. 34: 1-21. [*H. hypoleuca*]
- Szilvassy, L., and I. Petrasovits.** 1960. On the damage in rice fields caused by *H. griseola* in 1959. Novenytermesles 9(2): 171-174. (H, e, r)
- Szilvassy, L., and A. Szito.** 1982. Data on the biology and damage of the rice pests *Cricotopus bicinctus* Meigen and *Hydrellia griseola* Fallén. Novenyvedelem 18(12): 542-550. (H, e, g, r)
- Szito, A.** 1976. Data relating to the overwintering of *Hydrellia griseola* Fallén. Folia Entomol. Hungaria 29(1): 103-106. (H, e)
- Szito, A.** 1982. Role of temperature and precipitation in the trend of damage caused by larvae of the rice mosquito (*Cricotopus bicinctus* Meigen, Diptera: Chironomidae) and the rice fly (*Hydrellia griseola* Fallén, Diptera: Ephydriidae). Novenyvedelem 20(10): 439-444. (H, e, g, r)
- Szito, A.** 1985. Short-term prediction of damage caused by larvae of the rice mosquito (*Cricotopus bicinctus* Meigen) and the rice fly (*Hydrellia griseola* Fallén), pp. 157-167. In Risztermeszzi Tudomanyos Tanacskozas Szarvas, AGROINFORM, Budapest. (H)

- Takano, T., and Y. Fujisaki.** 1985. Emergence time of the smaller rice leaf miner adult, *Hydrellia griseola* Fallén. Annu. Rep. Soc. Pl. Protec. N. Japan 36: 14-16. (J)
- Takano, T., and K. Funabasama.** 1990. Ovipositing prevalence of smaller rice leaf miner, *Hydrellia griseola* Fallén, on rice plants transplanted early. Annu. Rep. Soc. Pl. Protec. N. Japan 41: 114-116. (J)
- Timon-David, J.** 1961. Contribution à l'étude d'écologie et du Peuplement des Iles de la Région Marseillaise. Colloques Inter. Centre Natl. Recherches Sci. XCIV. Le Peuplement des Iles Méditerranéennes et le Problème de L'Insularité. II. Liste des Diptères Captures à Ratonneau-Pomegues. pp. 228-230.
- Tomioka, T.** 1962. The effect of humidity on the development of eggs of the smaller leaf miner, *Hydrellia griseola* Fallén (Diptera: Ephydriidae). Bull. Hokkaido Pref. Agr. Exp. Sta. 9: 8-17. (J, e)
- Tomioka, T.** 1983. Ecological note on two species of *Hydrellia* (*H. tomiokai* and *H. griseola*, on rice in Japan). Annu. Rep. Soc. Pl. Protec. N. Japan 36: 13-16. (J)
- Tomioka, T., S. Okuyama, and S. Hachiya.** 1983. Two species of *Hydrellia* (*H. tomiokai* and *H. griseola*) infesting rice leaf in Hokkaido (Japan). Annu. Rep. Soc. Pl. Protec. N. Japan 36: 11-12. (J)
- Thier, R. W. and B. A. Foote.** 1980. Biology of mudshore Ephydriidae: Diptera. Proc. Ent. Soc. Washington 82(4): 517-536.
- Thygesen, T.** 1967. Kornbladfluen overvintring. Maanedsovers PlSygd. 429: 24.
- Tsacas, L.** 1959. Contribution à la connaissance des Diptères de Grèce, 1. Bull. Soc. Entomol. France 64: 123-130. F
- van der Wulp, F. M. and J. C. H. de Meijere.** 1898. Nieuwe Naamlijst van Nederlandsche Diptera. Tijds. Entomol. 41: 1-143.
- Vayssiére, P.** 1933. La mouche de Cresson (*Hydrellia nasturtii* Collin [Dipt. Ephydriidae]) et son parasite (*Ademon decrescens* Nees [Hym. Braconidae]). Bull. Soc. Entomol. France 38: 86-87.
- von Frauenfeld, G. R.** 1869. Ueber einige Pflanzenverwuster des Jahres 1869. Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 19: 601-604.
- von Oettingen, H.** 1935. Die wichtigsten grasbewohnenden Fliegenlarven Norddeutschlands. Nachr. über Schadlingsbekämpf 10(2): 62-70. (e)
- Wahlgren, E.** 1947. *Hydrellia*-Arten als Blattminierer in *Stratiotes aloides* L. (Dipt.). Opuscula Entomol. 12: 78-79.
- Walker, E.** 1856. Insecta Britannica. Diptera, vol. 3. John Edward Taylor, London.
- Washburn, F. L.** 1906. Additional Minnesota Diptera. Eleventh Annu. Rep. St. Entomol. Minnesota 1906: 79-82. [*H. hypoleuca*]
- Way, M. O., F. T. Turner, and J. K. Clark.** 1983. The rice leafminer, *Hydrellia griseola* (Fallén), a potential pest of rice in Texas. Southwest. Entomol. 8: 186-189.
- Weber, G., J. Gibbons, and K. Eichelkraut.** 1988. El manejo de *Hydrellia* spp. Arroz en las Americas 9(2): 11-14.
- Wilke, S.** 1924. Die graue gerstenminier Fliege, *Hydrellia griseola* Fall. (Diptera: Ephydriidae). Deut. Ent. Zeitschrift 6: 172-179.
- Willcocks, F. C.** 1925. The insect and related pests of Egypt. Part 2. Cairo Sultanic Ag. Soc., Tech Sec., 418 pp.
- Wirth, W. W.** 1965. Ephydriidae, pp. 734-759. In A. Stone, C. W. Sabrosky, W. W. Wirth, R. H. Foote, and J. R. Coulson (eds.), Catalog of the Diptera of America North of Mexico. U.S. Govt. Printing Office, Washington, D.C.
- Wirth, W. W., and A. Stone.** 1956. Aquatic Diptera, pp. 372-482. In R. L. Usinger (ed.), Aquatic insects of California. Univ. California Press, Berkeley.
- Zangheri, S.** 1956. Un dittero minatore del riso nel basso Ferrarese (*Hydrellia griseola* Fallén, Dipt. Ephydriidae). Boll. Soc. Ent. Italia 86(1-2): 12-16.
- Zatwarnicki, T.** 1988. Materials to the knowledge of the genus *Hydrellia* Robineau-Desvoidy (Ephydriidae: Diptera). Polskie Pismo Entomol. 58(3): 587-634. (p)
- Zhu, J. L.** 1981. Preliminary observations on the resistance of varieties of spring wheat to *Hydrellia griseola* (Fallén). Kunchong Zhishi 18: 213-214. (C)

Additional literature cited

- Altieri, M., J. Davis, and K. Burroughs.** 1983. Some agroecological and socioeconomic features of organic farming in California. Biol. Agric. Hort. 1: 97-107.
- Bock, I. R.** 1990. The Australian species of *Hydrellia* Robineau-Desvoidy (Diptera: Ephydriidae). Invert. Taxon. 3: 965-993.
- Buckingham, G. R., E. A. Okrah, and M. C. Thomas.** 1989. Laboratory host range tests with *Hydrellia pakistanae* (Diptera: Ephydriidae), an agent for biological control of *Hydrilla verticillata* (Hydrocharitaceae). Environ. Entomol. 18: 164-171.
- Buckingham, G. R., E. A. Okrah, and M. Christian-Meier.** 1991. Laboratory biology and host range of *Hydrellia balciunasi* (Diptera: Ephydriidae). Entomophaga 36: 575-586
- Deeming, J. C.** 1977. A new species of *Hydrellia* Robineau-Desvoidy (Diptera: Ephydriidae) attacking rice in southern Nigeria. Bull. Ent. Res. 67(2): 337-341.
- Isa, A. L.** 1989. Integrated pest management in Egypt, pp. 213-218. In Rice farming systems: new directions. IRRI, Los Banos, Philippines.
- Ismail, I. I., A. L. Isa, and E.-M. Farrag.** 1979. Notes on the biology of the rice leaf miner, *Hydrellia*

prosternalis Deeming (Diptera: Ephydidae). Agricultural Res. Rev., Cairo 57(1): 87-94. (a)
Williams, J. F., S. R. Roberts, J. E. Hill, S. C. Scardaci, and G. Tibbits. 1990. Managing water for weed control in rice. California Agric. 44(5): 7-10.

Acknowledgments

The following individuals assisted in developing the bibliography: Sheldon Cheney and Susan Wilzer

(USDA, National Agricultural Library); staff at the Hilton M. Briggs Library, South Dakota State University; Albert A. Grigarick and Drew Palrang (University of California at Davis) and David Tolk (USDA-ARS, Brookings, SD). I thank Gary R. Buckingham, D. L. Deonier, Albert A. Grigarick, Drew Palrang, and two anonymous reviewers for helpful suggestions on earlier versions of the manuscript.