

## POPULATION DYNAMICS OF PLANT PARASITIC NEMATODES AND THEIR RELATIONSHIPS WITH SOIL PHYSICO-CHEMICAL PARAMETERS DURING THE FALLOW PERIOD AFTER SUGARBEET AND LEADING INTO TOMATO CULTIVATION

P. Barbosa<sup>1</sup>, H. Adão<sup>2</sup> and M. Mota<sup>1</sup>

<sup>1</sup>NemaLab/ICAM, Universidade de Évora, 7002-554 Évora, Portugal

<sup>2</sup>IMAR-CIC, Universidade de Évora, 7002-554 Évora, Portugal

**Summary.** The main objective of this research was the study of the soil nematode community, and in particular plant parasitic nematodes (PPN), from a field located in Portugal's southern region, used for sugarbeet production. The study was performed from February to July 2003, covering part of the fallow period previous to tomato cultivation, the alternative crop in the rotation. The end of the fallow period in March and the soil preparation period in May were marked by a significant reduction in the numbers of PPN, whereas their numbers increased on the following tomato crop. The genus *Helicotylenchus* stood out as the most representative group, forming 90% of all PPN counted each month. The genus *Heterodera* was relatively abundant in the months following the previous sugarbeet crop, and numbers of the genus *Meloidogyne* increased during the tomato crop. The correlations between these groups and environmental parameters show that, apart from the direct influence of the host, pH, organic matter, temperature and soil moisture significantly influenced nematode abundance and community composition.

**Key words:** Crop rotation, plant parasitic nematodes, soil characteristics, sugarbeet.

Since its first cultivation and improvement at the beginning of 19<sup>th</sup> century, sugarbeet (*Beta vulgaris vulgaris* L.) has gradually achieved prominence as a crop of economic importance both worldwide and at national level. Presently, it is potentially one of the most important crops in terms of economical viability and agricultural vitality for many countries. In Portugal, it is an alternative to traditional crops, with rising importance from both agronomic and economic points of view and following the rules of the new European common agriculture policy (M. Espadinha, DAI, personal communication).

Nematodes are among the most important pests that affect this crop, being responsible for 10% of the losses in the total amount of sugar produced (Steele, 1984; Sasser, 1989). Several species of nematodes feed on the roots or occasionally in the shoot tissues; young seedlings are very susceptible and may suffer irreversible damage (Cooke, 1993). *Heterodera* and *Meloidogyne* are the main genera responsible for the loss of productivity in sugarbeet fields (Steele, 1984; Shurtleff and Averre, 2000; Starr *et al.*, 2002).

In general terms, research regarding relationships between phytoparasitic nematodes and soil variables from cultivated fields pays particular attention to the gender or species that is responsible for the greater amount of damage. However, and although the host plays a predominant role in the distribution and presence/absence of the individuals of this group, agricultural practices

result in alterations of the physico-chemical soil parameters. These alterations condition the composition and distribution of the nematode community associated with the crop under consideration (Yeates, 1999).

To study the changes in population densities of phytoparasitic nematodes associated with sugarbeet cultivation and their relationships with various soil factors, a survey was conducted during the three final months of the fallow period after sugarbeet cultivation (February to April) and the first three months of the tomato cropping period (May to July), in southern Portugal.

### MATERIALS AND METHODS

*Study area.* The study area is located in Portugal's southern region, with the region's typically temperate climate. The average air temperature was low during February 2003 (9.1 °C), gradually increasing after March (13.6 °C) until reaching unusually high values in June (22.6 °C) and July (22.9 °C). Precipitation was high in April (72 mm), abundant during February (60 mm) and March (51 mm), but very little from May (8 mm) to July (0 mm).

The field was a cultivated area of 1 ha of primary alluvium soil (AL type). Systematic sugarbeet cultivation had started about 28 years previously, with a rotation length of five years. Agricultural practices before and during the study are summarized in Table I.

*Soil sampling for nematodes.* Sampling was performed monthly from February to July 2003. The field was di-

---

<sup>1</sup> Contact author e-mail: pedronematology@gmail.com

**Table I.** Summary of the agricultural practices applied before and during the study.

| Date                     | Land management  | Remarks   |
|--------------------------|--|---|
| September 2002           | Spring sugarbeet harvest<br>Coarse harrowing.  |   |
| February 2003            | Fallow   | Soil drenched<br>Weeds<br>Residues of previous culture (leaves and old sugarbeet roots) |
| March 2003<br>April 2003 | Fallow   | Weeds<br>Residues of previous culture   |
| May 2003                 | Soil preparation<br>Addition of fertiliser (N, P, K)<br>Tomato seeding<br>Insecticide treatment<br>Drip irrigation |   |
| June 2003<br>July 2003   | Irrigation   |   |

vided into five equal areas. Nematode community analyses were made on the five areas, under the assumption that all plots were uniform regarding soil physico-chemical characteristics. For each area, a “composite sample” was obtained from fifteen sample points, collected from two plant rows, at a 0-20 cm depth. The collected soil was mixed thoroughly and eight sub-samples of 200 g each were used for analysis.

Nematodes were extracted by decanting and sieving (25, 60 and 325 mesh sieves), followed by centrifugal flotation (in sugar solution) (Caveness and Jensen, 1955). After extraction, nematodes were preserved in 4% formalin solution (Barker, 1985). Phytoparasitic nematodes were identified to genus level according to Brzeski (1998) and Siddiqi (2000) under an inverted microscope (Olympus® CK30). Nematode abundance was expressed as number of individuals per 200 g of soil.

*Soil sampling for physico-chemical parameter determination.* Soil used for determining physico-chemical parameters was collected on each of the six sampling dates from three points along the centre of the terrain. Samples were placed individually in plastic bags. Sub-samples from each bag were taken to determine: pH (H<sub>2</sub>O); nitrate content (N-NO<sub>3</sub><sup>-</sup>); extractable phosphorus (P<sub>2</sub>O<sub>5</sub>) and potassium (K<sub>2</sub>O); exchangeable cations, calcium (Ca<sup>+</sup>) and magnesium (Mg<sup>+</sup>); organic matter; soil moisture. A thermometer measured soil temperature (20 cm depth), after allowing twenty minutes for equilibration. Analysis of these parameters followed the usually employed methods (Midgley and Torrance, 1978; Thomas, 1982).

*Statistical analyses.* The data collected were analysed in three stages.

i) To compare the monthly mean abundance for each phytoparasitic genus, the Kruskal-Wallis non-parametric test was applied, followed by multiple comparisons

with the Dwass-Steel-Critchlow-Fligner method (Critchlow and Fligner, 1991).

ii) To determine temporal patterns in plant-parasitic nematodes and in soil parameters, two separate principal component analyses (PCA) with data transformation were employed (Manly, 1986; Legendre and Legendre, 1998).

iii) The relationships between plant parasitic nematodes and soil physico-chemical parameters were investigated using corresponded canonical analyses (CCA) (Manly, 1986; Legendre and Legendre, 1998).

To avoid statistical misinterpretations, the last two stages were repeated without the most abundant genus identified in the first analysis.

These multivariate analyses were carried out using the STATISTICA 6.0® and STATSDIRECT 2.4.1® statistical packages.

## RESULTS

*Soil physico-chemical parameters.* Results of the physico-chemical characterization are summarized in Fig. 1A and Fig. 1B. Soil temperature increased progressively throughout the study, from spring to summer (10 °C-27 °C). Soil moisture content was high from February (24.5%) to April (20.8%) due to the drenching of the soil by the heavy precipitation, and increased after May (12.8%) due to irrigation. The pH was maximum at the beginning of the study (7.6), decreased to a minimum in May (6.2), and increased after soil preparation for the tomato crop. The organic matter content (1.4-1.9%) remained the same, especially from May to July. Calcium (8.44-15.72 meq/100 g) and magnesium (0.87-2.09 meq/100 g) concentrations suffered mild oscillations, with the lowest value in June. Phosphorus (95-181 ppm) and potassium (90-169 ppm) concentrations decreased progressively from February to April (minimum





Fig. 1. A. Soil physico-chemical parameters from February to July 2003 ( $\pm$ standard deviation). Different scales.

value), increased significantly during May (maximum value) and decreased again until July. Nitrogen concentrations (12-166 ppm) varied in a similar way but with a highest value in June rather than May.

Results of the principal component analysis on soil parameters are shown in Fig. 2. The first two axes represent 75% of the total variation. The first factor (horizontal axis) with 55% of the variability mainly opposes phosphorus and potassium to soil moisture and calcium. The second factor with 20% of the variability mainly isolates magnesium. The remaining parameters occupy intermediate positions.


In the correlation circle, aggregation occurring between components with higher concentrations/values from February until April is evident. The same situation occurs with those parameters that increased from May to July (soil temperature, nitrogen, phosphorus and potassium – right side). In the first case, the group formed by soil moisture and pH is due to increased values after May. The grouping in the second case illustrates the significant boost in potassium and phosphorus concentrations after soil fertilisation in May, the steady rise of soil temperature as the season progressed, and the isolation of nitrogen is a result of the extremely high concentra-



**Fig. 1. B.** Soil physico-chemical parameters from February to July 2003 ( $\pm$ standard deviation). Different scales.

tion in June, also due to fertilisation (Fig. 2).


The seasonal changes in soil parameters were not a steady progression (Fig. 3). Results in April were closer to those in February/March due to an overall lowering of soil parameters (apart from soil temperature) and were very similar between months (i.e. the months during which the land was fallow). The same situation occurred in June when some parameters decrease in value (chemicals depleted due to cultivation of the crop) and tended towards the values observed in the initial



**Fig. 2.** Results of PCA on soil parameters from February to July 2003 at a 0-20 cm depth. **A.** Eigenvalues; **B.** Correlation circle F1-F2.

months. The high concentrations of a few of the parameters (mainly phosphorus, nitrogen and potassium) during field preparation and the sowing of a new crop, clearly isolated May from June.

The passage of time emphasized the conjunction of distinct soil parameters in different months. Soil physico-chemical characteristics were very dissimilar during the assay, with important contributions from different parameters in each month. The overall contribution was similar during the four months located in the vicinity of the horizontal axis of Fig. 3. In the two remaining months, some parameters had distinctly low values (soil moisture and pH in May) or high values (nitrogen in June) so that these months stand apart from the horizontal axis.



**Fig. 3.** Results of PCA on soil parameters from February to July 2003 at a 0-20 cm depth.

*Phytoparasitic nematodes.* During this study, ten genera from nine families of phytoparasitic nematodes were identified: *Helicotylenchus* (Hoplolaimidae, 96% of the total community), *Tylenchus* (Tylenchidae, 2%), *Tylenchorhynchus*, *Merlinius* (Telotylenchidae, 1%), *Pratylenchus* (Pratylenchidae, 1%), *Heterodera* (Heteroderidae, <1%), *Meloidogyne* (Meloidogynidae, <1%), *Trichodorus* (Trichodoridae, <1%), *Gracilacus* (Paratylenchidae, <1%) and *Criconemella* (Criconematidae, <1%). Even with the large discrepancies between genera relative abundance, results of analyses obtained with and without inclusion of the most abundant were very similar. The lower percentages for some genera are a consequence of the high numbers of *Helicotylenchus* and also because their densities varied month by month.

As shown in Fig. 4A, the patterns of monthly variation of the numbers of *Helicotylenchus* (Mean  $\pm$  SD) (344.1 individuals/200 g  $\pm$  115.4), *Tylenchus* (5.6  $\pm$  2.62), *Tylenchorhynchus* (4  $\pm$  3.79), *Pratylenchus* (3  $\pm$  2.03) and *Merlinius* (1.6  $\pm$  1.69) were similar. The mean numbers of nematodes fell from February to May, when the minimum abundance was reached; in June and July the numbers of individuals increased to the maximum abundances observed. The Kruskal-Wallis statistical test showed significant differences between months for each of the genera; *Helicotylenchus* (T = 215.1), *Tylenchus* (T = 102.5), *Tylenchorhynchus* (T = 140.3), *Merlinius* (T = 64.6), *Pratylenchus* (T = 942); P < 0.0001.

The densities of *Heterodera* (1.8  $\pm$  1.99) and *Meloidogyne* (0.6  $\pm$  1.23) followed opposite progressions (Figs 4A and 4B). For the former, the mean abundance decreased with time, whilst the presence of the latter increased after May. This is reflected in the significant differences between months; for *Heterodera*, T = 159.6 and, for *Meloidogyne*, T = 98.8; P < 0.0001. Values obtained for these two genera refer exclusively to juveniles.

For *Criconemella* (0.1  $\pm$  0.38), *Gracilacus* (0.3  $\pm$  0.64) and *Trichodorus* (0.5  $\pm$  0.96), the densities decreased in the first three months and they were not found in the last three (Fig. 4B). The Kruskal-Wallis test revealed significant differences between months for *Criconemella* (T = 22.3), *Gracilacus* (T = 46.4) and *Trichodorus* (T = 73.3); P < 0.0001.

The greatest number of nematodes occurred in July (556 individuals/200 g), followed by February (447/200 g) and May (195/200 g). The number of genera present was greatest (10) in February/March and least (6) in May. *Helicotylenchus* was clearly the most abundant genus in the entire study, whilst *Criconemella* was the least abundant.

The results of the principal component analysis on phytoparasitic nematodes demonstrate that the first two axes represented 97% of the total variation (Fig. 5). The first axis, with 54% of the variability, opposed *Tylenchus*, *Meloidogyne* and *Tylenchorhynchus*, with the higher positive coordinates, against *Heterodera* and *Gracilacus*. The second axis, with 43% of the variability, separated *Helicotylenchus* from *Merlinius* and *Praty-*

*lenchus*. The remaining genera occupied intermediate positions.

In a subsequent analysis from which the figures for *Helicotylenchus* were omitted (not reported), no significant differences were found. The first two axes represented 94% of the total variation. The greatest difference occurred with *Pratylenchus*, which shifted from its former group and became closer to *Tylenchus*. The remaining genera maintained essentially the same positions, with *Tylenchus* as the most abundant.

There seems to be a division between individuals with a final abundance lower than the starting one (first factor, left side) and those with a higher abundance at the end of the study. The first situation is represented by a group that includes *Criconemella*, *Gracilacus* and *Trichodorus*, all absent during the three final months, plus *Heterodera* alone, which was present in the final three months. The second situation is represented by a group formed by the most abundant genera, *Tylenchus* and *Helicotylenchus*, and a group formed by *Merlinius*, *Pratylenchus*, *Tylenchorhynchus* and *Meloidogyne*, whose abundances were very small or even null in some months.

The correlation matrix of the monthly factors for the phytoparasitic genera (Fig. 6) showed a distinct seasonal effect between February and March (winter), April (beginning of spring), May (end of spring), and June and July (summer). Factor 1 (the horizontal axis) consisted mainly of time variation, showing the seasonal traits of the community, as months are distributed along the axis. Factor 2 separated the months with lower abundance from those with a greater abundance; the end of spring illustrates the first situation and summer the second.

Thus, the communities from each seasonal group show significant differences, and mainly between winter and summer. The beginning and end of spring had intermediate positions, although the first was close to the winter observations and the second was distinct from all others.


*Relationships between phytoparasitic nematodes and soil physico-chemical parameters.* The eigenvalues of the canonical correspondence analysis between phytoparasites and soil parameters show that the first factor takes into account 56% of the total variation and that the second factor was responsible for 42% (Fig. 7). This analysis does not show significant differences when compared with the PCA of soil parameters, with phosphorus, nitrogen, potassium and soil temperature opposed to the other parameters; the relations between individual soil parameters are similar to previously.

When *Helicotylenchus* was not considered in the analysis, differences in the canonical analysis were almost null; this is to be expected since the PCA on phytoparasites did not reveal major differences. Apart from an inversion along the horizontal axis and a slight difference in the CCA total variation, all the relationships remained the same.


For phytoparasites, there was an evident separation

between those with a downward trend in population density in the final months (*Criconebella*, *Gracilacus*, *Trichodorus* and *Heterodera*) and those with increasing population densities in the same period, with a tighter grouping of the genera in the latter group.


The correlation matrix between phytoparasitic nematodes and soil physico-chemical parameters is summarized in Fig. 8. *Heterodera* showed the greatest number of correlations (five) with the nine parameters, and was highly negatively correlated with soil temperature. *Tri-*



**Fig. 4. A.** Mean number of nematodes/200 g ( $\pm$  standard deviation) for phytoparasitic genera from February to July 2003 at a 0-20 cm depth. Different scales. Kruskal-Wallis analysis by genera – different letters indicate significant differences ( $P < 0.05$ ) between months.



**Fig. 4. B.** Mean number of nematodes/200 g ( $\pm$  standard deviation) for phytoparasitic genera from February to July 2003 at a 0-20 cm depth. Different scales. Kruskal-Wallis analysis by genera – different letters indicate significant differences ( $P < 0.05$ ) between months.




**Fig. 5.** Results of PCA on phytoparasitic nematodes from February to July 2003 at a 0-20 cm depth. **A.** Eigenvalues; **B.** Correlation circle F1-F2.

*chodorus*, *Criconemella* and *Gracilacus* had four significant correlations, all with the same four parameters, namely soil temperature, soil moisture, pH and organic matter, and all strongly positively associated with organic matter. The remaining genera had only single positive relationships: *Tylenchus*, *Merlinius*, *Helicotylenchus* and *Pratylenchus* with pH, *Tylenchorhynchus* and *Meloidogyne* with soil temperature. Overall, the positive correlations were in the majority.


Correlations with soil moisture, pH and organic matter were always positive; on the other hand the only one with potassium was negative. The greatest number of correlations were with pH, a total of eight. There were no correlations with phosphorus, nitrogen, calcium or magnesium.

## DISCUSSION

*The phytoparasitic community.* The existence of a high density of phytoparasitic nematodes at the beginning of this study allowed us to postulate their strong presence




**Fig. 6.** Results of PCA on phytoparasitic nematodes from February to July 2003 at a 0-20 cm depth.



**Fig. 7.** Results of CCA between phytoparasitic nematodes and soil physico-chemical parameters from February to July 2003 at a 0-20 cm depth.

during the previous cropping period and/or the development of a series of adaptations that counteract the absence of a “living” host (Kable and Mai, 1968).

However, the later absence of a developing crop and the network of fungi and soil bacteria usually associated with roots contributed strongly to a lowering of phytoparasite abundance during the fallow period (Weaver *et al.*, 1995), with special relevance in the case of *Heterodera*, usually the main genus responsible for the loss of productivity in sugarbeet fields. Other factors may have contributed to the decline in the phytoparasite population, such as the poor adaptation of these organ-



**Fig. 8.** Correlation matrix between phytoparasitic nematodes and soil physico-chemical parameters ( $\alpha = 0.05$ ) from February to July 2003 at a 0-20 cm depth. The sizes of the circles (positive correlations) and the squares (negative correlations) are proportional to the degree of correlation (Pearson).

isms to the natural plant succession that occurs as plant cover moves from crop to various weeds (Donfack *et al.*, 1995); any re-adaptation to a different species of plant host may not be automatic, and in some cases may require a transitory period (Baujard and Martiny, 1995).

After a fallow period, *Criconebella*, *Gracilacus* and *Trichodorus* were no longer found and the abundance of *Heterodera* reached very low values. An opposite situation occurred with *Meloidogyne*; despite a low incidence from February until April, its presence increased after crop introduction. As values obtained for *Heterodera* and for *Meloidogyne* refer exclusively to juveniles, their population densities and potential effects are underestimated.

In this field, *Helicotylenchus* was undoubtedly the dominant phytoparasite with monthly abundances of more than 95% of the total community, clearly higher than the values obtained in other crops (corn, soybean) by Cares and Huang (1991), Mattos (1999) and Goulart *et al.* (2003), where they formed around 40% of the total.

For many years, the farming techniques in this area were based on traditional cropping, followed by a short fallow period and a new crop. The genus *Helicotylenchus* is very adaptable, even when the same crop is grown repeatedly, being able to survive and reproduce in frequently changing environments, even with different food sources (Kandji *et al.*, 2001; Gomes *et al.*, 2003; Goulart *et al.*, 2003). Being an ectoparasite, a damaging effect on the crop only occurs in the presence of a massive number of individuals; in our case, it is certainly possible that *Helicotylenchus* might have had a harmful effect on the new crop.

*Phytoparasitic nematodes and soil parameters.* In sum-



mary, we have demonstrated that there are well defined relationships between phytoparasitic nematodes and soil parameters. Nitrogen, phosphorus and potassium (traditionally incorporated in the fertilization) concentrations were very close in the PCA. When comparing results between PCA on phytoparasites and the correlations established, those for *Trichodorus*, *Criconebella*, *Gracilacus* and *Heterodera* were the most similar, and the negative effect of soil temperature strengthened the separation of these four genera from the others, particularly *Tylenchorhynchus* and *Meloidogyne*.

During this study, clear monthly changes occurred and these reflected the agricultural circumstances. The community structures for February/March and for June/July, this last with greater numbers of individuals, reflect what has occurred in terms of agronomical practices: fallow and cultivation. The other two months were in an intermediate position, being distinguished by the lesser number of individuals in April, given the reduction in soil nutrients, and in May due to land management practices.

*Heterodera*, *Gracilacus*, *Trichodorus* and *Criconebella*, present in greatest numbers at the beginning of the year, displayed an evident affinity with the amount of organic matter, but also with pH and soil moisture; in fact they were the only phytoparasites correlated with the last parameter mentioned. Higher soil temperatures and, to a minor extent, a greater concentration of nitrogen, phosphorus and potassium (from fertilization) after May were clearly adverse for them.

The increasing abundances of *Tylenchorhynchus* and *Meloidogyne* in June and July was influenced by the increasing soil temperature and to a lesser degree by a higher concentration of phosphorus, nitrogen and potassium, which counteracted the antagonistic effect of the pH. *Helicotylenchus*, *Merlinius*, *Tylenchus* and *Pratylenchus* showed a certain preference for high pH values and, to a minor extent, soil moisture. No parameter affected them to a significant degree, so their seasonality was more dependent on the host than on the soil.

Crop absence during part of the study may hypothetically create correlations between abundance of some genera and pH (Norton, 1991; Cadet *et al.*, 1994; Korthals *et al.*, 1996), soil moisture (McSorley, 1997; Yeates, 1998) and organic matter (Norton *et al.*, 1971). The positive correlations shown by the great majority of phytoparasitic nematodes with the increasing soil moisture and temperature may have been diluted due to the introduction of a tomato crop in April. Soil moisture was a limiting factor only in April/May since before that period the soil was drenched by rains and afterwards was permanently irrigated. Both situations of greater soil moisture match the highest nematode densities recorded (Ferris *et al.*, 1971; Wallace, 1973; Rickard and Barker, 1982).

*Impact of land management on nematodes.* The decline in the number of individuals during the fallow period (February/April) was at the time that a crop able to support phytoparasitic nematodes was absent, and there

were no land management practices that would provide soil enrichment in terms of nutrients or water. The following seems to have a noticeable effect on a few genera and to temporarily diminish the abundance of others. According to Yeates (1991), Neher (2001) and Gomes *et al.* (2003), some nematodes may have a high sensitivity to a specific type of environmental disturbance, declining to undetectable levels within a short period of time.

Seasonal changes with time, characterized by lower soil moisture and increased temperatures, were other factors that may have contributed to the reduced numbers of nematodes at the end of fallow (Gupta and Yeates, 1997). Soil moisture is not the only parameter to determine the densities of phytoparasitic nematodes found, and others, such as the presence of weeds after harvest, residues from the previous crop or nutrients accumulated (mainly nitrogen) by those plants, help to maintain high phytoparasite populations in the periods between crops (Anwar *et al.*, 1992; Wasilewska, 1997).

In May, land management in preparation for the tomato crop, with the subsequent removal of old roots/weeds and introduction of fertilizers and protective chemicals, drastically altered the situation, hastening the decline in nematode abundance (Freckman and Caswell, 1985; Bongers, 1990; Yeates and Hughes, 1990).

In the following months, changes induced by the presence of the crop and the constant irrigation benefited those nematodes that use tomato roots as a food resource (Barker and Campbell, 1981; Ettema and Bongers, 1993). Thus the situation in which phytoparasite densities were decreasing was reversed and their densities increased again.

## ACKNOWLEDGEMENTS

The authors gratefully acknowledge Mr Paulo Vieira for useful comments and corrections on the manuscript, Dr Russel Alpizar Jara for his great support and advice in the statistical treatments, Mrs Francisca Figo for all the technical assistance in the laboratory, and Mr Manuel Espadinha, Technical Director of DAI – Sociedade para o Desenvolvimento Agro-Industrial S.A., for all the information and support provided.

## LITERATURE CITED

- Anwar S.A., Rauf C.A. and Gorski S.D., 1992. Weeds as alternate hosts of phytonematodes. *Afro-Asian Journal of Nematology*, 2: 41-47.
- Baujard P. and Martiny B., 1995. Ecology and pathogenicity of the Hoplolaimidae from the Sahelian Zone of West Africa. 2. Laboratory studies on *Scutellonema cavenessi* Sher, 1964. *Fundamental and Applied Nematology*, 18: 335-345.
- Barker K.R., 1985. Sampling nematode communities. Pp. 3-17. In: An Advanced Treatise on Meloidogyne, vol. II (Barker K.R., Carter C.C. and Sasser J.N., eds). North Carolina State University Graphics, Raleigh, USA.

- Barker K.R. and Campbell C.I., 1981. Sampling nematode populations. Pp. 451-474. *In: Plant Parasitic Nematodes*, vol. III (Zuckerman B.M. and Rohde R.A., eds). Academic Press, New York, USA.
- Bongers T., 1990. The maturity index: an ecological measure of environmental disturbance based on nematode species composition. *Oecologia*, 83: 14-19.
- Brzeski M.W., 1998. *Nematodes of Tylenchina in Poland and temperate Europe*. Muzeum i Instytut Zoologii Polska Akademia Nauk, Warszawa, Poland, 397 pp.
- Cadet P., Thioulouse J. and Albrecht A., 1994. Relationships between ferrisil properties and the structure of plant parasitic nematode communities on sugarcane in Martinique (French West Indies). *Acta Ecologica*, 15: 767-780.
- Cares J.H. and Huang S.P., 1991. Nematode fauna in natural and cultivated cerrados of Central Brasil. *Fitopatologia Brasileira*, 16: 199-209.
- Caveness F.E. and Jensen H.J., 1955. Modification of the centrifugal-flotation technique for isolation and concentration of nematodes and their eggs from soil and plant tissue. *Proceedings of the Helminthological Society of Washington*, 22: 87-89.
- Critchlow D.E. and Fligner M.A., 1991. On distribution-free multiple comparisons in the one-way analysis of variance. *Communications of Statistical Theory and Methods*, 20: 127-139.
- Cooke D., 1993. Nematode parasites of sugarbeet. Pp. 133-170. *In: Plant Parasitic Nematodes in Temperate Agriculture* (Evans K., Trudgill D.L. and Webster J.M., eds). CAB International, Wallingford, UK.
- Donfack P., Floret C. and Pontanier R., 1995. Secondary succession in abandoned fields of dry tropical northern Cameroon. *Journal of Vegetal Science*, 6: 499-508.
- Ettema C.H. and Bongers T., 1993. Characterization of nematode colonization and succession in disturbed soil using the maturity index. *Biology and Fertility of Soil*, 16: 79-85.
- Ferris V.R., Ferris J.M., Bernard R.L. and Probst A.H., 1971. Community structure of plant-parasitic nematodes in soybean rotation fields. *Journal of Nematology*, 3: 339.
- Freckman D.W. and Caswell E.P., 1985. The ecology of nematodes in agroecosystems. *Annual Review of Phytopathology*, 23: 275-296.
- Gomes G.S., Huang S.P. and Cares J.E., 2003. Nematode community, trophic structure and population fluctuation in soybean fields. *Fitopatologia Brasileira*, 28: 258-266.
- Goulart A.C., Monteiro A.R. and Ferraz L.B., 2003. Comunidades de nematóides em cerrado com vegetação original preservada ou substituída por culturas. 2. Diversidade taxionômica. *Nematologia Brasileira*, 27: 123-128.
- Gupta V.R. and Yeates G.W., 1997. Soil microfauna as bioindicators of soil health. Pp. 201-234. *In: Biological Indicators of Soil Health* (Pankhurst C.E., Doube B.M. and Gupta V.R., eds). CAB International, Wallingford, UK.
- Kable P.F. and Mai W.F., 1968. Overwintering of *Pratylenchus penetrans* in a sandy loam and clay loam soil at Ithaca, New York. *Nematologica*, 14: 150.
- Kandji S.T., Ogol C.O. and Albrecht A., 2001. Diversity of plant-parasitic nematodes and their relationships with some soil physico-chemical characteristics in improved fallows in western Kenya. *Applied Soil Ecology*, 18: 143-157.
- Korthals G.W., Alexiev D.A., Lexmond T.M., Kammenga J.E. and Bongers T., 1996. Long-term effects of copper and pH on the nematode community in an agroecosystem. *Environmental Toxicology and Chemistry*, 15: 979-985.
- Legendre P. and Legendre L., 1998. *Numerical Ecology*, second English edition. Elsevier Science B.V., Amsterdam, The Netherlands, 853 pp.
- McSorley R., 1997. Relationships of crop and rainfall to soil nematode community structure in perennial agroecosystems. *Applied Soil Ecology*, 6: 147-159.
- Manly B.F., 1986. *Multivariate Statistical Methods – a primer*. Chapman & Hall, Ltd., London, UK, 159 pp.
- Mattos J.A., 1999. *Caracterização das comunidades de nematóides em oito sistemas de uso da terra nos cerrados de Brasil Central*. Universidade de Brasília, Brasília, Brazil, 113 pp.
- Midgley D. and Torrance K., 1978. *Potentiometric Water Analysis*. John Wiley and Sons, Inc, New York, USA, 409 pp.
- Neher D.A., 2001. Role of nematodes in soil health and their use as indicators. *Journal of Nematology*, 33: 161-168.
- Norton D.C., 1991. Biology and ecology of nematodes. Pp. 46-71. *In: Manual of Agricultural Nematology* (Nickle W.R., ed.). Marcel Dekker, New York, USA.
- Norton D.C., Frederick L.R., Ponchillia P.E. and Nyhan J.W., 1971. Correlations of nematodes and soil properties in soybean fields. *Journal of Nematology*, 3: 154.
- Rickard D.A. and Barker K.R., 1982. Nematode assays and advisory services. Pp 8-20. *In: Nematology in the Southern Region of the United States* (Riggs R.D., ed.). Arkansas Agricultural Experiment Station, Southern Cooperative Services Bulletin 276, Fayetteville, USA.
- Sasser J.N., 1989. *Plant-parasitic Nematodes: the farmer's hidden enemy*. Department of Plant Pathology and Consortium of International Crop Protection, North Carolina State University, Raleigh, USA, 115 pp.
- Shurtleff M.C. and Averre III, C.W., 2000. *Diagnosing plant disease caused by nematodes*. APS Press, St. Paul, Minnesota, USA, 187 pp.
- Siddiqi M.R., 2000. *Tylenchida parasites of plants and insects - second edition*. CABI Publishing, Wallingford, UK, 833 pp.
- Starr J.L., Bridge J. and Cook R., 2002. Resistance to plant-parasitic nematodes: history, current use and future potential. Pp. 1-22. *In: Plant Resistance to Parasitic Nematodes* (Starr J.L., Cook R. and Bridge J., eds.). CABI Publishing, New York, USA.
- Steele A.E., 1984. Nematode parasites of sugar beet. Pp. 507-545. *In: Plant and Insect Nematodes* (Nickle W.R., ed.). Marcel Dekker, Inc., New York, USA.
- Thomas G.W., 1982. Exchangeable cations. Pp. 159-165. *In: Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties - Agronomy Monograph n.º 9*, second edition (Page A.L., Miller R.H. and Keeney D.R., eds). Soil Science Society of America, Inc., Madison, USA.
- Wallace H.R., 1973. *Nematode Ecology and Plant Disease*. Edward Arnold (Publishers) Limited, London, UK, 228 pp.
- Wasilewska L., 1997. Soil invertebrates as bioindicators, with special reference to soil-inhabiting nematodes. *Russian Journal of Nematology*, 5: 113-126.
- Weaver D.B., Rodriguez-Kabana R. and Carden E.L., 1995. Comparison of crop rotation and fallow for management of *Heterodera glycines* and *Meloidogyne* spp. in soybean. *Journal of Nematology* 27(S): 585-591.
- Yeates G.W., 1991. Impact of historical changes in land use on the soil fauna. *New Zealand Journal of Ecology*, 15: 99-106.
- Yeates G.W., 1998. Soil nematode assemblages: regulators of ecosystem productivity. *Phytoparasitica*, 26: 97-100.
- Yeates G.W., 1999. Effects of plants on nematode community structure. *Annual Review of Phytopathology*, 37: 127-149.
- Yeates G.W. and Hughes K.A., 1990. Effect of three tillage regimes on plant and soil nematodes in an oats/maize rotation. *Pedobiologia*, 34: 379-387.